

LAND, FRESH-WATER, AND MARINE CRUSTACEA.

BY THOMAS SCOTT, F.L.S., Mem. Soc. Zool. de France, F.R.Ph.S.E.,
Corr. Mem. Geol. and Nat. Hist. Soes. of Glasgow, etc.

ORDER PODOPHTHALMA.

AUTHORITIES AND REFERENCES.

- (F. S. G.)=The Statistics of the work of the Fishery Board, S.S. "Garland."
- (H.)=Prof. J. R. Henderson "The Higher Crustacea of the Clyde" (*N. H. Soc. Glasg. Trans.*, vol. i. (N. S.) p. 315).
- (M.)=MS. records of Sir John Murray's S.Y. "Medusa."
- (A. M. N.)=Rev. Canon A. M. Norman, F.R.S., "A Revision of the British species of Schizopoda" (*Ann. and Mag. N. H.*, 1892).
- (R.)=Records of the late Dr. Robertson, Millport.
- (C.)=Dr. W. Y. Calman, Dundee, "On the genus *Pandalus*."
- (S.)=Thomas Scott, F.L.S.

NOMENCLATURE AND CLASSIFICATION FOLLOWED.

Rev. T. R. R. Stebbing, "History of the Crustacea," 1893 (Intern. Scient. Ser.).

Shall. w.=shallow water; l. m. w.=low water of ordinary or spring tides;
d. w.=deep water; bet. t. m.=between tide marks; h. w.=high water.
[Insufficiently attested or doubtful species are enclosed within square brackets.]

Sub-Order Brachyura.

Fam. Cancridae.

Cancer pagurus, Lin.—C. on rocky shores (H.).

Xantho hydrophilus (*Herbst.*).—One captured at mouth of estuary 15/6/1899

(F. S. G.). Fishery Board's Coll., Aberdeen.

Pilumnus hirtellus, (Lin.).—Portlooy, Cumbrae, under stones at l. w. (R.).

Apparently r.

Fam. Portunidae.

Carcinus maenas (*Penn.*).—C. bet. t. m., and in shall. w. inshore; sometimes in 10-20 f. upwards.

Portunus puber (*Lin.*).—Cumbrae; Lamlash B.; L. Fyne (H.): Shallow Bays, Barmore (S.).

P. depurator (*Lin.*).—The most c. *Portunus* in the Clyde (H.).

P. corrugatus (*Penn.*).—One off Tan Buoy, 8 f. (H.).

P. arcuatus, *Leach*.—F. of Clyde, not r.

P. pusillus, *Leach*.—Mod. c.; hard ground, down to 30 f. (H.): nr. Fort Matilda, Greenock (S.).

P. marmoreus, Leach.—Several localities (H.).

Portumnus latipes (Penn.)—Washed ashore, Irvine sands (R.).

Fam. Corystidae.

Corystes cassivelaunus, (Penn.)—Kames B., Cumbrae, sand below l. w. m. (R.) : small male at mouth of estuary, 15/6/99 (F. S. G.).

Atelecyclus septendentatus (Mont.)—One from Alland Bank, Millport B. (R.).

Fam. Ocypodidae.

Gonoplax rhomboides (Lin.)—Ayrshire coast (G. & M.) : nr. Ailsa Craig (trawled, F. S. G.) : shore, Ardeer (J. Smith).

Fam. Pinnotheridae.

Pinnotheres pisum (Lin.)—Not unc. in *M. modiolus*, occasionally in *C. edule* (H.).

Fam. Inachidae.

Macropodia rostratus (Lin.)—Not unc. in lower Clyde.

M. longirostris (Fabr.)—Mod. c., gen. dist. ; more c. than *M. rostratus*. [*Achaeus cranchii*, Leach.—One, off Cumbrae by Mr. Kennedy (R.) : U. L. Fyne, 10 f. (M.)].

Inachus dorsettensis (Penn.)—C. and gen. dist.

I. dorynchus, Leach.—Gen. dist., less c. than *I. dorsettensis*.

I. leptochirius, Leach.—F. of Clyde (A. M. N.) : bet. Cumbrae and Bute, 20 f., r. (R.).

Fam. Maiidae.

[*Maia squinado* (Herb.)—Off Dunoon, 1878 (Fleming) : off Ardrossan (Maj. Martin) : W. of Bute (Grieve) (See H.).]

Hyas araneus (Lin.)—C. and gen. dist., esp. inshore.

H. coarctatus, Leach.—C. and gen. dist., esp. in d. w.

Fam. Parthenopidae.

Eurynome aspera (Penn.)—Not c. ; 10-60 f. ; usually hard ground (H.) : off Fort Matilda 10 f. (S.).

Fam. Leucosiidae.

Ebalia tuberosa (Penn.)—Gen. dist. ; usually hard ground, 5-50 f. (H.) : off Fort Matilda (S.).

E. tumifacta (Mont.)—Lamlash B. (G. ; H.) : Alland Bank, Millport B. (R.) : Kilbrennan Sd., 15 f. (Brook).

E. cranchii, Leach.—Lamlash B. (L. : M.) : off Lion Rock Cumbrae (Alex. Patience) !

Sub-Order Macrura.

Fam. Lithodidae.

Lithodes maia (Lin.)—D. w. L. Fyne, and Clyde ; gen. dist. ; somewhat r. (H. ; F. S. G.).

Fam. Paguridae.

Eupagurus bernhardus (Lin.)—Abund., more inshore than in d. w.

E. prideauxi (Leach).—Mod. c., more in d. w. than inshore.

E. cuanensis (Thompson).—Skelmorlie Bank ; off Portincross ; Cumbrae ; Lamlash B. ; not very c. (H.).

E. pubescens (Kr.).—Mod. c. ; hard ground ; usually enveloped by *Hymenocidon tuberosa* (H.).

Anapagurus hyndmanni (Thompson).—F. of Clyde, 10-15 f. (A. M. N.) : off Cumbrae (R.) : E. L. Tarbert (S.).

A. laevis (*Thomps.*).—Gen. dist., 10-105 f. (H.): mod. freq. in Clyde and L. Fyne.

A. chiroacanthus (*Lillj.*).—Skelmorlie Bank; off Portincross; Tan Buoy; Lamlash B. (H.): E. L. Tarbert (S.).

Fam. Porcellanidae.

Porcellana platycheles (*Penn.*).—Under stones, nr. Cumbrae; Lamlash B. (H.): Lunderston B. (S.).

P. longicornis (*Lin.*).—C. and gen. dist., more freq. inshore; usually hard ground.

Fam. Galatheidae.

Galathea squamifera, *Leach*.—Under stones l. w., Cumbrae; Lamlash B.; off Fairlie (H.): U. L. Fyne (M.): E. L. Tarbert; Lunderston B.

G. nexa, *Embleton*.—L. Fyne, 20-40 f.; Tan Buoy, 7-8 f.; Lamlash B.; Skelmorlie Bank; etc. (H.); U. L. Fyne (M.).

G. dispersa, *Bate*.—Mod. c. and gen. dist.; usually hard ground.

G. intermedia, *Lillj.*.—C. hard ground, 10-20 f. (H.): L. Fyne 10-20, f. (M.).

G. strigosa, *Fabr.*.—Under stones at l. w. spring tides, Cumbrae; young dredged off Roseneath Pt., 15 f., hard ground (H.): Strachur B. (S.).

Munida rugosa (*Fabr.*).—Gen. dist. and mod. c., usually d. w.

Fam. Thalassinidae.

Jaxea nocturna, *Nardo*.—Adult fragments twice (1899) in fish stomachs nr. Ailsa Craig (F. G. Pearcey): Young at times freq. in bottom tow-net (F. S. G.).

Fam. Thaumastochelidae.

Calocaris macandreae, *Bell*.—L. Fyne, 40-105 f.; off Tarbert; off Wemyss B., 40 f.; off Brodick B., 93 f. (H.).

Fam. Palinuridae.

Palinurus vulgaris, *Latr.*.—Off Campbeltown; L. Fyne (H.): r.

Fam. Nephropsidae.

Nephrops norvegicus (*Lin.*).—C., usually d. w.

Astacus gammarus (*Lin.*).—Rocky shores nr. mouth of Firth, c., (H.): E. L. Tarbert, etc.

Fam. Crangonidae.

Crangon vulgaris (*Lin.*).—Mod. c.; shall. w. inshore, on sandy, weedy bottom.

C. allmanni, *Kinahan*.—C. and gen. dist., mostly in d. w.
Egeon sculptus (*Bell.*).—Lamlash B., 5 f., two (A. M. N.); off Muggie Pt., L. Cumbrae, 20 f., one (H.).

Pontophilus spinosus, *Leach*.—Not c.; many localities as far up as Cloch L.H., usually soft ground 20-60 f. (H.).

Cheraphilus nanus (*Kr.*).—Off Cumbrae (R.): Ballantrae Bank (F. S. G.).

C. echinulentus, *M. Sars*.—Off Skate I., L. Fyne, 105 f., one (H.): not very r. at mouth of estuary (F. S. G.).

C. neglectus, *G. O. Sars*.—Occasionally in E. L. Tarbert and vicinity, 5-20 f. (S.).

Fam. Nikidae.

Nika edulis, *Risso*.—One off Skate I., 105 f.; one off Brodick 93 f. (H.): occasionally in shrimp-trawl in seaward part of estuary (F. S. G.).

Fam. Hippolytidae.

Caridion gordoni (*Bate*).—Lamlash B. (A. M. N.): not c. throughout Firth; d. w. (H.).

Hippolyte varians, Leach.—C. ; rock pools l. w., and lam. zone (H.) : c. E. L. Tarbert.

H. prideauxiana, Leach.—Shrimp trawl nr. mouth of estuary, 11/96 (F. S. G.). [*H. fascigera*, Gosse.—E. L. Tarbert, amongst *zostera*, not r. Now considered to be a form of *H. varians*.]

Spirontocaris spinus (Sowerby), var. *Hippolyte securifrons*, Norm.—Is mod. c. Typical *S. spinus* at mouth of estuary, but r. (vide Stebbing).

S. cranchii (Leach).—L. Fyne ; off Cumbrae ; off Craigmore ; Hawk's Neb, Bute (H.) : E. L. Tarbert (S.).

S. pusiola (Kr.).—Off Cumbrae (R.) : L. Goil ; L. Long (H.).

S. gaimardii (M. Edw.).—Gen. dist., prob. not very r. (H.).

Fam. Pandalidae.

Pandalus montagui, Leach.—C. and gen. dist. ; largest and most highly coloured from d. w.

P. bonnieri, Caulfield.—L. Fyne ; L. Long (Calman).

P. propinquus, G. O. Sars.—L. L. Fyne ; L. Long, 40 f. (Calman).

Pandalina brevirostris, (Rathke).—C. and gen. dist. (H.).

Fam. Palaemonidae.

Leander squilla (Lin.).—C. ; rock pools (H.).

Palaemonetes varians (Leach).—Cumbrae at l. w. (R.) : Inverkip (S.).

Fam. Pasiphaeidae.

Pasiphaea sivado (Risso).—Off Skate I. 105 f. (H.) : off Strachur 70 f. (M.) : mid-water tow-net, L. L. Fyne ; not r. at mouth of estuary (F. S. G.).

Sub-Order Schizopoda.

ARRANGEMENT AND NOMENCLATURE FOLLOWED.

Rev. Canon A. M. Norman, F.R.S., D.C.L., "Revision of the Brit. Species of Schizopoda" (*Ann. and Mag. N. H.*, Ser. 6, vol. ix., pp. 454-464, and Ser. 6, vol. x., pp. 143-166 : and pp. 241-263, 1892).

Fam. Euphausiidae.

Rhoda raschii (M. Sars) (= *Boreophausia raschii*).—Gen. dist., sometimes c. esp. in L. Fyne.

R. inermis (Kr.).—Clyde (Brook ; Hoyle). Off Sanda I. 20-27 f. (F. S. G., 15/12/98).

Nyctiphanes norvegica (M. Sars).—Gen. dist., usually more c. in L. Fyne than elsewhere.

Thysanoessa neglecta (Kr.).—In bottom tow-net bet. Ailsa Craig and Sanda I. 20 f. (F. S. G., 16/1/99).

Fam. Mysidae.

Siriella clausii, G. O. Sars.—E. L. Tarbert ; L. Gilp ; r. (F. S. G.).

S. brooki, Norman.—E. L. Tarbert, 1886 (S.).

S. armata (M. Edw.).—Cumbrae, in surface tow-net, r. (R.) : E. L. Tarbert ; L. Gilp (S.).

Gastrosaccus spinifer (Goës).—F. of Clyde (R.) ; nr. Sanda I. ; nr. Ailsa Craig 20-30 f. ; S. of Davaar I. 7-10 f. (F. S. G.).

Anchialus agilis, G. O. Sars.—Nr. Sanda I. 20-27 f. (F. S. G., 15/12/98).

Heteromysis formosa, S. J. Smith.—Dredged, Tarbert Bank, L. L. Fyne

(F. S. G., 1899) : r.

Erythrops elegans, G. O. Sars.—Gen. dist. from U. L. Fyne to seaward limits of estuary ; rarely captured except in bottom tow-net.

E. serratus, G. O. Sars.—Dist. somewhat similar to *E. elegans*.

Mysidopsis didelphys (*Norman*).—Off Cloch L.H. 43 f. (H.): dist. from U. L. Fyne to mouth of estuary.
M. gibbosa, *G. O. Sars*.—Gen. dist. but not very c.
M. angusta, *G. O. Sars*.—L. Fyne and nr. Ailsa Craig; not c. (F. S. G.)
Leptomyysis gracilis, *G. O. Sars*.—U. L. Fyne; nr. Ailsa Craig, etc.
L. lingvura, *G. O. Sars*.—Gen. dist., but not very c.
Hemimysis lamornae (*Couch*).—Port Glasgow; L. Goil (R.): E. L. Tarbert; nr. Ailsa Craig; Davaar I. (F. S. G.).
Praunus flexuosus (*Müller*).—More or less c.; usually more so inshore than in d. w.
P. inermis (*Rathke*).—Cumbrae in Lam. zone (H.): E. L. Tarbert, etc. (F. S. G.).
[P neglectus (*G. O. Sars*).—Mouth of L. Fyne 60 f. (H.): never met with by (S.).]
Schistomyysis spiritus, *Norman*.—E. of Sanda I., surface tow-net 24/1/99 (F. S. G.).
S. ornata, (*G. O. Sars*).—Head of L. Fyne; bet. Ailsa Craig and Sanda I.; scarce.
S. arenosa (*G. O. Sars*).—E. L. Tarbert; shallow water, not r.; Ettrick B., Bute (1887) (S., A. M. N.).
Neomysis vulgaris (*J. V. Thompson*).—Dhu Loch, nr. Inveraray, not r.; in stomach of 15 spined stickleback nr. head of L. Fyne (S.). (Bell suggests this = *Mysis integer* recorded by Leach from L. Ranza.)

ORDER EDRIOPHTHALMA.

Sub-Order Sympoda.¹

ARRANGEMENT AND NOMENCLATURE.

Prof. G. O. Sars, *Crustacea of Norway*, vol. iii.

AUTHORITIES.

- (R.)=Dr. Robertson, F.G.S., "Thirteen Cumacea from the Firth of Clyde" (*Nat. Hist. Soc. Glas. Trans.*, vol. iii. (N. S.), p. 47, 1889).
(S.)=Thomas Scott, F.L.S., "Scottish Cumaceans" (*Ann. Scot. N. H.*, Oct. 1900). Rev. T. R. R. Stebbing, *A History of the Crustacea*.
(A. M. N.)=Rev. Canon A. M. Norman, D.C.L., F.R.S., "On a Crangon, some Schizopoda and Cumacea, new or rare in the British Seas" (4th Ann. Rep. Fish. Bd. for Scot., 1886).

[¹ See Stebbing on change of "Cuma" and "Cumacea" to "Bodotria," and "Sympoda," in Memoir *On Crustacea brought by Dr. Willey from the South Seas* (Camb. Univ. Press) Dec. 1900.]

Fam. Bodotriidae.

- Bodotria arenosa**, *Goodsir*.—Blackwaterfoot, Arran, 4-5 f. (R.).
B. scorpioides (*Mont.*).—Dredged, Ballantrae Bank, 6/9/98 (S.).
B. pulchella, *G. O. Sars*.—Dredged nr. Sanda I. (S.).
Iphinoë trispinosa (*Goodsir*).—Plentiful Kames B., Millport; L. Ranza (R.); U. L. Fyne; Kilbrennan Sd. (S.).
I. serrata, *Norman*.—Kames B., Bute, 1887; off Ballantrae, 1899 (S.).
I. tenella, *G. O. Sars*.—Clyde (Stebbing).
Cumopsis edwardsii (*Spence Bate*) (= *C. goodsiri*), *V. Ben.*.—Shore pools Millport (S.).
C. longipes (*Dohrn*).—Shore pools Millport (R. ; S.)

Fam. Vaunthompsoniidae.

- Vaunthompsonia cristata**, *Sp. Bate*.—E. L. Tarbert 1886; Ballantrae Bank, 15-20 f. 1899 (S.).

Fam. Lampropidae.

Lamprops fasciata, *G. O. Sars*.—W. of Helensburgh Pier (R.): E. L. Tarbert; Hunterston Sands (S.), freq.
Hemilamprops rosea (*Norman*).—U. L. Fyne; Kilbrennan Sd. 1899, r. (S.).
H. cristata, *G. O. Sars*.—F. of Clyde 1860 by Robertson (A. M. N.).

Fam. Leuconidae.

Leucon nasica, *Kroyer*.—U. L. Fyne; off Arran, E. side 43 f. (S.).
Eudorella emarginata (*Kroyer*).—Rothesay B., 12 f. (R.): d. w. bet. Arran and Ayrshire (S.).
E. truncatula (*Sp. Bate*).—L. Fyne; nr. Ailsa Craig (S.).
Eudorellopsis deformis (*Kroyer*).—Nr. Ailsa Craig; off Sanda I. (S.).

Fam. Diastylidae.

Diastylis rathkii (*Kroyer*).—E. L. Tarbert, 1886-87 (S.), r.
D. rugosa, *G. O. Sars*.—L. Ranza, 5 f.; Blackwaterfoot, Arran, 20 f. (R.): E. L. Tarbert, 1886 (S.).
D. lucifera (*Kroyer*).—Clyde district, mod. r. (S.).
D. rostrata (*Goodsir*).—Blackwaterfoot, 20 f. (R.): Kames B., Bute, 15 f. (S.).
D. tumida (*Lilly*).—R., Blackwaterfoot, 20 f. (R.).
D. spinosa, *Norman*.—Dredged Blackwaterfoot; W. side of Helensburgh Pier l. w. (R.).
Diastyloides bispinata, *G. O. Sars*.—Blackwaterfoot, 20 f. (R.): L. Fyne; off E. of Arran 40 f.; nr. mouth of estuary 64 f. (S.).
Leptostylis villosa, *G. O. Sars*.—Bet. Arran and Ayrshire coast, 40 f. 1896 (S.).

Fam. Pseudocumidae.

Pseudocuma cercaria (*V. Ben.*).—Kames B., Millport (R.): freq. and gen. dist.
P. similis, *G. O. Sars*.—D. w. nr. seaward limit of estuary, 12/1899 (S.).

Fam. Nannastacidae.

Nannastacus unguiculatus, *Sp. Bate*.—Kames B., Millport, m. c. (R.): L. Fyne; Kilbrennan Sd., &c. (S.).
Cumella pygmaea, *G. O. Sars*.—Cumbrae (R.): L. Fyne; Whiting B.; Kilbrennan Sd., &c. (S.).

Fam. Campylaspidae.

Campylaspis rubicunda (*Lilly*).—U. L. Fyne, 30-40 f.; E. of Arran, 40 f.; E. of Sanda I., 30 f. (S.).
C. sulcata, *G. O. Sars*.—Single specimen off Cumbrae 20 f., 15/8/88 (A. M. N.).
C. costata, *G. O. Sars*.—U. L. Fyne 30-40 f.; Kames B., Bute, 15-20 f. (S.): off Cumbrae, 20 f. (A. M. N.).

Sub-Order Isopoda.

ARRANGEMENT AND NOMENCLATURE.

Sars, Prof. G. O., *Crustacea of Norway*, vol. ii.

AUTHORITIES.

(R.)=Robertson, Dr. D., F.G.S., F.L.S., "Catalogue of the Amphipoda and Isopoda of the Clyde," Parts i. and ii. (*N. H. Soc. Glas. Trans.*, vol. ii. iii. (N. S.)).
 Stebbing, Rev. T. R. R., *History of the Crustacea*.
 (A. M. N.)=Norman, Rev. Canon, A.M., D.C.L., F.R.S., "British Isopoda Chelifera" (*Ann. and Mag. N. H.*, Ser. 7, vol. iii., 1899).

(S.)=Scott, Thos., F.L.S., "Notes on some Scottish marine Isopods" (*Ann. Scot. N. H.*, 1898).

EXPLANATION OF ABBREVIATIONS.

(R. I., 70)=reference to page 70, Part I. of Robertson's Catalogue.

(R. I., 24, 73, II, 15.)=recorded three times under different or corrected names.
This formula used both in Isopoda and Amphipoda.

Fam. Apseudidae.

Apseudes talpa (*Mont.*)—Roots of *Laminaria* Tan, Cumbrae, 7 f. (R. II., 28).
Fam. Tanaidae.

Tanais cavolinii, *M.-Edw.* (*T. vittatus*, *Bate and Westw.*).—Farland Pt., Cumbrae (A. M. N.): amongst barnacles at l. w. E. L. Tarbert (S.): Cumbrae (R. I., 70).

Paratanais batei, *G. O. Sars.*—Cumbrae (R. I., 70): L. L. Fyne; Whiting B. (S.).

Leptognathia longiremis, (*Lilly*).—L. Fyne (S.).

L. rigida, (*Bate & Westw.*).—Nr. Tan Buoy 7-8 f. (R. I., 70).

L. breviremis (*Lilly*).—Bet. Arran and Ayrshire, 40 f.; Tarbert Bank; L. L. Fyne, 25 f. (S.).

L. brevimana (*Lilly*).—Off Sanda, I. 20-30 f.; Ballantrae Bank, 14 f. (S.).

Tanaopsis laticaudata, *G. O. Sars.*—Nr. Cumbrae (A. M. N.: R. II., 29): L. Gair (L. Fyne), 2-3 f.; Campbeltown L. 5 f. (S.).

Pseudotanais forcipatus (*Lilly*).—Tarbert Bank; L. L. Fyne 25 f. (S.).

Fam. Gnathiidae.

Gnathia maxillaris (*Mont.*).—Gen. dist.; female more c. than male.
G. halidaii (*Sp. Bate*).—Kames B., Cumbrae (R. I., 71).

Fam. Aegidae.

Aega tridens, *Leach*.—Off S. end of Cumbrae; nr. Arran (R. I., 72): Rothesay B. (S.): parasitic on fishes.
[*Ae. rosacea* (*Risso*), (*Ae. bicarinata*, *Leach*).—L. Fyne 15 f. (R. I., 72).]

Fam. Cirolanidae.

Cirolana borealis (*Lilly*), (*C. spinipes*, *Bate and Westw.*).—Cumbrae; L. Cumbrae, N. side, 27 f.; mouth of L. Fyne 37 f. (R. I., 72): not r. w. from d. w.

C. cranchii, *Leach*.—Farland Pt., Cumbrae, 14 f. r. (R. I., 72).

Conilera cylindracea (*Mont.*).—Bet. Bute and Cumbrae, 50 f. (R. I., 73): F. of Clyde (A. M. N.).

Eurydice achata (*Slabber*), (*E. pulchra*, *Leach*).—Fairly freq. shall. sandy bays (R. I., 74).

Fam. Sphaeromidae.

Sphaeroma serratum (*Fabr.*).—Taken on Ayr sands (R. I., 81).

S. curtum, *Leach*.—Off Portincross, Ayrshire, 6 f. (R. I., 82).

S. rugicauda, *Leach*.—Brackish pools; Hunterston (R. I., 82): Inverkip; Langbank, c. (S.).

S. hookeri, *Leach*.—Brackish pool, S. side of L. Cumbrae (R. II., 29).

Dynamene rubra, *Mont.*.—Cumbrae (A. M. N. vide R. I., 82).

D. viridis, *Leach*.—Bet. t. m., Portloy, Cumbrae (R. II., 29).

Naesa bidentata (*Adams*).—With the last at Portloy (R. I., 84). [Note.—For last three species and *D. montague*, *Leach*, vide Stebbing, p. 361.]

Fam. Limnoriidae.

Limnoria lignorum (*Rathke*).—C. on woodwork of piers and jetties (R. I., 77).

Fam. Idotheidae.

- Idothea baltica* (*Pallas*).—(*I. tricuspidata*, *R.*).—C. and gen. dist. (R. I., 79).
I. pelagica, *Leach*.—Amongst weeds at l. w. Cumbrae (R. I., 79): rocks nr. l. w., E. L. Tarbert (S.).
I. emarginata (*Fabr.*).—Amongst weeds at l. w., Cumbrae (R. I., 80).
I. linearis (*Pennant*).—In a sandy pool at l. w., Cumbrae (R. I., 80).
Zenobiana prismatica, *Risso*, (*Idotea parallela*, *R.*).—Dredged bet. Tan Buoy and Hawk's Neb, Bute, 50 f. (R. I., 80).
Stenosoma acuminata (*Leach*), (*Idotea acuminata*, *R.*).—Nr. l. w., Cumbrae; sandy pools, Kilchattan B. (R. I., 81).

Fam. Astacillidae.

- Astacilla longicornis* (*Sowerby*), (*Arcturus longicornis*, *R.*).—gen. dist. (R. I., 78).
[*A. gracilis* (*Goodsir*).—(R. I., 78), is the male of *A. longicornis* (*Sars.*)].
Arcturella dilatata, *G. O. Sars*. (*Leacia dilatata*, *R.*).—Off Blackwaterfoot, Arran, 20 f. (R. II., 28).

Fam. Asellidae.

- Asellus aquaticus* (*Lin.*).—C. in fresh-water ponds, canals, &c. (R. I., 77)

Fam. Janiridae.

- Janira maculosa*, *Leach*.—L. w., Portloy, Cumbrae; mouth of L. Fyne 40-60 f.; off Dunoon 13 f. (R. I., 76); E. L. Tarbert; Campbeltown L., &c. (S.).
Jaera marina (*Fabr.*).—C. in shall. w., and under stones bet. t. m. (R. I., 74).
[*J. nordmanni* (*Rathke*).—(R. I., 75). Sars suggests that the form described by Spence Bate under this name may not be the true Rathkian species but the male of *J. marina*].

Fam. Munnidae.

- Munna boeckii*, *Kr.*.—Whiting B. (S.).
M. kroyeri, *Goodsir*.—Cumbrae, from nests of *Lima hians* (R. I., 76).
[*M. Whiteana*, *Sp. Bate*.—(R. I., 75), is the female of *M. kroyeri* (*Sars.*)].
Paramunna bilobata, *G. O. Sars*.—F. of Clyde (A. M. N.): E. L. Tarbert; Kilbrennan Sd., &c. (S.).
[*Leptaspidea brevipes* (*Sp. Bate*).—Off Cumbrae (R. I., 76). Norman suggests this is a *Pleurogonium*.]
Pleurogonium rubicundum, *G. O. Sars*.—F. of Clyde (A. M. N.). At mouth of estuary 64 f. Dec. 7th, 1899 (F. S. G.).
P. inerme, *G. O. Sars*.—Off Farland Pt., Cumbrae, 23 f. (R. II., 29): Kilbrennan Sd.; Campbeltown L. (S.).
P. spinosissimum, *G. O. Sars*.—Off Farland Pt., Cumbrae (R. II., 29).

Fam. Munnopsidae.

- Eurycope phallangium*, *G. O. Sars*.—L. Fyne, r., 30-40 f. (S.).

Fam. Oniscidae.

- Ligia oceanica* (*Lin.*).—Under stones, &c.; more or less c. in various places.
(Note:—This and the following eight species are land Isopods, and may be found under stones, old wood, &c.)

Fam. Trichoniscidae.

- Trichoniscus pusillus*, *Brandt*.—Cumbrae (R. I., 86): Rothesay, nr. Aquarium (S.). Prob. not unc. if carefully sought for.
T. roseus, *Koch*.—E. Tarbert (S.).

Fam. Oniscidae.

- Oniscus asellus* (*Lin.*).—Gen. dist. and mod. c.
Philoscia muscorum (*Scopoli*).—Widely dist. and not very unc. (R. I. 85).
Porcellio scaber, *Latr.*—C. and gen. dist.
P. pictus, *Brandt*.—Cumbrae (R.): Ayrshire (Smith): nr. Campbeltown, 1896 (S.).
Cylisticus convexus (*De Geer*), (*Porcellio armadilloides*, *R.*).—Kilwinning (Smith, vide R. I. 88): Lanarkshire; Rothesay (S.).
Armadillidium vulgare (*Latr.*).—Mouth of Garnock (Smith, see R. I. 88).

Fam. Bopyridae.

- Cryptothir balani* (*Sp. Bate*).—One taken from a *Balanus*, on shore, E. L. Tarbert, 1886 (S.).
Pseudione affinis, *G. O. Sars*.—On a *Pandalus montagui* in shrimp trawl nr. Sanda I., 5/97 (F. S. G.).
P. hyndmanni (*Bute & Westw.*).—L. Fyne, 1886; and Clyde (S.).
P. crenulata, *G. O. Sars*.—From a small *Munida rugosa* in shrimp trawl, nr. Sanda I., 5/97 (F. S. G.).
Pleurocrypta hendersonii, *Giard & Bonnier*.—On *Galathea dispersa*. Clyde (Henderson, vide Stebbing).
Atheleges paguri (*Rathke*).—On *Eupagurus bernhardus*, L. Fyne; E. L. Tarbert.
Phryxus abdominalis (*Kr.*).—On *Spirontocaris spinosus* in shrimp trawl, nr. Sanda I. (F. S. G.).

Fam. Dajidae.

- Aspidophryxus peltatus*, *G. O. Sars*.—On backs of *Erythrops* nr. Davaar I., &c.

NOTE.—A recent monograph on the Bopyridae by M. Jules Bonnier gives the following Clyde records. Bonnier often differs from Sars as to what are distinct species.

- Bopyroides sarsi*, *Bonnier*.—Branchial cavity of a typical *Spirontocaris spinosus* (*Sowerby*), dredged nr. mouth of Clyde estuary (F. S. G., Aug. 30th, 1899). Distinct from *B. hippolytes* (*Kr.*).
Pseudione affinis, *G. O. Sars*.—Branchial cavity of *Pandalus leptorhynchus*, Kinahan. Firth of Clyde (*Hoyle*).
P. hoylei (*Giard & Bonnier*).—Branchial cavity of *Pandalus montagui*, Leach. Firth of Clyde (Henderson; Hoyle; F. S. G.; Patience). According to Sars *P. hoylei* is identical with *P. affinis*.
P. hyndmanni (*Bate & Westwood*).—Branchial cavity of *Eupagurus bernhardus* (*Lin.*). Clyde (Henderson and myself). See remarks by Prof. Sars and M. Bonnier.
P. insignis (*Giard & Bonnier*) (= *P. crenulata*, *G. O. Sars*, of my list).—According to M. Bonnier the *Pseudione* of *Munida rugosa* is a different sp. from *P. crenulata*, *G. O. Sars*, found on *Munida tenuimana*; Sars considers them the same species.
P. sp..—From the branchial cavity of *Pandalina brevirostris* (*Rathke*). Clyde (Henderson, as *Bopyrus sp.*).
^(?) *P. sp.*.—From the branchial cavity of *Eupagurus prideauxi* (*Leach*). Clyde (Henderson, as *Bopyrus sp.*).
Pleurocrypta hendersoni, *Giard & Bonnier* (= *P. marginata*, *G. O. Sars*, Scott, in 18th Ann. Rept. Fishery Board for Scotland, Pt. III, p. 403). From branchial cavity of *Galathea dispersa*, Bate. F. of Clyde (Henderson; F. S. G.; Patience). According to M. Bonnier the *P. marginata*, *G. O. Sars*, from Norwegian specimens is different from that on English specimens of the same decapod.

- P. intermedia**, Giard & Bonnier (=P. microbranchiata, G. O. Sars).—Branchial cavity of *Galathea intermedia*, Lilly, F. of Clyde (Henderson, as *Bopyrus* sp.).
- Atheleges paguri** (Rathke).—On abdomen of *Eupagurus bernhardus* (Lin.). F. of Clyde (Henderson (as *Phryxus paguri*) ; F. S. G.).
- Phryxus abdominalis** (Kr.).—On abdomen of *Spirontocaris spinus* (Sowerby), *S. securifrons*, Norm., and *Pandalus montagui*, Leach. F. of Clyde (Henderson ; F. S. G. ; Patience). M. Bonnier doubts whether the *Phryxus* found on these species are identical with *abdominalis*, Kroyer, found on *Spirontocaris gainardi* (M.-Edw.). Prof. G. O. Sars, on the other hand, regards them as belonging to that species.
- P. cranchii**, Giard & Bonnier.—On abdomen of *Hippolyte cranchii*, Leach. F. of Clyde (Henderson as *Phryxus* sp.).

Sub-Order Amphipoda.

ARRANGEMENT FOLLOWED.

Prof. G. O. Sars, *Crustacea of Norway*, vol. i.

AUTHORITIES.

- Robertson, Dr. D.; F.G.S., F.L.S.—“Catalogue of the Amphipoda and Isopoda of the Clyde” (*N. H. Soc. Glas. Trans.*, N.S.), vol. ii. p. 1, vol. iii. p. 199.
 Norman, Rev. Canon A. M., D.C.L., F.R.S., “A Revision of the British Amphipoda,” *Ann. and Mag. N. H.*
 Stebbing, Rev. T. R. R., Papers on British Amphipoda.
 Walker, A. O., Papers on British Amphipoda.

Fam. Hyperidae.

- Hyparia galba** (Mont.).—In folds of *Rhizostoma pulmo*, Cumbrae (R. I, 64) : E. L. Tarbert (S.). F. of Clyde (A. M. N.).
Hyperoche tauriformis (B. & W.).—Off Sanda ; nr. Ailsa Craig ; Kilbrennan Sd. ; U. L. Fyne (S.).
Parathemisto obliqua (Kr.).—Nr. Sanda, in surface net (R. I, 65) : off Ailsa Craig (F. S. G.).

Fam. Orchestiidae.

- Talitrus locusta** (Pallas).—Sandy shores, abund. (R. I, 10). U. L. Fyne (S.).
Orchestia littorea (Mont.).—Under stones nr. and above h. w. m., freq. (R. I, 11).
O. mediterranea, Costa.—Cumbrae ; mouth of R. Garnock (R. I, 13) : bet. Fairlie and Hunterston (S.). Inveraray (A. M. N.).
Hyale nilssoni (Rathke).—Kames B., Millport (R. I, 13).
H. lubbockiana (Bate).—Largs Pier ; tide pools Cumbrae (R. I, 13-14) : E. L. Tarbert ; Fintry B., Bute (S.).

Fam. Lysianassidae.

- Normanion quadrimanus** (B. & W.).—Off Farland Pt., Cumbrae (R. II, 12) : Nr. Davaar Is., 1898 ; Tarbert Bank, L. L. Fyne, 1901 (F. S. G.).
Acidostoma obesum (Bate).—Off Farland Pt., Cumbrae (R. I, 17).
Lysianax septentrionalis, Della Valle.—Cumbrae ; off Skelmorlie Buoy ; W. end of L. Cumbrae (R. I, 15-16) : E. L. Tarbert ; U. L. Fyne (F. S. G.).
L. ceratinus A. O. W.—F. of Clyde (A. M. N.).
L. atlanticus (M. Edw.).—Nr. Tan Buoy, Cumbrae (R. I, 17).
Socarnes vahli (Kr.).—N. side of Clach Rock, Cumbrae, 6 or 7 f. (R. II, 5).
S. erythrophthalmus, Robertson.—Tan, Cumbrae ; Castle-B., L. Cumbrae ; Rothesay ; Millport B. (R. II, 6).

- Sophrosyne robertsoni**, Stebbing and Robertson—(Clyde R. II, 9) : U. L. Fyne (M.).
- Perrierella audouiniana** (Bate).—In a sponge, mouth of L. Fyne, 40-60 f. (R. I, 16; II, 7) : U. L. Fyne (M.) : Tarbert Bank ; E. of Sanda, 30 f. (F. S. G.).
- Callisoma hopei**, A. Costa [= *Callisoma crenata* (Bate)].—L. Fyne, 80 f. ; Kilchattan B., 30 f. (R. I, 19). Gen. dist.
- Hippomedon denticulatus** (Bate).—Nr. Tan Buoy ; Castle-B., L. Cumbrae, &c. (R. I, 18 ; II, 9). Gen. dist.
- Orchomene humilis** (A. Costa) (= *Anonyx edwardsii*, Bate and Westwood).—Tan Buoy ; L. Striven ; L. Goil ; L. Fyne (R. I, 17 ; II, 11) ; Whiting B. (S.).
- Tryphospha pinguis** (Boeck).—Dredged w. of Tan Buoy, 18 f. (R. II, 9).
- T. nana** (Kr.).—W. of Tan Buoy, 14 f. (R. II, 10) : off Ballantrae, rather r. (F. S. G.).
- T. minuta** (Kr.).—Minard, U. L. Fyne (M.).
- Tryphosella sarsi**, Bonnier.—Off Farland Pt., Cumbrae, 23 f. (R. II, 11) : off Largabruach ; off Inveraray, U. L. Fyne (F. S. G.).
- T. höringii**, (Boeck).—Cumbrae, on back of a codfish (R. I, 92).
- T. nanoides** (Lillj.).—Cumbrae, on back of a codfish (R. I, 92).
- [**T. pusilla**, G. O. Sars.—Tan buoy 12 f. (R. II, 12)].
- Tryphosites longipes** (Bate).—Off N.E. corner of L. Cumbrae 14 f. ; bet. Kilchattan and Cumbrae 66 f. (R. I, 18) : Kilbrennan Sd. ; E. L. Tarbert, &c. (F. S. G.).
- Hoplonyx cicada** (Fabr.).—Bet. the Allans, Cumbrae, 2-4 f. (R. I, 18) : Tarbert Bank, L. L. Fyne, 20-25 f. ; rather r. (F. S. G.).
- H. similis**, G. O. Sars.—Off Farland Pt., Cumbrae (R. II, 12).
- Lepidepecreum carinatum** (Bate).—L. Ranza B. 5 f. ; S. E. of Bute 10-12 f. (R. I, 91).
- Euonyx chelatus**, Norman.—Off Mull of Kintyre 49 f. (R. I, 90) : several dredged nr. mouth of estuary, 11/95 (F. S. G.).

Fam. Pontoporeiidae.

- Bathyporeia pelagica**, Bate.—Sandy shores bet. t. m., various places (R. I, 42) : Cumbrae (A. M. N.).
- B. norvegica**, G. O. Sars.—Cumbrae (A. M. N.).
- B. robertsoni**, Bate.—Cumbrae (A. M. N.).
- Haustorius arenarius** (Slabber).—Sandy shores nr. l. w. m., mod. c. (R. I, 23) : Cumbrae (A. M. N.).
- Urothoe elegans**, Bate.—Cumbrae (A. M. N.).
- U. marinus**, Bate.—Bet. t. m. Irvine and Fairlie ; Lamlash B. ; Kames B. ; Fintry B. ; Balloch B., Cumbrae (R. I, 29-30) : Barmore, L. L. Fyne (S.).
- U. brevicornis**, Bate.—F. of Clyde (R., vide A. M. N.).
- Argissa hamatipes** (Norman).—Kilchattan B., Bute, 8-9 f. (R. II, 12) : not unc. in seaward part of estuary.

Fam. Phoxocephalidae.

- Phoxocephalus holboelli** (Kr.).—Dredged bet. the Allans, taken nr. l. w. m. Balloch B., Cumbrae (R. I, 24) : Millport B. bet. t. m. (A. M. N.).
- P. fultoni**, T. Scott.—Cumbrae, l. w. m. (R.) : L. Ryan (Fullerton) : F. of Clyde (A. M. N.).
- P. simplex** (Bate).—Tan Buoy 6 f. (R. I, 23).
- P. oculatus**, G. O. Sars.—(I think R. obtained this sp. in the Clyde).
- Harpinia neglecta**, G. O. Sars.—L. Ranza, 6-8 f. ; L. Fyne 80 f. (R. I, 24) : Clyde, gen. dist. (A. M. N.).
- H. crenulata**, G. O. Sars.—Campbeltown L. (S.) : off Farland Pt. ; Cumbrae (A. M. N.) : r.

Fam. Ampeliscidae.

- Ampelisca typica** (*Bate*).—R. records this as *A. gaimardii* (*Spence Bate*) from various places (R. I, 20): Cumbrae (A. M. N.).
- A. tenuicornis**, *Lilly*.—L. Fyne 80 f.; bet. Tan Buoy and Bute; L. Goil 12 f. (R. I, 21-22): Campbeltown L. (A. M. N.).
- A. assimilis**, *Boeck*.—Row B., Gareloch, 10-12 f. (R. I, 22): off Sanda (F. S. G.).
- A. brevicornis** (*A. Costa*).—Off S. shore, Cumbrae; opposite Largs, 15 f.; off Fairlie 20 f.; l. w. Crarae, L. Fyne, &c. (R. I, 21): nr. Ailsa Craig; off Sanda I. (F. S. G.).
- A. macrocephala**, *Lilly*.—Off Brodick 80 f. (R. I, 22): Cumbrae (A. M. N.).
- A. spinipes**, *Boeck*.—Off Cumbrae L.H. 60 f.; off Holy I. 36 f.; nr. Skate I. 80-100 f. (R. I, 91): Cumbrae (A. M. N.).
- Haploops tubicola**, *Lilly*.—Off L. Cumbrae 60 f.; L. Striven 35 f.; off Dunoon 15-25 f.; off Fairlie 6-8 f. (R. I, 22, 92): off Sanda I. (A. M. N.).
- H. setosa**, *Boeck*.—Off Skate I., L. Fyne, 100 f., two specimens (R. I, 23).

Fam. Stegocephalidae.

- Stegocephaloides christianiensis** (*Boeck*).—L. Fyne 40-70 f. (R. II, 20): nr. head of L. Fyne (F. S. G.).

Fam. Amphilochidae.

- Amphilochus manudens**, *Bate*.—Off Tan Buoy 7-8 f.; off Cumbrae L.H., 60 f. (R. I, 28).
- A. tenuimanus**, *Boeck*.—Off Farland Pt., Cumbrae, 17 f. (R. I, 28): Ballantrae Bank (F. S. G.).
- Amphilochoides serratipes** (*Norman*).—Off Blackwaterfoot 20 f.; off Clach Rock, Cumbrae (R. II, 20-22): off Sanda; at Ballantrae Rk.; off Ailsa Craig (F. S. G.).
- Gitanopsis bispinosa** (*Boeck*).—Off Blackwaterfoot 20 f. (R. II, 20).
- Gitana sarsii**, *Boeck*.—Kilchattan B. 5 f. (R. II, 2): Whiting B., Arran; off Sanda I. 20 f. (F. S. G.).
- Cyproidia damnoniensis**, *Stebbing*.—Off Blackwaterfoot; off Farland Pt., Cumbrae (R. II, 21).
- Stegoplax brevirostris** (*T. and A. Scott*).—Tarbert Bank, L. Fyne, r. (S.).

Fam. Stenothoidae.

- Stenothoe marina**, *Bate*.—Cumbrae at l. w.; off Dunoon (R. I, 15): Whiting B.; off Ailsa Craig (F. S. G.).
- S. monoculoides** (*Montagu*).—Various places about Cumbrae (R. I, 14): F. of Clyde (A. M. N.).

Fam. Metopidae.

- Metopa alderi** (*Bate*).—Bet. Cumbrae L. H. and Arran (R. I, 90).
- M. borealis**, *G. O. Sars*.—Off Farland Pt., Cumbrae, 20 f. (R. II, 21): E. of Arran 30-40 f. (F. S. G.).
- M. rubrovittata**, *G. O. Sars*.—Off the N. side of L. Cumbrae, 10-15 f. (R. II, 21).
- [**M. affinis**, *Boeck*.—W. of Tan Buoy; Mull of Kintyre (R. I, 90).]
- M. pusilla**, *G. O. Sars*.—Off Farland Pt., Cumbrae, 20-24 f. (A. M. N.).
- Metopella nasuta**, (*Boeck*).—Off Farland Pt., Cumbrae, 19 f. (R. II, 22).

Fam. Cressidae.

- Cressa dubia**, *Bate*.—W. of Tan Buoy, 7-8 f. (R. I, 15): Whiting B.; off Sanda I. (F. S. G.); Cumbrae (A. M. N.).

Fam. Leucothoidae.

Leucothoë spinicarpa (*Abildgaard*).—Nr. Tan Buoy, 5 f.; L. Fyne, 92 f. (R. I, 39); Whiting B. (F. S. G.).
 [*L. incisa*, *Robertson*.—L. w. Balloch B.; Farland Pt., Cumbrae, 20 f. (R. II, 23) (?); L. Gair (L. Fyne) (F. S. G.). Norman considers this only a form of the next species.]

L. lilljeborgii, *Boeck*.—L. Gair; head of U. L. Fyne; r. (F. S. G.).

Fam. Oediceridae.

Monoculodes carinatus, *Bate*.—Several places about Cumbrae (R. I, 25).
M. borealis, *Boeck*.—Off Blackwaterfoot, (R. II, 15).
M. tuberculatus, *Boeck*.—In d. w. E. of Arran, r. (F. S. G.).
M. packardi, *Boeck*.—L. Striven, 40 f. (R. II, 14); L. Fyne, 30-60 f.; E. of Arran, 40 f.; not very r. (F. S. G.).
Periocolodes longimanus (*Bate*).—Gen. dist., not unc. in d. w. (vide R. I, 26).
Pontocrates altamarinus (*Bate*).—Callum's Bay, Bute, 16 f. (R. I, 27); U. L. Fyne (S.).
P. norvegicus, *Boeck*.—Shall. w. in sandy bays freq. (R. I, 26).
Synchelidium brevicarpum (*Bate*).—Nr. Cumbrae; L. Fyne; Blackwaterfoot (R. I, 25; II, 16).
Halimedon parvimanus (*Bate*).—Various places (R. I, 24, 93; II, 15); gen. dist.; not r. in d. w.

Fam. Pleustidae.

Paramphithoë bicuspis (*Kr.*).—In surface tow-net after sunset, Cumbrae (R. I, 36).
P. monocuspis, *G. O. Sars*.—Off Sanda I., r. (F. S. G.).
P. assimilis, *G. O. Sars*.—Off Cumbrae, 15-20 f. (R. I, 94).
Stenopleustes nodifer, *G. O. Sars*.—Off N. side of L. Cumbrae, 20 f.; off Fairlie Perch, 20 f. (R. II, 18).

Fam. Epimeridae.

Epimeria cornigera (*Fab.*).—Not unc. in d. w. (R. I, 32). Mouth of estuary, and in d. w. bet. Arran and Ayrshire coast (F. S. G.).

Fam. Iphimedidae.

Iphimedia obesa (*Rathke*).—Gen. dist.; not very r.
I. minuta, *G. O. Sars*.—Off Farland Pt., Cumbrae, 20 f. (R. II, 23).
I. elbanæ, *Bate*.—Dredged nr. mouth of estuary (F. S. G.).
Odius carinatus (*Bate*).—Off Garnock Beacon, 5½ f. (R. II, 31).

Fam. Laphystiidae.

Laphystius sturionis (*Kr.*).—Kilbrennan Sd. (R. I, 93); Rothesay B. (S.). Parasitic on Cod, Anglerfish, etc.

Fam. Syrrhoidae.

Syrrhoë fimbriatus, *Stebbing and Robertson*.—Clyde (R. II, 22).
Bruzelia typica, *Boeck*.—W. of Tan Buoy, 12-20 f. (R. II, 22).

Fam. Eusiridae.

Eusirus longipes, *Boeck*.—W. of Tan Buoy, 10 f.; L. Goil, 12 f. (R. I, 38); nr. Sanda I.; head of L. Fyne, etc. (F. S. G.).

Fam. Calliopiidae.

Apherusa bispinosa (*Bate*).—C. in Clyde (R. II, 17); Kilbrennan Sd.; off Ailsa Craig, etc. (F. S. G.).
A. jurinii (*M. Edw.*).—Bet. t. m., Farland Pt., Cumbrae (R. II, 17).

Calliopius laeviculus (*Kr.*).—Mod. c., Kames B., Balloch B., Cumbrae
(R. I, 37, 39, 94).

C. rathkei (*Zaddoch*).—Amongst weedy stones at l. w. Cumbrae (R. I, 38).

Fam. Atyidae.

Paratylus swammerdami (*M.-Edw.*).—C. and gen. dist.

P. vedlomensis (*Bate*).—Not unc. in Clyde (R. I, 36).

P. uncinatus, *G. O. Sars*.—Kilbrennan Sd., 4/96 (F. S. G.)

Fam. Dexaminidae.

Dexamine spinosa (*Mont.*).—Mod. c. all round our shores (R. I, 32).

D. thea, *Boeck*.—Portloy, Cumbrae; Clach Rock, and off Hunterstone,
opposite Millport (R. I, 33, 34); head of L. Fyne (F. S. G.).

Tritaeta gibbosa (*Bate*).—Freq., by dip-net among seaweed, &c. (R. I, 34):

nr. Davaar I. (F. S. G.).

Guernea coalita (*Norm.*).—Taken twice off Millport (R. II, 24): gen. dist.
esp. in d. w., scarce (F. S. G.).

Fam. Melphidippidae, *Stebbing* (1899).

Melphidippa goesi, *Stebbing*.—On N. side of L. Cumbrae 20 f. (R. II, 19).

Melphidippella macera (*Norm.*).—N. side of L. Cumbrae (R. II, 19): Kil-
brennan Sd.; off Ailsa Craig (F. S. G.).

Fam. Gammaridae.

Amathilla homari (*Fabr.*).—In surface-net after sunset, Cumbrae, and
Sanda B. (R. I, 41, 92); Bet. t. ms., E. L. Tarbert (S.).

Gammarus marinus, *Leach*.—C. under stones bet. t. ms., and in tidal pools
(R. I, 50).

G. locusta (*Lin.*).—Bet. t. ms., and dredged nr. Cumbrae, 6-35 f. (R. I, 50, 94).
Gen. dist.

G. duebeni (*Lillj.*).—Fresh water lochs nr. Campbeltown (S.).

G. pulex (*de Geer*).—C. in slow streams and ditches (R. I, 52).

Melita palmata (*Mont.*).—Clyde (A. M. N.) 1889.

M. obtusata (*Mont.*).—C. and gen. dist. (R. I, 47, 48).

M. gladiosa, *Bate*.—Taken bet. the Allans, Cumbrae, 4-5 f. (R. II, 18).

Eriopisa elongata (*Bruzel*).—Taken in Clyde by Norman (R. II, 25).

Maera othonis (*M.-Edw.*).—In various parts of Clyde and L. Fyne (R. II, 53):
nr. Davaar I. (F. S. G.).

M. semiserrata (*Bate*).—Dredged twice at Tan Buoy (R. I, 52).

M. loveni (*Bruzel*).—Off L.H., L. Cumbrae, 55-60 f.; off Arran, 80 f.
(R. I, 52).

Gammarella brevicaudata, (*M.-Edw.*).—Off Hunterstone (R. I, 47; II, 24).

Megaluropus agilis, *Norm.*.—Garrison B., and Kames B., nr. Millport (R. II,

25); nr. Sanda I, 5/9/99 (F. S. G.).

Cheiocratus sundewalli (*Rathke*).—Various places (R. I, 45, 46); nr. Lar-
gabruach, U. L. Fyne; Tarbert Bank (F. S. G.).

C. intermedius, *G. O. Sars*.—E. L. Tarbert (S., 1886); L. Gair, (L. Fyne);
Tarbert Bank, 17-20 f. (F. S. G.).

C. assimilis, (*Lillj.*).—L. L. Fyne, 104 f.; off Farland Pt., Cumbrae, 20 f.
(R. II, 18); Tarbert Bank, 20-25 f. (F. S. G.).

Fam. Lilljeborgiidae.

Lilljeborgia pallida, *Bate*.—Off Farland Pt., Cumbrae, 20 f. (R. II, 23).

L. kinahani (*Bate*).—Nr. Tan Buoy, 7 f. (R. I, 46); Tarbert Bank, 20 f.
(F. S. G.).

Fam. Aoridae.

[*Phaedra antiqua* (*Bate*).—In surface-net off the Allans, Millport B., after
sunset (R. I, 94).]

Microdeutopus anomalus (*Rathke*).—Off Tan Buoy, &c., Cumbrae (R. I, 41). Whiting B. (S.).

M. versiculatus (*Bate*).—Dredged a little beyond l. w., Cumbrae (R. I, 42). *M. danmoniensis* (*Bate*).—One S. of Tan Buoy (R. I, 40): on E. side and in centre of U. L. Fyne 10-17 f. (M.).

Aora gracilis (*Bate*).—Gen dist., but mod. r.

Autonoë websteri (*Bate*).—Under stones nr. l. w., Portloy, Cumbrae, (R. I, 40).

Fam. Photidae.

Leptocheirus hirsutimanus (*Bate*).—Clach Rock, nr. Tan Buoy; bet. Fairlie and Millport (R. I, 42): off Sanda I. (F. S. G.).

L. pilosus, *Zaddach*.—Castle B., L. Cumbrae, 2-3 f. (R. II, 27): Tarbert Bank, L. L. Fyne, 20-25 f. (F. S. G.).

Gammaropsis maculata, *Johnstone*.—Otter Spit, L. Fyne; nr. Cumbrae (R. I, 48). Gen dist.

G. palmata (*Stebbing and Robertson*).—Cumbrae; Clyde one (Steb. and Rob.): off Ballantrae, 24/9/98; Kilbrennan Sd. (F. S. G.).

Megamphopus cornutus, *Norm*.—Off Farland Pt.; Castle B., Cumbrae; off Blackwaterfoot (R. II, 24, 26).

Microprotopus maculatus, *Norm*.—Garrison B.; Kames B., Cumbrae (R. II, 26).

Photis longicaudata (*Bate*).—Millport B.; Clach Rock, Cumbrae; off S. end of Bute, 16 f. (R. I, 95): Kilbrennan Sd. (F. S. G.).

Podoceropsis sophiae, *Boeck*.—Skelmorlie Bank, 10 f.; off Skelmorlie Pt., 39 f.; S. of L. Cumbrae (R. I, 60, 61).

P. excavata (*Bate*).—Gen. dist. and mod. c. (R. I, 60).

Fam. Amphithoidae.

Amphithoë rubricata (*Mont*).—Mod. c.: gen. dist. (R. I, 53, 54).

Pleonexes gammaroides, *Bate*.—Portloy and Kames B., Cumbrae; off L. Cumbrae 8 f. (R. I, 56; II, 27).

Fam. Ischyroceridae.

Ischyrocerus minutus, *Lilly*.—W. of Clach Rock; off Farland Pt.; on the Allans, Cumbrae (R. II, 28).

Jassa pusilla (*G. O. Sars*).—Amongst algae on Millport pier (R. II, 27). Off Sanda I. (F. S. G.).

J. falcata (*Mont*).—Millport pier, amongst algae; on Tan Buoy; Portloy and bet. the Allans, Cumbrae; off Dunoon 4 f. (R. I, 57).

J. dentex (*Czerniawski*) (= *Podocerus herdmani*, *A. O. Walker*).—Off Sanda I. [A. O. Walker now thinks this is only a form of *J. falcata*.]

J. ocia (*Bate*).—N. of Tan Buoy 8 f.; off Clach Rock 6 f. (R. I, 58).

Microjassa cumbrensis (*Stebbing and Robertson*).—Off Farland Pt. 20 f. (R. II, 27).

Parajassa pelagica (*Leach*) [= *Janassa capillata* (*Rathke*)].—On Millport pier; Kames B.; Tan Buoy, Cumbrae; off Dunoon (R. I, 57).

Erichthonius abditus (*Templeton*).—N. of Cloch L.H. 6 f. (R. I, 58): Campbeltown L.; nr. Ailsa Craig (F. S. G.).

E. difformis, *M.-Edw*.—L. w., Portloy; off Farland Pt. Cumbrae 17 f. (R. I, 59).

Fam. Corophiidae.

Siphonoecetes colletti, *Boeck*.—W. of Cumbrae 10 f. (R. I, 59).

Corophium grossipes (*Zin*).—Brackish water; Langbank; Rosneath; Balloch B., Cumbrae; mouth of Garnock (R. I, 61): Inverkip, c. (S.).

C. crassicornis, *Bruzel*.—Balloch B., Cumbrae; shore pool, L. Cumbrae, c. (R. I, 62): Campbeltown L. (F. S. G.).

C. bonelli, *M.-Edw.*—Gen. dist. (Whiting B. etc.).
C. affine, *Bruzel.*—Clyde (A. M. N.); E. L. Tarbert (S.).
Unciola planipes, *Norm.*—Off Sanda I. (F. S. G., 1/9/98).

Fam. Cheluridae.

Chelura terebrans, *Philippi*.—All round coasts in submerged timbers of piers and jetties (R. I, 63).

Fam. Dulichiidae.

Dulichia porrecta, *Bate*.—Off Farland Pt., Cumbrae (R. II, 28).
D. monacantha, *Metzger*.—One found in stomach of haddock caught in Clyde (1897) (S.).
D. tuberculata, *Boeck*.—Dredged nr. Mull of Kintyre 49 f. (R. I, 95).

Fam. Caprellidae.

Phtisica marina, *Slabber*.—Gen. dist. (R. I, 65, 66): freq. Tarbert Bank (F. S. G.).
Protella phasma (*Mont.*).—Gen. dist., and not very r. (R. I, 66): Tarbert Bank freq. (F. S. G.).
Pariambus typicus (*Kr.*).—Gen. dist., not unc. on *Asterias rubens* (R. I, 69).
Caprella linearis (*Lin.*).—Cumbrae among weedy stones l. w.; off Garroch-head, Bute 15 f. (R. I, 67).
C. septentrionalis (*Kr.*).—Cumbrae (R.).
C. acanthifera, *Leach*.—Cumbrae bet. the Allans amongst zostera, plentiful R. I, 68).
[*C. tuberculata*, *Guerin*.—Cumbrae, rare (R. I, 69).]

ORDER BRANCHIOPODA.

Sub-Order Phyllocarida.

Fam. Nebaliidae.

Nebalia bipes (*Fabr.*).—Clyde; L. Fyne, various places (S.).

Sub-order Cladocera.

ARRANGEMENT FOLLOWED.

Prof. C. L. Herrick, *Crustacea of Minnesota*, 1884.

AUTHORITIES.

Baird, W., *Monog. of the Brit. Entomostraca*, 1849.
(A. M. N.) Norman, Rev. Canon A. M., and Prof. G. S. Brady, *A Monograph of Brit. Entomostraca belonging to the Bosminidae, Macrothricidae, and Lynceidae*, 1867.
Brady, Prof. G. S., on *The Brit. species of Entomostraca belonging to the Daphnia and other allied species*, 1898.
Scott, Thomas, F.L.S., "The Fauna of the inland waters of Scotland" (*Ann. Reps. Fish. Bd. for Scotland*).

Fam. Sididae.

Sida crystallina (*Müller*).—Bardowie L., nr. Glasgow; L. Lomond; Greenan L., Bute; Park L., Campbeltown; L. Doon, Ayrshire.
Diaphanosoma brachyurum (*Lievin*).—Bardowie L.; Park L.

Fam. Holopedidae.

Holopedium gibberum, *Zaddach*.—L. Arklet; L. Doon, c. during summer.

Fam. Daphnidæ.

Ceriodaphnia reticulata (*Jurine*).—Possil Marsh.

C. quadrangula, *Müller*.—Bardowie L.; St. Germains' L., Glasgow; Dhu Lochan, nr. L. Lomond; Park L., nr. Campbeltown.

C. laticaudata, *Müller*.—Possil Marsh, rather r.

C. sp..—L. Lomond; L. Doon; appears to be r.

Scapholeberis mucronata, *Müller*.—Bardowie L.; Dhu Lochan, not c.

Simocephalus vetulus, *Müller*.—Possil Marsh; L. Lomond; Greenan L.; Park L.; L. Doon.

Daphnia lacustris, *G. O. Sars*.—L. Lomond, with and without a vertex tooth.

var. *aquilina*, *G. O. Sars*.—L. Arklet, freq.

D. (?) longispina, *Leydig*.—L. Fad, Bute.

var. *nasuta*, *G. O. Sars*.—L. Doon, Ayrshire.

D. (?) jardini, *Baird*.—St. Germains' L., Glasgow. A crested form which may belong either to this or to *D. galeata*, *G. O. Sars*.

Fam. Bosminidæ.

Bosmina longirostris (*Müller*).—Gen. dist. esp. in smaller lochs and ponds.

B. longispina, *Leydig*.—Not unc. in larger lochs, as L. Lomond and L. Doon.

Fam. Macrothricidæ.

Lathonura rectirostris (*Müller*).—L. Doon—the shore on W. side.

Streblocercus minutus, *G. O. Sars*.—Dhu Lochan, nr. L. Lomond, r.

Drepanothrix dentata (*Euren*).—Cumbrae (R.): L. Arklet; L. nr. L'ranza; L. Doon, &c., not unc.

Acantholeberis curvirostris (*Müller*).—Cumbrae (R.): L. Arklet; L. nr. L'ranza, not r.

Ilyocryptus sordidus (*Lieven*).—Cumbrae (R.): Possil Marsh; L. Arklet; L. Doon; Park L., Campbeltown, not r.

Fam. Lynceidæ.

Eurycerus lamellatus (*Müller*).—C. and gen. dist.

Acroporus harpae, *Baird*.—Gen. dist. and mod. c.

Camptocercus macrurus (*Müll.*).—L. Lomond; L. Doon, freq. The Camptocercus found in Scotland is usually ascribed to *C. macrurus*, *Müll.*, but seems to agree better with Schoedlers' *C. rectirostris*.

Alonopsis elongatus, *G. O. Sars*.—C. and gen. dist.

Ledigia quadrangularis (*Leydig*).—Pond nr. Rutherglen (J. Steel)! St. Germains' L., not c.

Graptoleberis testudinaria (*Fischer*).—L. Arklet; L. Doon; freq.

Alona guttata, *G. O. Sars*.—Gen. dist., freq.

A. costata, *G. O. Sars*.—St. Germains' Loch; L. Lomond (shore nr. Bal-maha); not c.

A. quadrangularis (*Müller*).—Gen. dist., mod. c.

A. affinis (*Leydig*).—L. Lomond; L. Doon; L. behind L'ranza; Park L., Campbeltown &c., freq.

A. intermedia, *G. O. Sars* (= *A. neglecta*, *T. S.*)—L. Arklet; L. Doon, not c.

A. rustica, *T. Scott*.—L. Arklet; L. Lomond; L. Doon; Park L.

A. rectangula, *G. O. Sars*.—Nr. E. Tarbert, L. Fyne; Park L., Campbeltown.

Alonella exigua (*Lilljeborg*).—C. and gen. dist.

A. nana, *Baird*.—L. Arklet; L. Lomond; L. Doon, &c., not r., but being so small is easily overlooked.

Pleuroxus trigonellus (*Müller*).—L. Lomond; L. Doon, not r.

P. laevis, *G. O. Sars*.—Possil Marsh is the only station I know of.

P. uncinatus, Baird.—Pond nr. Rutherglen (J. Steel): Possil Marsh; L. Lomond.

Harporthynchus falcatus, G. O. Sars.—L. Arklet; L. Lomond; L. Doon, not r.

Peracantha truncata (Miller).—L. Arklet; Dhu Lochan; Greenan L., Bute; L. Doon.

Chydorus sphaericus, Müller.—C. and gen. dist.

C. globosus, Baird.—Possil Marsh; Bardowie L., nr. Glasgow; L. Lomond; Park L., nr. Campbeltown.

C. (?) latus, G. O. Sars.—Only observed in L. Arklet.

C. barbatus (G. S. Brady).—In most Lochs within Clyde area.

D. coelatus, Schoedler.—L. Arklet; L. Doon; Park L., not c.

Anchistropus emarginatus, G. O. Sars.—Paisley Canal in 1863 (R.). Not observed since.

Monospilis tenuirostris (Fischer).—L. Lomond (shore bet. Balmaha and mouth of R. Endrick), rather r.

Fam. Polyphemidae.

Polyphemus pediculus (Lin.).—Possil Marsh; L. Arklet; L. Lomond; Park L.

Bythotrephes longimanus, Leydig.—L. Arklet; L. Lomond; L. Doon, not unc.

Podon polyphemoides, Leuckart.—F. of Clyde; L. Fyne, sometimes pretty abund.

P. intermedius, Lilljeborg.—Dist. similar to preceding but much less c.

Evadne nordmanni, Loren.—Dist. similar to *Podon*; c., sometimes very abund.

Fam. Leptodoridae.

Leptodora hyalina, Lilljeborg.—L. Arklet; L. Lomond; L. Doon, not unc., esp. during summer.

Sub-Order Branchiura.

Argulus foliaceus, Lin.—Plentiful a few years ago on the Greyling, *Thymallus vulgaris* in the upper waters of R. Clyde (Jas. Steel)!

ORDER OSTRACODA.

ARRANGEMENT AND NOMENCLATURE FOLLOWED.

(B. & N.)=Prof. G. S. Brady and Rev. Canon A. M. Norman, "The marine and fresh water Ostracoda of the N. Atlantic and North-Western Europe (Roy. Soc. Dub. Sci. Trans., vol. iv. (S. ii.) Pt. ii. and vol. v. (S. ii.) Pt. xii.).

R.=Robertson, Dr. D., F.G.S., F.L.S. Ostracoda in "Notes on the fauna and flora of the W. of Scotland," 1876.

"Fresh and Brackish water Ostracoda" (N. H. Soc. Glas., vol. iv., 1879).

B. & R.=Brady and Robertson.

B. C. & R.=Brady, Crosskey, and Robertson.

Fam. Cyprididae.

Cypria exculta (S. Fischer).—Somewhat local but not very r.; Possil Marsh (R.); Cumbrae (B. & N.); Dalry (J. Smith): L. Doon, &c. (S.).

C. ophthalmica (Jurine).—C. and gen. dist.

Cyclocypris laevis (O. F. Müller).—Mod. c.; liable to be confounded with next species.

C. serena (Koch).—One of the most c. and gen. dist.

- C. globosa* (*G. O. Sars*).—Nr. L. Ascog and L. Greenan, Bute (B. & N.): L. Lomond; L. Arklet; L. Doon (S.).
- Scotia browniana* (*Jones*).—Marshy ground at side of L. Fad, Bute (B. & N.): Binend L. 1877 (J. Smith).
- Cypris fuscata*, *Jurine*.—Frankfield L., nr. Glasgow; brackish ponds, Cumbrae (R.): Dalry (J. Smith): L. Doon (S.).
- C. incongruens*, *Ramdohr*.—Pond nr. Paisley; brackish pool, Hunterston &c. (R.): sometimes very abund.
- C. virens* (*Jurine*).—Pond nr. Crosshill; Frankfield L.; Callum's Tarn, Bute (R.): Dalry (J. Smith): Pools nr. Fort Matilda, Greenock; L. Ascog, Bute, &c.
- C. reticulata*, *Zaddoch*.—Possil Marsh; side of Paisley Canal (R.).
- C. obliqua*, *Brady*.—Callum's Tarn, Bute; Cragengower Tarn, Cumbrae (R.): Binend L. (J. Smith): L. Lomond (S.).
- C. prasina*, *Fischer*.—Pond on shore at Hunterston; Pond nr. Ferry Ho., Cumbrae (R.): Brackish pools at Langbank (S.).
- Herpetocypris reptans* (*Baird*).—Mod. c. and gen. dist.
- H. strigata* (*Müller*).—Burnside L. nr. Glasgow; ponds at Port Glasgow; Cumbrae, &c. (B. & N.).
- H. tumefacta* (*B. & R.*).—Cumbrae; L. Greenan; L. Fad; L. Ascog, Bute; Tarbert, &c. (B. & N.).
- Ilyodromus robertsoni* (*B. & N.*).—Side of L. Fad (B. & N.): L. Doon (S.).
- Cypridopsis villosa* (*Jurine*).—L. Lomond; Frankfield L.; ponds, Cumbrae (R.): Dalry, &c. (J. Smith): L. Fad; L. Ascog, Bute (B. & N.): L. Doon.
- C. newtoni* (*B. & R.*).—Little L. and Pilmuir Dam, Renfrewshire; Cumbrae; Bute (B. & N.).
- C. aculeata* (*Liljeborg*).—Brackish pools, S.W. end of Cumbrae (R.): Tarbert; Langbank (S.).
- Pionocypris vidua* (*Müller*).—Pond at Paisley; L. Fad (R.): Kilwinning (J. Smith): L. Lomond.
- P. obesa* (*B. & R.*).—L. Lomond; old dam at Govan Colliery, Glasgow (R.): Kilwinning (J. Smith).
- Potamocypris fulva*, *Brady*.—Paisley Canal; Cumbrae; Kilwinning (R.): Dalry; Pilmuir Dam; L. Goin (J. Smith): L. Fad, and Ascog, Bute; Tarbert.
- Paracypris polita*, *G. O. Sars*.—Off Tarbert, 25 f. (B. & N.): Tarbert Bank, 20 f. (S.).
- Notodromus monacha* (*Müller*).—Cumbrae; L. Cumbrae (R.): Kilwinning (J. Smith): Pools nr. L. Fad, Bute (S.).
- Candonia candida* (*Müller*).—C. and gen. dist.
- C. lactea*, *Baird*.—Paisley Canal (R.): Pilmuir Dam (J. Smith): Park L. and Achy-Lochy, Campbeltown.
- C. compressa* (*Koch*).—Mod. c.; more or less gen. dist.
- C. rostrata* (*B. & N.*).—Hairmyres, nr. E. Kilbride (B. & N.): Possil Marsh; L. Doon (S.).
- C. fabaeformis* (*Fischer*).—Hairmyres; Crosslea, nr. Paisley; Cumbrae, etc. (B. & N.): Dalry, Eaglesham Dams, etc. (J. Smith): L. Lomond.
- C. acuminata* (*Fischer*).—Kilmuir Dam, Renfrewshire; Tarbert (B. & N.): Eaglesham Dam, Craigton Dam, Harelaw Dam (J. Smith): L. Lomond (S.).
- C. euptectella*, *Robertson*.—Tarn, L. Cumbrae; Callum's Tarn, Bute (R.): Nockewart, Dalry (J. Smith): Pools nr. Pt. Glasgow; Mossclavie Pond, nr. Greenock (S.).
- C. hyalina* (*B. & R.*).—Frankfield L.; Possil Marsh, etc. (R.): Long Loch (J. Smith): Loch Lomond; L. Arklet; L. Doon (S.).
- Candonopsis Kingsleii* (*B. & R.*).—Mod. c. and gen. dist.

- Ilyocypris biplicata* (Koch).—Mod. c., but somewhat local.
Pontocypris mytiloides (Norman).—Gen. dist. throughout Estuary.
P. acupunctata, Brady.—Nr. Pt. Glasgow 1892, Campbeltown L. 1897 (S.).
P. trigonella, G. O. Sars.—Gen. dist., not r.
Argilloecia cylindrica, G. O. Sars.—Inveraray; Largs; Campbeltown (R.); off Greenock; L. Long (B. & N.): off Tarbert (S.).

Fam. Bairdiidae.

- Bairdia inflata*, Norman.—Cumbrae; Lamlash (R.); L. Fyne, off Tarbert; Campbeltown L. (S.).
B. complanata, Brady.—L. Fyne (A. M. N.).

Fam. Darwinulidae.

- Darwinula stevensoni* (B. & N.).—L. Lomond (shore S. of Balmaha) (S.).

Fam. Cytheridae.

- Cythere lutea*, Müller.—C. and gen. dist.
C. pellucida, Baird.—Brackish water, freq. (Langbank; Inverkip; Hunterston shore, etc.).
C. confusa (B. & N.).—Mod. c. and gen. dist.
C. porcellanea, Brady.—More or less gen. dist.
C. macallana (B. & R.).—Off Fairlie, 5 f. (A. M. N.): off Cumbrae; Roseneath (R.).
C. tenera, Brady.—C. (R.). Not very c. (S.).
C. (?) semipunctata, Brady.—L. Fyne, off Tarbert; bet. the Cumbraes, 25 f. (A. M. N.): off Sanda I., mod. r.
C. badia, Norman.—Kames B.; Cumbrae; off Bute (R.).
C. crispata, Brady.—Freq. (R.): L. Tarbert; Campbeltown L., etc.
C. gibbosa (B. & R.).—Kames B., Cumbrae; L. Gilp; L. Tarbert (B. & N.): Brackish pools nr. Langbank (S.).
C. rubida, Brady.—Kames B., Cumbrae; Clachland Pt., Arran (B. & N.).
C. albomaculata, Baird.—Freq. (R.): mod. c. and gen. dist.
C. robertsoni, Brady.—Freq. (R.): Tarbert Bank; Campbeltown L., etc.
C. convexa, Baird.—Freq. and gen. dist.
C. marginata, Norman.—Off Tarbert, L. Fyne, 25 f. (A. M. N.): Tarbert Bank; rather r. (S.).
C. limicola, Norman.—Girvan (R.): Cumbrae; L. Cumbrae (A. M. N.): Campbeltown L. (S.).
C. cunieformis, Brady.—Freq. (R.).
C. navicula, Norman.—Off Port Glasgow 1892; off Sanda I. 1897.
C. cluthae, B. C. & R.—L. Fyne, 20 f. (B. & N.).
C. villosa (G. O. Sars).—C. and gen. dist.
C. pulchella, Brady.—Kames B., Cumbrae; Kilchattan B. (B. & N.).
C. quadridentata, Baird.—Off Cumbrae (R.): Tarbert Bank, 25 f.; not r.
C. emaciata, Jones.—Freq. (R.): Tarbert Bank, 25 f.
C. tuberculata (G. O. Sars).—Gen. dist., c.
C. emarginata (G. O. Sars).—Off Inveraray (R.): Tarbert Bank (prob. sub-fossil), not c. (S.).
C. finmarchica (G. O. Sars).—Off Inveraray (R.): Campbeltown L. 1897 (S.).
C. angulata (G. O. Sars).—Various places in F. of Clyde and L. Fyne (B. & N.).
C. dunelmensis (Norman).—Mod. freq. and gen. dist.
C. antiquata, Baird.—Various places in F. of Clyde (B. & N.): Tarbert Bank; Campbeltown L.
C. whitei (Baird).—Kilchattan B.; Campbeltown L. (R.): Tarbert Bank.
C. jonesii (Baird).—Gen. dist.; not r.
Limnicythere inopinata (Baird).—Not unc. in Fresh-water Lochs and ponds.

- L. sancti-patricii* (*B. & R.*).—L. Libo ; side of Bishop L. (R.).
Cytheridea elongata, *Brady*.—Freq. (R.) : nr. Fort Matilda ; Campbeltown L., &c.
C. papillosa, *Bosquet*.—C. (R.) : off Greenock ; Roseneath ; Rothesay (B. & N.) : L. Fyne.
C. punctillata, *Brady*.—C. (R.) : off Roseneath ; Rothesay ; Inveraray ; Skipness ; Tarbert (B. & N.).
C. torosa (*Jones*).—Langbank ; Pt. Glasgow ; Greenock (R.). A brackish water species.
C. lacustris (*G. O. Sars*).—Glasgow and Paisley Canal (W. of Pollokshields) ; Clyde, nr. Langbank (R.) : Pilmuir Dam (J. Smith) : L. Lomond (B. & N.).
C. (?) subflavescens, *Brady*.—L. Fyne ; nr. Rothesay ; bet. the Cumbraes (B. & N.).
Eucythere declivis (*Norman*).—Freq. and gen. dist.
Krithe bartonensis (*T. R. Jones*).—C. (R.). Clyde nr. Greenock ; L. Fyne ; off Rothesay, &c. (B. & N.).
Loxoconcha impressa, *Baird*.—C. and gen. dist.
L. guttata (*Norman*).—Freq. and gen. dist. (B. & N.).
L. viridis (*Müller*).—Brackish pools at Inverkip (1880) ; Campbeltown L. (1897) (S.). Irvine sands (J. Smith).
L. multifora (*Norman*).—C. (R.). L. Fyne (A. M. N.) : Cumbrae and Rothsay B. (B. & N.) : Campbeltown L. (S.).
L. pusilla, *B. & R.*—Greenock ; Cumbrae (R.) : Campbeltown L. (S.).
L. tamarindus, (*Jones*).—C. and gen. dist.
L. fragilis, *G. O. Sars*.—Clyde nr. Greenock and Pt. Glasgow (R.) : [probably fossil, not unc. in Artic shell beds bet. these two places].
Xestoleberis aurantia (*Baird*).—Kames B., Cumbrae (R.) : Tarbert Bank ; E. L. Tarbert ; Campbeltown L. ; not r.
X. depressa, *G. O. Sars*.—Gen. dist. throughout estuary.
Cytherura gibba (*Müller*).—Freq. (R.), but mostly in brackish water.
C. cornuta, *Brady*.—Gen. dist. and mod. c.
C. sella, *G. O. Sars*.—Gen. dist. and mod. c.
C. acuticostata, *G. O. Sars*.—Gen. dist. but scarcely so c. as the last.
C. striata, *G. O. Sars*.—Gen. dist. Freq. (R.).
C. angulata, *Brady*.—Rock-pools, Arran (A. M. N.) : L. Fyne ; Campbeltown L. ; not so c. as some other *Cytherurae* (B. & N.).
C. undata, *G. O. Sars*.—Gen. dist. freq. throughout estuary.
C. producta, *Brady*.—F. of Clyde (B. & N.) : L. Fyne, not c.
C. nigrescens (*Baird*).—Gen. dist. throughout estuary.
C. simplex, *B. & N.*—St. Ninian's B., Bute ; off Fairlie (B. & N.) : Campbeltown L. (S.).
C. concentrica, *B. C. & R.*—Off Fairlie (A. M. N.). Appears to be r.
C. similis, *G. O. Sars*.—L. Fyne ; Rothesay B. ; off Fairlie (B. & N.) : Campbeltown L. (F. S. G.).
C. fulva, *B. & R.*—Off Greenock ; L. Fyne ; F. of Clyde (B. & N.) : Campbeltown L. (F. S. G.).
C. clathrata, *G. O. Sars*.—Callum's B., Bute (R.) : Tarbert Bank, not c. (F. S. G.).
C. cellulosa, *Norman*.—Gen. dist. ; freq. (very small).
C. bodotria, *T. Scott*.—Greenock Bank, off Pt. Glasgow, 1892 (S.).
Cytheropteron latissimum (*Norman*).—Gen. dist. throughout estuary.
C. nodosum, *Brady*.—F. of Clyde, freq. (R.) : off Tarbert (A. M. N.) : Campbeltown L. (F. S. G.).
C. inflatum, *B. C. & R.*—L. Fyne (A. M. N.).
C. punctatum, *Brady*.—Inverkip ; Cumbrae ; L. Riddon ; off Girvan, r. (R.) : off Tarbert, 25 f. (A. M. N.).

- C. alatum**, *G. O. Sars*.—Off Tarbert, (A. M. N.) ; Fintry B. ; Kilchattan B. ; Rothesay B. (B. & N.) : off Sanda I. (F. S. G.).
- C. angulatum**, *B. & R.*—Off Tarbert (A. M. N.) : Roseneath ; Cumbrae ; Bute (R.) : off Sanda I. (F. S. G.).
- C. depresso**, *B. & N.*—Campbeltown L. (1897) : r. in estuary (S.).
- C. humile**, *B. & N.*—Off Fort Matilda, Greenock (B. & N.) : U. L. Fyne ; Kilbrennan Sd. (S.).
- Bythocythere constricta**, *G. O. Sars*.—Off Roseneath, etc. ; L. Long ; L. Fyne (B. & N.).
- B. turgida**, *G. O. Sars*.—Off Roseneath ; Rothesay B. ; Kilchattan B. (B. & N.) : off Tarbert (S.).
- B. simplex** (*Norman*).—Gen. dist., but not very c.
- Pseudocythere caudata**, *G. O. Sars*.—Mod. freq. throughout estuary, and L. Fyne.
- Selcrochilus contortus** (*Norman*).—Gen. dist. throughout estuary.
- Cytherideis subulata**, *Brady*.—Cumbrae ; Bute ; Ardrossan, r. (R.) : Campbeltown L. (F. S. G.).
- Cytherois fischeri** (*G. O. Sars*).—Gen. dist., but more freq. inshore amongst algae, Inverkip ; Campbeltown L. (F. S. G.).

Fam. Paradoxostomatidae.

- Paradoxostoma variabile** (*Baird*).—C. and gen. dist.
- P. ensiforme** *Brady*.—Mod. c. ; but more so off shore than in shall. w.
- P. abbreviatum**, *G. O. Sars*.—Gen. dist., but not very plentiful.
- P. obliquum**, *G. O. Sars*.—Cumbrae ; Bute ; Arran, r. (R.) : Off Tarbert (F. S. G.).
- P. normani**, *Brady*.—Campbeltown L. 1897 (S.). Somewhat r. in Clyde.
- P. pulchellum**, *G. O. Sars*.—L. Goil ; Cumbrae ; Arran ; Ardrossan, r. (R.) : L. Fyne, 1885 (S.).
- P. hibernicum**, *Brady*.—Cumbrae ; Bute ; Arran ; Campbeltown, r. (R.).
- P. arcuatum**, *Brady*.—Off Pt. Glasgow, r. (S.).
- P. orcadense**, *B. & R.*—White B., Cumbrae ; off Glensauno, r. (R.).
- P. hodgei**, *Brady*.—Off Callum's Bay, Bute (R.) : off Tarbert, 25 f. (A. M. N.).
- P. flexuosum**, *Brady*.—Freq. in Clyde (R.) : L. Fyne ; Campbeltown L. (F. S. G.).
- P. affine**, *T. Scott*.—Off Inveraray, 25-40 f. (A. M. N.).
- Machaerina tenuissima** (*Norman*).—Off Skelmorlie ; Fairlie ; nr. Cumbrae ; Kilchattan B. (B. & N.) : off Tarbert (F. S. G.).

Fam. Asteropidae.

- Asterope mariae** (*Baird*).—Off Tarbert (A. M. N.) : Cumbrae (R.) : U. L. Fyne ; East L. Tarbert.
- A. teres** (*Norman*).—Lamlash B. (A. M. N.) : Cumbrae (R.) : E. L. Tarbert.

Fam. Cypridinidae.

- Philomedes brenda** (*Baird*).—Nr. Arran (E. side) 40 f. 1896 (S.).
- P. interpuncta** (*Baird*).—Gen. dist., in surface and bottom tow-nets, as well as dredge.
- P. macandrei** (*Baird*).—Nr. Cumbrae (R.).

Fam. Polycopidae.

- Polycopis orbicularis**, *G. O. Sars*.—Gen. dist. throughout estuary.
- P. punctata** *G. O. Sars*.—“Some specimens dredged in L. Fyne are doubtfully referable to this species” (B. & N.).

Fam. Cytherellidae.

- Cytherella serrulata**, *Brady and Norman*.—Nr. Sanda I., 1896 (S.).

ORDER COPEPODA.

ARRANGEMENT AND NOMENCLATURE FOLLOWED.

- Brady, Prof. G. S. (1) *Monog. of the free and semi-parasitic Copepoda of the Brit. Islands (Ray Society, 1878-80).*
 (2) *Revision of the Brit. species of the fresh-water Cyclopidae and Calanidae. Nat. Hist. Trans., N.D. and N.C., vol. xi., Pt. 1 (1891).*
 Giesbrecht, Dr. W. (1) *Die Pelagischen Copepoden des Golfs von Neapel.*
 (2) *Die Asterocheriden des Golfs von Neapel.*
 Canu, Dr. Eugene, *Les Copepodes du Boulonnais.*
 Schmeil, Dr. Otto, *Deutschlands freilebende süßwasser Copepoden.*
 List of species from Irish Sea (*Brit. Assoc. Rep.*, Sect. D, Liverpool, 1896).
 Drs. W. Giesbrecht and O. Schmeil, *Copepoda Gymnoplea*, Das Tierreich (6 Lieferung).
 Dr. Basset Smith, *A systematic description of the parasitic copepoda found on fishes (Proc. Zool. Soc. Lond., 1899).*
 Scott, T. & A., *Revis. of Brit. Copep. belonging to the Gen. Bradya and Ectinosoma (Tr. Lin. Soc., 2nd ser.; Zool., vol. vi., Pt. 5 (1896))*: papers in *Ann. and Mag. Nat. Hist.*, and *Ann. Scot. Nat. Hist.* (1892-98).
 Scott, T., papers in *F.B. Reports*, 1886-1900.

Fam. Calanidae.

- Calanus finmarchicus* (*Gunn*).—Throughout estuary, surface and bottom.
Paracalanus parvus, *Claus*.—Off Minard, U. L. Fyne; Kilbrennan Sd.; Ayr B.; off Sanda I. etc., not c.
Pseudocalanus elongatus, *Boeck*.—Gen. dist. and mod. c. (surface and bottom townets).
Stephus gyrans (*Giesbrecht*).—L. Gair (U. L. Fyne), r., dredged.
S. minor, *T. Scott*.—Otter Spit (U. L. Fyne); off Sanda I., r., dredged.
S. fultoni, *T. Scott*.—Otter Spit (U. L. Fyne); Kilbrennan Sd., r., dredged.
Pseudocyclopia crassicornis, *T. Scott*.—Otter Spit (U. L. Fyne), r., dredged.
P. caudata, *T. Scott*.—Otter Spit; Ayr B.; Ballantrae Bank, r., dredged.
Bradyidius armatus (*Vanhoffen*).—Gen. dist. throughout estuary.
Euchaeta norvegica, *Boeck*.—U. L. Fyne, very c.; much less so in seaward part of estuary; chiefly in bottom townet.
Scolecithrix hibernica, *A. Scott*.—U. L. Fyne; d. w. E. of Arran; off Davaar I.; nr. Ailsa Craig (in bottom townet).
S. pygmaea, *T. Scott*.—U. L. Fyne; Kilbrennan Sd.; Whiting B.; nr. Ailsa Craig (in bottom townet), not c.

Fam. Centropagidae.

- Centropages typicus*, *Kr.*.—Gen. dist. and mod. freq.
C. hamatus (*Lillj.*).—Gen. dist., c.
Isias clavipes, *Boeck*.—Off Cumbrae; off Portincross (Brady); Machrie B. (Brook); off Ailsa Craig.
Diaptomus gracilis, *G. O. Sars*.—C. in fresh-water lochs, as L. Doon, Ayrshire; L. Lomond, etc.
D. hircus, *Brady*.—Tangy L., nr. Campbeltown.
D. laciniatus, *Lillj.*.—L. Doon, Ayrshire; freq. during summer.
Temora longicornis (*Müller*).—Gen. dist. throughout estuary.
Eurytemora velox (*Lillj.*).—Cumbrae (Brady); pond at Rutherford (J. Steel); Fossil Marsh; marshy ground at Langbank; brackish pools at Hunterstone.
Metridia lucens, *Boeck*.—E. L. Tarbert (Calderwood); Largabruach, U. L. Fyne; nr. Ailsa Craig.
Paramisophria cluthae, *T. Scott*.—Largabruach, U. L. Fyne; Tarbert Bank, L. L. Fyne, r.

Fam. Candaciidae.

Candacia pectinata, *Brady*.—Whiting B.; mid-channel E. of Pladda I.; off Sanda I., not very c.

Fam. Pontellidae.

Labidocera wollastoni (*Lubbock*).—Machrie B. (Brook): nr. Ailsa Craig; off E. of Arran; not c.; 12-26 f.

Anomalocera patersoni (*Temple*).—Gen. dist. sometimes mod. c.

Parapontella brevicornis (*Lubbock*).—E. L. Tarbert (Calderwood): Ayr B.; Machrie B.; off Davaar I.; nr. Ailsa Craig.

Acartia clausii, *Giesbrecht*.—Gen. dist. and mod. c. throughout estuary.

Fam. Misophriidae.

Misophria pallida, *Boeck*.—Portincross (Brady): Largabruach, U. L. Fyne; Otter Spit, r.

Fam. Cyclopidae.

Oithona helgolandica, *Claus*.—Gen. dist., mod. c.

Cyclopina littoralis, *Brady*.—Largabruach, and Head of U. L. Fyne; not c.

C. gracilis, *Claus*.—Loch Gair; off Inveraray, Hunterstone; not c.

Cyclops strenuus, *Fischer*.—C. in most fresh-water lochs.

C. leuckarti, *Claus*.—L. Lomond, not c.

C. dybowskii, *Lande*.—L. Lomond—shore S. of Balmaha.

C. bicuspidatus, *Claus*.—L. Doon; Achy L. (nr. Campbeltown); Possil Marsh.

var. *lubbockii*, *Brady*.—Brackish pools at Langbank.

C. languidus, *G. O. Sars*.—L. Doon, Ayrshire; not c.

C. vernalis, *Fischer*.—L. Lomond; L. Doon.

C. bisetosus, *Rehberg*.—L. Lomond (shore); pools at Langbank; Garry L., Ailsa Craig.

C. viridis (*Jurine*).—L. Lomond; L. Fad; L. Doon; Park L., nr. Campbeltown; mod. c.

C. fuscus (*Jurine*).—L. Arklet; Park L.

C. albidus (*Jurine*).—L. Lomond; L. Doon; pond at Castlemilk; Possil Marsh.

C. serrulatus, *Fischer*.—C. and gen. dist. in lochs and ponds throughout the district.

C. macrurus, *G. O. Sars*.—L. Arklet.

C. affinis, *G. O. Sars*.—Achy L., nr. Campbeltown.

C. nanus, *G. O. Sars*.—Dhu Lochan, nr. L. Lomond; L. Doon.

C. fimbriatus, *Fischer*.—Gen. dist. and mod. c.

C. phaleratus, *Koch*.—Bardowie L.; not c.

C. aequoreus, *Fischer*.—Brackish pools at Langbank and Hunterstone.

Thorellia brunnea, *Boeck*.—Gen. dist. throughout estuary, freq.

Fam. Ascidicolidae.

Notodelphys allmani, *Thorell*.—E. L. Tarbert; Tarbert Bank (inside large Ascidians, as are most of the species in this family).

N. agilis, *Thorell*.—E. L. Tarbert; Tarbert Bank 20-25 f. (F. S. G.).

N. prasina, *Thorell*.—E. L. Tarbert; Tarbert Bank; Kilbrennan Sd.

Doropygus porcicauda, *Brady*.—E. L. Tarbert; off Largabruach, U. L. Fyne; Tarbert Bank (F. S. G.).

D. pulex, *Thorell*.—In ascidians, Tarbert Bank; nr. Millport in *Corella parallelogramma* (Gray).

D. (?) gibbosa, *Thorell*.—In ascidians Tarbert Bank (F. S. G.).

Botachus cylindratus, *Thorell*.—Same localities as the last; Whiting B.

Notopterophorus papilio, *Hesse*.—In large Ascidiants in E. L. Tarbert; not unc.; Kilbrennan Sd. 29/10/96 (F. S. G.).

Ascidicola rosea, Thorell.—In large Ascidiarians dredged nr. head of L. Fyne ; Tarbert Bank (F. S. G.).

Enterocola fulgens, van Beneden.—In the intestines of small Ascidiarians dredged at Tarbert Bank (F. S. G.).

Gunenotophorus globularis, Costa.—In material dredged nr. Sanda I. 12/98 (F. S. G.).

Botryllophilus (?) *ruber*, Hesse.—Associated with *Botryllus* sp., dredged Tarbert Bank, 1899 ; not very c. (F. S. G.).

Lamippi proteus, Claparede.—Parasitic in *Aleyonium digitatum* in various parts of Clyde area.

Fam. Harpacticidae.

(NOTE.—The Copepods belonging to this family—both marine and freshwater—are for the most part found on or near the bottom.)

Longipedia coronata, Claus.—Freq. and gen. dist.

L. minor, T. & A. Scott.—Largabruach, U. L. Fyne ; off Sanda I. (with ova). *Canuella perplexa*, T. & A. Scott.—Ayr B. ; Cumbrae, bet. tide marks ; Inverkip, in shore pools.

Ectinosoma atlanticum (B. & R.).—Gen. dist., sometimes abund. in tow-net.

E. sarsi, Boeck.—Gen. dist., and not unc.

E. melaniceps, Boeck.—Gen. dist., not r.

E. herdmanni, T. & A. Scott.—L. Gair and Largabruach, U. L. Fyne ; shore at Hunterston.

E. curticone, Boeck.—Dubh L., nr. Inveraray ; Largabruach ; shore pools at Hunterston ; Ayr B.

E. normani, T. & A. Scott.—Ayr B., r.

E. erythrops, G. S. Brady.—Otter spit, U. L. Fyne, not c.

E. gracile, T. & A. Scott.—Shore pools at Inverkip ; dredged nr. Sanda I., r.

E. pygmaeum, T. & A. Scott.—Tarbert Bank 20 f., Kilbrennan Sd., r.

Bradya typica, Boeck.—Gen. dist., not very c.

B. minor, T. & A. Scott.—Shore at Hunterston ; Ayr B.

B. elegans, T. & A. Scott.—L. Gair and Largabruach, U. L. Fyne.

B. hirsuta, T. & A. Scott.—Ayr B., r.

B. similis, T. & A. Scott.—Largabruach ; Kilbrennan Sd. ; Ayr B.

Zosima typica, Boeck.—Lowburn, and Largabruach, U. L. Fyne ; Kilbrennan Sd., not r.

Tachidius brevicornis (G. F. Muill.).—Cumbrae, nr. Millport (Shore) ; Hunterston in brackish water pools.

T. littoralis, Poppe.—Shore pools, Dumbarton and Hunterston.

Robertsonia tenuis (B. & R.).—Rothesay B.

Amymone sphaerica, Claus.—E. L. Tarbert ; Kilbrennan Sd. ; Whiting B. ; nr. Sanda I.

A. nigrans, T. & A. Scott.—Otter Spit, L. Fyne 12/1/99.

Stenhelia hispida, Brady.—Off Portincross, Ayrshire 15 f. (Brady).

S. ima, Brady.—Off Callum's Hole, Bute ; off Glen Sannox (Brady) ; E. L. Tarbert ; off Inveraray.

S. intermedia, T. Scott.—Kilbrennan Sd.

S. hirsuta, I. C. Thompson.—Tarbert Bank ; Campbeltown L.

Ameira longipes, Boeck.—Off Portincross, Ayrshire (Brady).

A. longicaudata, T. Scott.—U. L. Fyne ; Kilbrennan Sd., off Sanda I.

A. exilis, T. & A. Scott.—Shore pools at Hunterston.

A. exigua, T. Scott.—Kilbrennan Sd.

A. longiremis, T. Scott.—L. Gair ; Kilbrennan Sd.

Jonesiella spinulosa (B. & R.).—Largabruach ; Kilbrennan Sd. ; Campbeltown L.

J. fusiformis (B. & R.).—Tarbert Bank 20 f. ; off Sanda I.

- Delavalia palustris*, *Brady*.—Brackish pools, Dumbarton ; Hunterston.
D. reflexa, *B. & R.*.—Nr. Sanda I., rather r.
D. robusta, *B. & R.*.—Largabruach ; Kilbrennan Sd., nr. Sanda I.
D. mimica, *T. Scott*.—L. Gair ; Kilbrennan Sd., Ayr B.
D. aemula, *T. Scott*.—Ayr B. ; Campbeltown L., not c.
D. giesbrechti, *T. & A. Scott*.—Ayr B. ; Campbeltown L.
 var . . . ?—Brackish-water pools, Hunterston.
Pseudotachidius coronatus, *T. Scott*.—Off Lowburn, Hd. of L. Fyne ; off Skate I., 105 f. ; bet. Arran and Heads of Ayr, 40 f.
Cervinia bradyi, *Norman*.—D. w. E. of Arran, very r.
Canthocamptus staphylinus (*Jurine*).—C. in fresh water lochs, ponds, ditches, &c.
C. palustris, *G. S. Brady*.—Brackish pools, Dumbarton ; Langbank ; Hunterston ; Dubh L., nr. Inveraray.
C. parvus, *T. & A. Scott*.—Brackish pools, Hunterston, not c.
C. minutus, *Claus*.—L. Lomond ; L. Doon ; Park L. and Achy-L., nr. Campbeltown.
C. inornatus, *T. Scott*.—L. Arklet ; Park L., not c.
C. schmeilii, *Mrazek*.—L. Lomond ; L. Doon ; Park L.
C. hirticornis, *T. Scott*.—Brackish pools at Langbank.
C. cuspidatus, *Schmeil*.—L. Fad, Bute, r.
C. crassus, *G. O. Sars*.—Freq. in most fresh-water lochs and ponds.
C. pygmaeus, *G. O. Sars*.—More or less freq. in lochs, ponds, ditches.
C. zschokkei, *Schmeil*.—Dhu Lochan, nr. L. Lomond ; L. Fad, Bute ; Black L. and Park L., nr. Campbeltown.
Moraria anderson-smithi, *T. & A. Scott*.—L. Arklet ; L. Lomond ; L. Doon ; Garry L. (Ailsa Craig) ; Black L., nr. Campbeltown.
M. brevipes (*G. O. Sars*).—L. Doon, not c.
M. poppei (*Mrazek*).—L. Fad, Bute ; Shore pools at Hunterston, r.
Maraenobiotus vejvodovskii, *Mrazek*.—L. Doon, amongst shore vegetation.
Psyllocamptus fairliensis, *T. Scott*.—Shore pools bet. Fairlie and Hunterston.
 ✕ *Mesochra lilljeborgii*, *Boeck*.—Brackish-water pools at Hunterston.
 ✕ *M. spinicauda*, *T. & A. Scott*.—Shore pools, nr. Millport.
Neobradyla pectinifer, *T. Scott*.—Ballantrae Bank, r.
Tetragoniceps consimilis, *T. Scott*.—Ballantrae Bank ; Otter Spit, L. Fyne.
T. incertus, *T. Scott*.—Dredged bet. Sanda and Bennan Hd. 1896.
T. macronyx, *T. Scott*.—Dredged, Ballantrae Bank, 18 f.
Diosaccus tenuicornis (*Claus*).—Cumbrae, surface net ; Portincross, dredged (Brady) : E. L. Tarbert ; Kilbrennan Sd.
Laophonte serrata (*Claus*).—E. L. Tarbert ; Kilbrennan Sd. r.
L. horrida (*Norman*).—Off Cumbrae and Portincross (Brady) : U. L. Fyne ; Kilbrennan Sd., not unc.
L. thoracica, *Boeck*.—Off Inveraray and Largabruach, U. L. Fyne.
L. similis (*Claus*).—E. L. Tarbert ; L. Fyne (Calderwood). Shore pools, Hunterston.
L. curticauda, *Boeck*.—Amongst weeds at L. Cumbrae (Brady) : shore pools Hunterston.
L. longicaudata, *Boeck*.—Otter Spit ; Kilbrennan Sd.
L. lamellifera (*Claus*).—Portincross, Ayrshire ; off Cumbrae and Bute (Brady) ; Otter Spit ; Ayr B.
L. intermedia, *T. Scott*.—Shore pools Hunterston.
L. hispida (*B. & R.*).—Off Glen Sannox ; off Cumbrae and Bute (Brady) : Largabruach, U. L. Fyne ; Ayr B.
L. littoralis, *T. & A. Scott*.—Shore pools Hunterston.
L. mahammed, *Richard*.—Brackish pools Langbank.
L. depressa, *T. Scott*.—Tarbert Bank 25 f. not c.

- L. simulans*, *T. Scott*.—Off Inveraray; Kilbrennan Sd.; freq. in crevices of decaying wood.
- L. denticornis*, *T. Scott*.—Machrie B.; Kilbrennan Sd.
- Pseudolaophonte spinosa* (*J. C. Thompson*).—Dredged nr. Otter Spit 15-18 f., Jany. 99.
- Laophontodes typicus*, *T. Scott*.—Nr. Otter Spit; Carradale B.
- L. bicornis*, *A. Scott*.—Machrie B., not c.
- Pontopolites typicus*, *T. Scott*.—L. Gair; Largabruach, U. L. Fyne; Kilbrennan Sd.
- Normanella dubia* (*B. & R.*).—L. Gair; Largabruach; Machrie B.; Carradale B.; Ayr B.
- Cletodes limicola*, *Brady*.—Kilbrennan Sd.
- C. longicaudata*, *B. & R.*.—Off Portincross, Ayrshire (Brady): E. L. Tarbert; L. Gair; Largabruach; Kilbrennan Sd.; Campbeltown L.
- C. propinquua*, *B. & R.*.—Shore pools Hunterston.
- C. linearis* (*Claus*).—r.; E. L. Tarbert; bet. Sanda and Bennan Hd.
- C. similis*, *T. Scott*.—Otter Spit.
- C. curvirostris*, *T. Scott*.—Carradale B.; Campbeltown L.
- C. lata*, *T. Scott*.—Kilbrennan Sd.; Campbeltown L.; nr. Davaar I.
- C. irrasa*, *T. Scott*.—Kilbrennan Sd.
- C. tenuipes*, *T. Scott*.—r.; Carradale B., off Sanda I.
- C. hirsutipes*, *T. Scott*.—Carradale B.
- Itunella tenuiremis* (*T. Scott*).—Nr. Lowburn and Largabruach, U. L. Fyne; shore pools Hunterston.
- Leptopsyllus minor*, *T. Scott*.—Shore pools, nr. Millport, not c.
- L. herdmani*, *I. C. Thompson & A. Scott*.—Shore pools, nr. Millport, not c.
- Enhydrosoma curvatum*, *B. & R.*.—Off Portincross; Cumbrae; Bute (Brady): L. Gair; Ayr B.
- Pseudocletodes vararensis*, *T. & A. Scott*.—Nr. Sanda I. twice.
- Nannopus palustris*, *Brady*.—Brackish pools Inverkip and Hunterston.
- Platyhelipus littoralis*, *Brady*.—Dumbarton; Ayr B.; Carradale B.; Hunterston, shore pools.
- Huntemannia jadensis*, *S. A. Poppe*.—Shore pools nr. Millport; Inverkip.
- Heteropsyllus curticaudatus*, *T. Scott*.—Carradale B., r.
- Dactylopus pectinatus*, *T. Scott*.—Nr. head of L. Fyne (18 f.).
- D. tiboides*, *Claus*.—Off L. Cumbrae, and Portincross (Brady); E. L. Tarbert; Kilbrennan Sd., etc.
- D. similis*, *Claus*.—Off Portincross (Brady); L. Fyne; Whiting B.; Campbeltown L.
- D. stromii* (*Baird*).—Nr. Cumbrae (Brady); Ayr B.; Whiting B., etc.
- D. flavus*, *Claus*.—Off Callum's Hole, Bute (Brady); head of L. Fyne; Otter Spit; Ayr B.
- D. brevicornis*, *Claus*.—Whiting B.; off Sanda I.
- D. minutus*, *Claus*.—E. L. Tarbert (16 f.); off Sanda I.
- Thalestris mysis*, *Claus*.—E. L. Tarbert; Machrie B.
- T. helgolandica*, *Claus*.—Clyde (A. M. N.); Whiting B.; Kilbrennan Sd.
- T. rufocincta*, *Norman*.—Off Cumbrae, and Portincross (Brady); Otter Spit; Whiting B., freq.
- T. harpactoides*, *Claus*.—Shore pools Hunterston.
- T. clausii*, *Norman*.—L. Fyne (Calderwood); off Cumbrae (Brady): Machrie B.; Ayr B.
- T. violascens*, *Claus*.—Nr. the 'Allans' Cumbrae (Brady), r.
- T. hibernica*, *B. & R.*.—Off Cumbrae (Brady), r.
- T. longimana*, *Claus*.—Off Portincross (Brady); Machrie B.
- T. peltata* (*Boeck*).—Otter Spit, rather r.
- T. forcipulus*, *Claus*.—Off Inveraray, and Largabruach, U. L. Fyne; Machrie B.; Shore pools Hunterston, not very r.

- Pseudothalestris pygmaea**, *T. & A. Scott*.—Machrie B.; Ballantrae Bank, r.
Westwoodia nobilis (*Baird*).—Off Portincross, 10-30 f. (Brady); E. L. Tarbert (Calderwood); shore at Hunterston.
- Harpacticus chelifer** (*Müller*).—Gen. dist., freq.
- H. fulvus**, *Fischer*.—Brackish pools, Hunterston.
- H. flexus**, *B. & R.*.—Kames B., Cumbrae, 1-2 f. (Brady); E. L. Tarbert; shore at Hunterston.
- Zaus spinatus**, *Goodsir*.—E. L. Tarbert; L. Fyne (Calderwood); Otter Spit; Machrie B., freq.
- Z. goodsiri**, *Brady*.—Off Portincross, 15 f. (Brady); Otter Spit; off Sanda I.
- Alteutha bopyroides**, *Claus*.—E. L. Tarbert; L. Fyne (Calderwood).
- A. crenulata** (*Brady*).—Off Sanda I.; Ballantrae Bank.
- Eupelta purpurocinctum** (*Norman*).—E. L. Tarbert (Calderwood); Ayr B.; N.-E. of Sanda I. Mod. freq. in Clyde.
- Peltidium purpureum**, *Philippi*.—E. L. Tarbert; off Largabruach; nr. Lowburn, L. Fyne, rather r.
- Porcellidium fimbriatum**, *Claus*.—Cumbrae; Portincross (Brady); E. L. Tarbert (Calderwood); Machrie B. Freq. on fronds of *Laminaria*.
- P. viride** (*Philippi*).—Cumbrae; Portincross (Brady).
- P. subrotundum**, *Norman*.—L. Fyne (M.); Portincross (Brady).
- Idya furcata** (*Baird*).—E. L. Tarbert (Calderwood); Cumbrae; Portincross (Brady); Machrie B.
- I. gracilis**, *T. Scott*.—L. Gair; off Inveraray; off Lowburn (U. L. Fyne); shore pools Hunterston.
- I. longicornis**, *T. & A. Scott*.—E. L. Tarbert; Machrie B.; Campbeltown L.
- I. minor**, *T. & A. Scott*.—Largabruach, U. L. Fyne; Machrie B.
- I. cluthae**, *T. Scott*.—Nr. head of L. Fyne; off Ailsa Craig.
- Scutelidium fasciatum**, *Boeck*.—E. L. Tarbert (Calderwood).
- Cylindropsyllus laevis**, *Brady*.—E. L. Tarbert; Ballantrae Bank.
- C. minor**, *T. Scott*.—Ballantrae Bank 18 f., not c.
- C. fairliensis**, *T. Scott*.—Brackish pools at Hunterston, rather r.
- Leptocaris minutus**, *T. Scott*.—Brackish pools at Hunterston, r.

Fam. Monstrillidae.

Monstrilla danae, *Claparède*.—U. L. Fyne, sometimes freq.; Kilbrennan Sd.

Fam. Corycaeidae.

Corycaeus anglicus, *Lubbock*.—In surface net nr. Ailsa Craig, 29/5/99.

Fam. Lichomolgidae.

- Modiolicola insignis**, *Aurivilius*.—Commensal in shell of *M. modiolus*.
- Lichomolus albens**, *Thorell*.—Dredged nr. Otter Spit.
- L. agilis**, *Leydig*.—Kilbrennan Sd., on branchial plumes of *Doris sp.*
- L. maximus**, *I. C. Thompson*.—Freq. in Clyde; commensal with *Pecten maximus* and *P. opercularis*.
- L. fuciculus**, *Brady*.—Nr. Largabruach, U. L. Fyne.
- L. furcillatus**, *Thorell*.—Off Inveraray, r.
- L. forcipulus**, *Thorell*.—E. L. Tarbert; Tarbert Bank, in branchial chamber of large Ascidians.
- Sabelliphilus sarsi**, *Claparède*.—Kilbrennan Sd.; off Sanda I.; on branchial plumes of *Sabella sp.*
- Hermanella arenicola** (*Brady*).—Off Battery Pt., Cumbrae (Brady); Nr. Otter Spit, 8-15 f.; L. Gilp, 2-3 f.; not c.
- Pseudanthessius liber** (*B. & R.*).—S. of Bute, 16 f. (Brady); nr. Lowburn and Iveraray.
- P. thorellii** (*B. & R.*).—From various parts of estuary.
- Clausia cluthae**, *T. & A. Scott*.—Tarbert Bank, 20-25 f.; Ayr B.

Fam. Asterocheridae.

- Asterocheres lilljeborgi*, Boeck.—Tarbert Bank, 20-25 f., r.
A. echincola (Norman).—Gen. dist., freq. in water passages of *Suberites*.
A. simulans (T. Scott).—L. Fyne and Clyde, r.
A. violaceus (Claus).—Nr. Otter Spit; off Pladda; r.
Dermatomyzon nigripes (B. & R.).—E. L. Tarbert (Calderwood): off Calum's Hole, Bute; Portincross (Brady).
Rhynchomyzon purpurocinctum (T. Scott).—Otter Spit, 8-15 f., Jany./99.
Scotomyzon gibberum (T. & A. Scott).—Nr. head of L. Fyne on *Echinus esculentus*.
Acontiophorus scutatus (B. & R.).—Off Portincross (Brady): Otter Spit; Machrie B.; off Sanda I.
Scottocheres elongatus (T. & A. Scott).—Off Sanda I., r.
Neopontius angularis, T. Scott.—Otter Spit; Kilbrennan Sd., r.
Cribropontius normani (B. & R.).—L. Gair, r.
Bradypontius magniceps (Brady).—Off Portincross (Brady): E. L. Tarbert (Calderwood): nr. Head of L. Fyne.
Dyspontius striatus, Thorell.—Whiting B., r.
Artotrogus orbicularis, Boeck.—Tarbert Bank, 20 f., very r.
Parartotrogus richardi, T. & A. Scott.—Nr. Otter Spit, 15-18 f. (Jany. 99).
Eurynotopsyllus insolens, T. Scott (= *Eurynotus insolens*, T. Scott).—Kilbrennan Sd., 86 (F. S. G.); very r.
Cancerina confusa, T. Scott.—Tarbert Bank, very r., 1899 (F. S. G.).

COPEPODA PARASITA.

CLASSIFICATION ADOPTED: Dr. Basset Smith, "Systematic description of parasitic Copepoda found on fishes," (*Proc. Zool. Soc. London*, April 1899).

Fam. Ergasilidae.

- Bomolochus soleae*, Claus.—Back of *Solea vulgaris*, captured in Clyde (F. S. G.). Freq. in nostrils of Cod.

Fam. Caligidae.

- Caligus curtus*, O. F. Müll.—On various fishes, Cod, Saithe, etc., mod. c.
C. rapax, M.-Edw.—On various fishes, also in tow-net; c. and gen. dist.
C. diaphanus, Nordmann.—Chiefly in gill cavity of Gurnards, *Trigla gurnardus*; *T. lineata*, etc.
Lepeophtheirus pectoralis (O. F. Müll.).—Not unc. under pectoral fins of flat fishes, as the Plaice, *Pleuronectes platessa*, and Flounder, *P. flesus*.
L. thompsoni, Baird.—Freq. on the gills of Turbot, *Bothus maximus*, (F. S. G.).
L. stromi, Baird.—On Salmon nr. Girvan (Duthie, Fishery Officer).
L. (?) obscurus, Baird.—On gills of Brill, *Bothus rhombus*, Clyde (F. S. G.).
Treibius caudatus, Kr.—On *Raia batis* and *R. clavata*, F. of Clyde; freq.

Fam. Diclosteiidae.

- Clavella labracis*, van Ben.—On gills of stripped wrasse *Labrus mixtus*, nr. Ayr, Jan. 1900.
Cycnus pallidus (van Ben.).—On gills of Conger, *Conger niger*, F. of Clyde; not r. (F. S. G.)

Fam. Lernaeidae.

- Lernaea branchialis*, Lin.—Not unc. on gills of Haddocks, Whitings and other Gadoids (F. S. G.).

Fam. Chondracanthidae.

- O**ralien asellinus (*Lin.*).—Not unc. on gills of Gurnards, flat fishes, etc.
Chondracanthus cornutus, (*O. F. Mill.*).—On gills of Plaice, *Pleuronectes platessa*; Clyde (F. S. G.).
C. clavatus, *Basset Smith*.—Mod. freq. on gills of Lemon Soles, *Pleuronectes microcephalus*, F. of Clyde (F. S. G.).
C. soleae, *Kr.*.—On gills of Black Sole, *Solea vulgaris*, F. of Clyde (F. S. G.).
C. flurae, *Kr.*.—On gills of Long Rough Dabs *Drepanopsetta platessoides*, F. of Clyde.
C. merluccii, *Holten*.—Clinging to roof and sides of mouth of Hake *Merluccius merluccius*, Clyde.
C. lophii, *Johnston*.—Inside gill pouches of Angler-fish, *Lophius piscatorius*, Clyde (F. S. G.).
C. zeus, *De la Roche*.—Occasionally on gills of John dory, *Zeus faber*, Clyde (F. S. G.).
C. limanda, *Kr.*.—On gills of common Dab, *Pleuronectes limanda*, Clyde (F. S. G.).
C. ornatus, *T. Scott*.—On gills of a Spotted dragonet, *Cadionymus maculatus*, Clyde 1899 (F. S. G.).

Fam. Lernaeopodidae.

- C**haropinus dubius, *T. Scott*.—On gills and gill-arches of *Raia circularis*, Clyde, Dec., 99 (F. S. G.).
Lerneopoda galei, *Kr.*.—Attached bet. ventral fins of *Galeorhinus galeus* (Duthie); also on *Scyllorhinus caniculus* (F. S. G.).
L. bidiscalis, *W. F. de V. Kane*.—Attached to end of claspers of *Galeorhinus galeus*, Clyde Jany. 1900 (Duthie).
L. ciuthae, *T. Scott*.—On gills of *Raia fullonica*, Clyde, April 1897 (F. S. G.).
Brachiella insidiosa, *Heller*.—On gills of Hake, *Merluccius merluccius*, Clyde (F. S. G.).
B. trigiae, *Claus*.—From gills of *Trigla lineata*, nr. Ailsa Craig 1897.
B. merluccii, *Basset-Smith*.—On gill-rakers of *Merluccius merluccius*, Clyde (F. S. G.).
A. anchorella, *Rugosa*, *Kr.*.—On gills of *Anarrhichus lupus*, not unc. in Clyde (F. S. G.).
A. uncinata, *O. F. Mill.*.—Not unc. on Whiting and other gadoids in Clyde (F. S. G.).
A. stellata, *Kr.*.—On skin of Hake *Merluccius merluccius*, Clyde, 18/12/99 (F. S. G.).

Fam. Choniostomatidae.

- A**spidoecia normani, *Giard and Bonnier*.—Living upon *Erythrops serrata*, G. O. Sars, Clyde, rather r. (S.). (See Dr. H. J. Hansen's Monog. on the Choniostomatidae).

ORDER THORACICA.

Fam. Balanidae.

- B**alanus porcatus, *Costa*.—U. L. Fyne, on shells of *Mytilus* (M.). Gen. dist., freq.
B. crenatus, *Brug.*.—U. L. Fyne (M.). Bet. Corswall and Mull of Kintyre (F. S. G.), freq.
B. balanoides (*Lin.*).—U. L. Fyne (M.). Gen. dist., freq.
B. hameri (*Ascan.*).—U. L. Fyne in 12-20 f., and also on the shore (M.). Occasionally by dredging.

Fam. Verrucidae.

Verruca stromia (*O. F. Müll.*).—Freq. on stones and dead shells.

Fam. Lepadidae.

Scalpellum vulgare, *Leach*.—Attached to Zoophytes; not very r. in Clyde.

ORDER RHIZOCEPHALA.

Fam. Sacculinae.

(NOTE.—Several Sacculines have been obtained in the Clyde; but without a knowledge of their development it would be unsafe to say whether they are distinct or not, and with one or two exceptions I am therefore unable to ascribe them to particular species).

Sacculina carcinii (*Thomps.*).—On abdomen of shore crab, *Carcinus maenas*, not unc. in some parts of the Clyde.

S. sp..—On abdomen of *Galathea dispersa*, Clyde 1895-1896, r. (F. S. G.).

S. sp..—On abdomen of *Munida rugosa* (*Fabr.*). Clyde 1896-1897 (F. S. G.): 1899 (Alex. Patience). Not very r. in seaward part of estuary.

S. sp..—On the abdomen of *Inachus dorsettensis*, dredged bet. Arran and Ayrshire, 20-43 f., 1897 (F. S. G.), apparently r.

S. sp..—On the abdomen of *Stenorhynchus rostratus*, in the trawl net E. of Arran in 20-43 f., 9/97 (F. S. G.).

Peltogaster paguri (*Rathke*).—On abdomen of *Eupagurus bernhardus*, E. L. Tarbert (1886), etc. (S.).

D.
D.
F.=
H.
Ho
S.=

M:
Ro
§=
J.]
(?)

T
Pol
Ap
Eu
E.
He
Lat
Lei
L.
Gad

Eui
Lat
Act
Evi
Hai

H. :

PYCNOGONIDAE.

BY M. LAURIE.

Pycnogonum littorale, *Fabr.*.—Mod. c.

Phoxichilidium sp.—Cumbrae, among sea weed, l. w. (R.).

Pallene brevirostris, *Johnst.*.—L. w. Allans, Cumbrae (R.): Lamlash B. (H.).

Nymphon gracile, *Fabr.*.—L. Cumbrae, Lam Zone and l. w. (R.): Lamlash B. (H.).