

PROCEEDINGS
OF THE
ZOOLOGICAL SOCIETY
OF LONDON.



PART XXII.

1854.

PRINTED FOR THE SOCIETY ;
SOLD AT THEIR HOUSE IN HANOVER SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,
PATERNOSTER ROW.

39. **HELIX SANNIO.** *H. testa sinistrorsa, angustissime umbilicata, turbinato-depressa, solida, oblique costulato-striata, vix nitidula, superne pallide flavicante; spira conoidea, obtusula; anfract. 6 $\frac{1}{2}$, vix convexiusculis, lente accrescentibus, ultimo non descendente, antice rugato, peripheria carinato, subtus convexo, cinnamomeo; apertura diagonali, lunari, intus alba, nitida; perist. incrassato, breviter reflexo, marginibus distantibus, basali substricto, superne sinuato, in laminam triangularem reflexo.*

Diam. maj. 38, min. 32 $\frac{1}{2}$, alt. 21 mill.

Hab. —?

40. **HELIX MANSUETA.** *H. testa umbilicata, depressa, tenuiuscula, striatula et sub lente minutissime granulata, non nitente, nigro-rufa vel cinnamomea; spira brevissime conoidea, vertice subtili; sutura impressa; anfract. 5, convexiusculis, sensim accrescentibus, ultimo antice vix descendente, peripheria obsolete angulato, circa umbilicum apertum, mediocrem saturate fasciato; apertura obliqua, lunari; perist. roseo, breviter expanso, marginibus distantibus, columellari superne dilatato, fornicato-patente.*

Diam. maj. 15 $\frac{1}{2}$, min. 13 $\frac{1}{2}$, alt. 8 mill.

Hab. Moreton Bay, Australia (Mr. Strange).

41. **HELIX MARMATENSIS.** *H. testa umbilicata, depressa, tenuiuscula, vix striatula, albida, superne maculis castaneis fasciatim dispositis ornata; spira plana, vertice subtili vix prominente; anfract. 4 $\frac{1}{2}$, convexis, ultimo rotundato, antice vix descendente, circa umbilicum angustum obsolete compresso; apertura obliqua, rotundato-lunari; perist. tenui, marginibus convergentibus, dextro breviter expanso, columellari superne dilatato, patente.*

Diam. maj. 12, min. 10 $\frac{1}{2}$, alt. 6 mill.

Hab. Marmato, New Granada.

42. **HELIX SCEPASMA.** *H. testa umbilicata, conoidea, solida, ruguloso-striata, opaca, alba; spira regulariter conoidea, vertice acutiusculo; sutura subimpressa; anfract. 7, vix convexiusculis, lente accrescentibus, ultimo antice paululum descendente, peripheria acute carinato, basi circa umbilicum mediocrem, profundum turgido; apertura diagonali, lunari; perist. subincrassato, angulatim reflexo, marginibus vix convergentibus, columellari breviter patente.*

Diam. maj. 20, min. 18, alt. 11 mill.

Hab. —?

4. DESCRIPTIONS OF FOUR NEW SPECIES OF BULIMUS FROM THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

1. **BULIMUS VITELLINUS**, Pfr. *B. testa umbilicata, globoso-conica, solidula, oblique striata, nitidula, luteo-fulva; spira conica, apice obtusa; anfract. 6, rotundatis, ultimo spiram æquante, medio obtuse angulato, circa umbilicum apertum, non pervium subcom-*

presso; apertura diagonali, lunato-ovali, intus submargaritacea; perist. tenui, recto, margine columellari leviter arcuato, sursum valde dilatato, fornicato-patente.

Long. 22, diam. 16 mill.

Hab. Natal.

2. **BULIMUS PICTUS**, Pfr. *B. testa anguste umbilicata, ovato-conica, tenuiuscula, sublaevigata, pallide lutea, fasciis spadiceis, interruptis strigisque sparsis elegantissime subtessellata; spira conica, acutiuscula; sutura levissime crenulata; anfract. 7, convexiusculis, ultimo $\frac{2}{5}$ longitudinis æquante, basi rotundato, infra medium et circa umbilicum fasciis non interruptis signato; apertura parum obliqua, elliptico-ovali; perist. simplice, recto, margine dextro regulariter arcuato, columellari leviter arcuato, sursum dilatato, fornicato-patente.*

Long. 23, diam. 11 mill.

Hab. Peru.

3. **BULIMUS RADIOSUS**, Pfr. *B. testa semi-obtecte perforata, oblongo-conica, tenuiuscula, striatula, lineis impressis, spiralibus, subconfertis decussatula, albida, strigis epidermidis virenti-fulvæ radiata; spira conica, acuta; anfract. $5\frac{1}{2}$, superis planis, sequentibus convexis, ultimo spira paulo breviore, infra medium obsolete angulato; apertura parum obliqua, oblonga; perist. simplice, margine dextro recto, superne arcuato, columellari subtorto, superne dilatato, fornicato-reflexo.*

Long. 21, diam. 10 mill.

Hab. New Hebrides.

4. **BULIMUS GALAPAGANUS**, Pfr. *B. testa compresse umbilicata, ovato-turrita, tenui, striatula, sericea, pellucida, cornea-cerea; spira convexo-turrita, acutiuscula; sutura submarginata; anfract. 8, convexis, ultimo $\frac{1}{3}$ longitudinis subæquante, juxta umbilicum compresso; apertura subverticali, oblongo-ovali; perist. simplice, margine dextro vix expansiusculo, superne arcuato, cum columellari introrsum subtorto, patente angulum formante.*

Long. $15\frac{1}{2}$, diam. 6 mill.

Hab. Galapagos Islands.

5. MONOGRAPHS OF ACTÆON AND SOLIDULA, TWO GENERA OF GASTEROPODOS MOLLUSCA WITH DESCRIPTIONS OF SEVERAL NEW SPECIES FROM THE CUMINGIAN COLLECTION.
BY ARTHUR ADAMS, F.L.S., SURGEON R.N.

Genus ACTÆON, Montfort.

Head in the form of a quadrate disk, bilobed in front, with broad posterior tentacular lobes; eyes sessile, on the middle of the head; mantle included; branchial plume single.

Operculum horny, sub-spiral.