

(in) SIPHONOSTOMATOIDA
Vanbenedenia, valid.
Malm, 1860.

Hernaeopodidae.
Hernaeopodoida
V. kröyeri.

Syn Vanbenedenia, Malm, 1860

To Prof. P. J. Van Beneden.

Head short. Broader than long, not-
markedly attenuated, but plainly sepa-
rated from body. Body elongated,
indistinctly segmented: no anal laminae.
Second max'ds short and thick.

Egg-cases long & filiform: as long as the body.
Suctorial mouth elongate, obtusely conical, wholly
covered above by the thoracic legs, armed at the base
below with the 2^o max'ds. Thoracic legs short-
approximated their entire length, scarcely movable.
Male unknown - (Malin, 1860, p. 620).

VANBENEDENIA MALM, 1860

- CASTRO ROMERO + BAEZA KUROKI 1986 c: 232

- KABATA 1992: 124, DIAGNOSIS

Vanbenedenia chimaerae (Heegaard, 1962)

-- -- Kabata, 1964, p. 254-267.

--, Kabata, 1968, p. 505 see Tracheliastes chimaerae
Heegaard, 1962

--, Kabata, 1970, p. 42

--, Kabata & Cousens, 1972, passim.

Lernaeopodidae.

Vaubenedenia Kröyeri, Malin, ♂. ♀.

- —. Malin, 1860, p. 620. (no figures)
- —. Malin, 1863, p. 114, plate I.
- —. Olsson, 1869, p. 39: pl. II, figs. 16+17. ♂.
- —. Benedeen, 1870, p. 21.

Host. Dorsal fin of Chimaera monstrosa

Mediterranean, Kattegat.

Coasts of Sweden & Norway.

— — —. Carus, 1885, p. 375.
— — —, B-Smidt. 1899, p. 497.
— — — Wilson, 1915, p. 644.

— — —, Kabata, 1958

— — —, Kabata, 1960, pp. 731-735, figs. 1-3. Faroe Is.

— — —, Tambs-Lyche, 1958

. Off. coast of Western Norway.

--, Kabata, 1970, p. 84

-- KABATA 1988b: 6, ORIGIN OF NAME

-- KABATA 1989: 339

V. KROEYERI, KABATA 1992: 124, FIG. 27 F-H, SYNONYMY,
HOST LIST