

***Axonolaimus spinosus* (BÜTSCHLI)**

Anoplostoma spinosa BÜTSCHLI 1874, p. 37, pl. IV—V, fig. 20 a—c; *Axonolaimus spinosus* (BÜTSCHLI) G. SCHNEIDER 1906, p. 10, fig. 2 a—b; *Axonolaimus spinosus* (BÜTSCHLI) DE MAN 1922, p. 233, fig. 21 a—c; *Axonolaimus spinosus* (BÜTSCHLI) ALLGÉN 1947, p. 16—17

Localities and material. — Falkland Islands. St. ~~42a~~: 1 ♂, 1 ♀; South Georgia, St. ~~22b~~: 3 ♂♂, 7 ♀♀; St. ~~28~~: 1 ♂

In the collections from the above localities there are some males of an *Axonolaimid*, which, in the shape of their head as well as of their tail and their spicules seem to be rather typical representatives of this Northern species.

Geographical distribution. — Norway: Sörflatanger (ALLGÉN 1934), Trondheimsfjord (ALLGÉN 1933), Sweden: West Coast, vicinity of the Zool. Stat. Kristineberg (ALLGÉN 1929), Southern Kattegatt (ALLGÉN 1934, 1947), the Sound (ALLGÉN 1927, 1929, 1935), South coast (ALLGÉN 1934), Finland: Gulf of Finland (G. SCHNEIDER 1906, 1926, 1927), Russia: Bay of Neva (FILIPJEV 1930), Germany: Bay of Kiel (BÜTSCHLI 1874).