

### **Chromadora nudicapitata BASTIAN**

*Chromadora nudicapitata* BASTIAN 1865, p. 168, pl. XIII, fig. 230—232; *Chromadora natans* BASTIAN 1865, p. 168, pl. XIII, fig. 236—238

Localities and material. — Fuegian Archipelago, St. 64: 1♂, 1♀; Falkland Islands, St. 40: 1♀; St. 46: 3♀♀; St. 51: 1♀; South Georgia, St. 18: 1♀; St. 22c: 7♂♂, 8♀♀, 1 juv.; St. 22d: 9♂♂, 9♀♀; St. 28: 4♀♀.

All the specimens, recorded from the above localities, are in every respect typical representatives of this since long known North-Sea Species. The males have consequently 5 preanal papillae in a typical arrangement.

Geographical distribution. — *Norway*: Oslofjord (BÜTSCHLI 1874, ALLGÉN 1931), Herdla (ALLGÉN 1932), Trondheimsfjord (ALLGÉN 1933), Tarva (ALLGÉN 1934), Hitra (Hestnes) and Vega (ALLGÉN 1940, 1943), Marøy-Rörvik (ALLGÉN 1946a), *Sweden*: West Coast, vicinity of the Zool. Stat. Kristineberg (ALLGÉN 1929), South Kattegatt (ALLGÉN 1934, 1947), the Sound (ALLGÉN 1935), *Germany*: North Frisian Islands: Amrum (SCHULZ 1936, 1938), *North Sea* and Channel (DE MAN 1888), *Holland*: Coast of Zeeland (DE MAN 1906, 1907), Zuider Sea (DE MAN 1922), *Belgium* (SCHUURM. STEKH. and DE CONINCK 1933), *Brit. Islands*: England (BASTIAN 1865), Ireland (SOUTHERN 1914), *Mediterranean*, Western Part, Coast of France: Cette (ROUVILLE 1904), *Adriatic Sea*: Quarnero (DADAY 1901), Eastern Part: Alexandria (SCHUURM. STEKH. 1943), *United States of America*: California (ALLGÉN 1947); Coast of Chile (GERLACH 1953), *South Sea*: Campbell Island (ALLGÉN 1932).

The male from St. 64 — on the whole a typical representative of the species under consideration ( $L = 0,697$  mm;  $a = 25,82$ ;  $\beta = 6,97$ ;  $\gamma = 10,25$ ) — differs from the other males in the presence of only 2 preanal papillae. The last papilla is situated  $54 \mu$  in front of the anus, the anterior papilla  $27 \mu$  still longer in front. Probably the two first papillae and the hindmost papilla of the 5 papillae in typical males are not developed in this male.

## List of Localities

- St. 1. Off the Coast of Uruguay. Black-grey clay.  $33^{\circ} 0' S$ . —  $51^{\circ} 10' W$ . 80 m. 12. 12. 1901.  
Number of species found: 8; Number of specimens found: 21.
- St. 2. Off the Coast of North Argentina.  $37^{\circ} 15' S$ . —  $56^{\circ} 8' W$ . Sand-mixed gravel. 100 m. 23. 12. 1901.  
Number of species found: 12; Number of specimens found: 26.
- St. 3. Fuegian Archipelago.  $54^{\circ} 43' S$ . —  $64^{\circ} 8' W$ . Rubble stones and gravel. 36 m. 6. 1. 1902.  
Number of species found: 20; Number of specimens found: 37.
- St. 5. Graham Region. S. East of the Seymour Sound.  $64^{\circ} 20' S$ . —  $56^{\circ} 38' W$ . Sand and gravel. 150—  
200 m. 16. 1. 1902.  
Number of species found: 1; Number of specimens found: 10.
- St. 6. Graham Region.  $64^{\circ} 36' S$ . —  $57^{\circ} 42' W$ . Stones and gravel. Mud-sample. 125 m. 20. 1. 1902.  
Number of species found: 29; Number of specimens found: 40.
- St. 7. Graham Region.  $65^{\circ} 56' S$ . —  $54^{\circ} 35' W$ . Stone-mixed mud. 920 m. 22. 1. 1902.  
Number of species found: 3; Number of specimens: 7.
- St. 8. Graham Region. Position of the station as well as depth uncertain.  $64^{\circ} 5' S$ . —  $56^{\circ} 37' W$ . Loose  
clay. 360 m. 11. 2. 1902.  
Number of species: 9; Number of specimens: 33.
- St. 11. Graham Region.  $65^{\circ} 19' S$ . —  $56^{\circ} 48' W$ . Gravel-mixed clay. 400 m. 18. 2. 1902.  
Number of species: 31; Number of specimens: 68.
- St. 15. Falkland Islands. Port William.  $51^{\circ} 40' S$ . —  $57^{\circ} 49' W$ . Macrocystis-Formation. 10 m. 31. 3. 1902.  
Number of species found: 44; Number of specimens: 150.
- St. 18. South Georgia. Mouth of the Westford, Cumberland Bay.  $54^{\circ} 15' S$ . —  $36^{\circ} 25' W$ . Loose clay. 250  
m. Bottom temp. + 1,2 C. 22. 4. 1902.  
Number of species: 13; Number of specimens found: 125.
- St. 21. South Georgia. Mouth of the Possession-Bay.  $54^{\circ} 8' S$ . —  $37^{\circ} 3' W$ . Clay. 200 m. 9. 5. 1902. Bottom  
temp. + 1,5 C.  
Number of species found: 15; Number of specimens found: 79.
- St. 22. South Georgia. Off the May-Bay.  $54^{\circ} 17' S$ . —  $36^{\circ} 28' W$ . Clay with some algae. 75 m. Bottom temp.  
+ 1,5 C. 14. 5. 1902.  
Number of species found: 26; Number of specimens found: 85.

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.  
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.  
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.  
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.  
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.  
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.  
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.  
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.  
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.  
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.  
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.  
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.  
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.  
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.  
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.  
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.  
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.  
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.  
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.  
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.  
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.  
Number of species found: 63; Number of specimens found 247.

- St. 49. Falkland Islands. Berkeley Sound.  $51^{\circ} 35' S$ . —  $57^{\circ} 56' W$ . Shells and stones. 25—30 m. 10. 8. 1902.  
Number of species found: 27; Number of specimens found: 58.
- St. 51. Falkland Islands. Port William.  $51^{\circ} 40' S$ . —  $57^{\circ} 42' W$ . Sand. 22 m. 3. 9. 1902.  
Number of species: 45; Number of specimens found: 245.
- St. 53. Falkland Islands. Port William.  $51^{\circ} 40' S$ . —  $57^{\circ} 47' W$ . Sand and gravel. 12 m. 3. 9. 1902.  
Number of species found: 65; Number of specimens found: 372.
- St. 54. Falkland Islands. Stanley Harbour.  $51^{\circ} 42' S$ . —  $57^{\circ} 50' W$ . Ooze with shells. 10 m. 3. 9. 1902.  
Number of species found: 2; Number of specimens found: 7.
- St. 55. Falkland Islands. Port Albemarle.  $52^{\circ} 11' S$ . —  $60^{\circ} 26' W$ . Sandy bottom with algae. 40 m. 8. 9. 1902.  
Number of species found: 33; Number of specimens found: 113.
- St. 56. Falkland Islands. Port Albemarle. Albemarle Harbour.  $52^{\circ} 9' S$ . —  $60^{\circ} 33' W$ . Sandy bottom with algae. 15 m. 8. 9. 1902.  
Number of species found: 15; Number of specimens found: 40.
- St. 57. Falkland Islands. Port Albemarle. Albemarle Harbour.  $52^{\circ} 8' S$ . —  $60^{\circ} 33' W$ . Sand. 18—30 m. 11. 9. 1902.  
Number of species found: 21; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland.  $52^{\circ} 29' S$ . —  $60^{\circ} 36' W$ . Sand and gravel. 197 m. 11. 9. 1902.  
Number of species found: 23; Number of specimens found: 93.
- St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank.  $53^{\circ} 45' S$ . —  $61^{\circ} 10' W$ .  
Crushed shells with stones 137—150 m. 12. 9. 1902.  
Number of species found: 20; Number of specimens found: 70.
- St. 62. Fuegian Archipelago. Beagle-Channel.  $54^{\circ} 53' S$ . —  $67^{\circ} 56' W$ . Sand-mixed clay. 140 m. 16. 9. 1902.  
Number of species found: 12; Number of specimens found: 63.
- St. 64. Fuegian Archipelago. North side of the Beagle Channel between Ushuaia and Lapataia.  $54^{\circ} 52' S$ . —  $68^{\circ} 25' W$ . Shells and algae. 35 m. 13. 10. 1902.  
Number of species found: 33; Number of specimens found: 192.
- St. 67. Fuegian Archipelago. Ushuaia.  $54^{\circ} 49' S$ . —  $68^{\circ} 18' W$ . Ooze. 6 m. 16. 10. 1902.  
Number of species found: 9; Number of specimens found: 42.