

***Desmodora microchaeta* ALLGÉN**

ALLGÉN 1929, p. 456, fig. 17 a—c

Localities and material. — Fuegian Archipelago, St. 64: 1♀; Falkland Islands, St. 15: 2♀♀; St. 40: 1♀; St. 41: 1♀; St. 47: 1♂, 1 juv.; South Georgia, St. 28: 1♀, 1 juv.; St. 34: 1♀.

Geographical distribution. — Norway: Herdla (ALLGÉN 1932), Bud (ALLGÉN 1939), Tarva (ALLGÉN 1934), Fröya (ALLGÉN 1946), Terningen, Ingöy Sea, and Knivskjærodden (ALLGÉN 1940), Vega, Rodöy, Gåsnes and Stappen (ALLGÉN 1940, 1943), Sweden: West Coast, vicinity of the Zoological Station Kristineberg (ALLGÉN 1929), Mediterranean: Western Part, coast of France, Banyuls-sur-Mer (ALLGÉN 1942), Hawaii: Honolulu (ALLGÉN 1951).

List of Localities

- St. 1. Off the Coast of Uruguay. Black-grey clay. $33^{\circ} 0' S$. — $51^{\circ} 10' W$. 80 m. 12. 12. 1901.
Number of species found: 8; Number of specimens found: 21.
- St. 2. Off the Coast of North Argentina. $37^{\circ} 15' S$. — $56^{\circ} 8' W$. Sand-mixed gravel. 100 m. 23. 12. 1901.
Number of species found: 12; Number of specimens found: 26.
- St. 3. Fuegian Archipelago. $54^{\circ} 43' S$. — $64^{\circ} 8' W$. Rubble stones and gravel. 36 m. 6. 1. 1902.
Number of species found: 20; Number of specimens found: 37.
- St. 5. Graham Region. S. East of the Seymour Sound. $64^{\circ} 20' S$. — $56^{\circ} 38' W$. Sand and gravel. 150—
200 m. 16. 1. 1902.
Number of species found: 1; Number of specimens found: 10.
- St. 6. Graham Region. $64^{\circ} 36' S$. — $57^{\circ} 42' W$. Stones and gravel. Mud-sample. 125 m. 20. 1. 1902.
Number of species found: 29; Number of specimens found: 40.
- St. 7. Graham Region. $65^{\circ} 56' S$. — $54^{\circ} 35' W$. Stone-mixed mud. 920 m. 22. 1. 1902.
Number of species found: 3; Number of specimens: 7.
- St. 8. Graham Region. Position of the station as well as depth uncertain. $64^{\circ} 5' S$. — $56^{\circ} 37' W$. Loose
clay. 360 m. 11. 2. 1902.
Number of species: 9; Number of specimens: 33.
- St. 11. Graham Region. $65^{\circ} 19' S$. — $56^{\circ} 48' W$. Gravel-mixed clay. 400 m. 18. 2. 1902.
Number of species: 31; Number of specimens: 68.
- St. 15. Falkland Islands. Port William. $51^{\circ} 40' S$. — $57^{\circ} 49' W$. Macrocystis-Formation. 10 m. 31. 3. 1902.
Number of species found: 44; Number of specimens: 150.
- St. 18. South Georgia. Mouth of the Westford, Cumberland Bay. $54^{\circ} 15' S$. — $36^{\circ} 25' W$. Loose clay. 250
m. Bottom temp. + 1,2 C. 22. 4. 1902.
Number of species: 13; Number of specimens found: 125.
- St. 21. South Georgia. Mouth of the Possession-Bay. $54^{\circ} 8' S$. — $37^{\circ} 3' W$. Clay. 200 m. 9. 5. 1902. Bottom
temp. + 1,5 C.
Number of species found: 15; Number of specimens found: 79.
- St. 22. South Georgia. Off the May-Bay. $54^{\circ} 17' S$. — $36^{\circ} 28' W$. Clay with some algae. 75 m. Bottom temp.
+ 1,5 C. 14. 5. 1902.
Number of species found: 26; Number of specimens found: 85.

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.
Number of species found: 63; Number of specimens found 247.

- St. 49. Falkland Islands. Berkeley Sound. $51^{\circ} 35' S$. — $57^{\circ} 56' W$. Shells and stones. 25—30 m. 10. 8. 1902.
Number of species found: 27; Number of specimens found: 58.
- St. 51. Falkland Islands. Port William. $51^{\circ} 40' S$. — $57^{\circ} 42' W$. Sand. 22 m. 3. 9. 1902.
Number of species: 45; Number of specimens found: 245.
- St. 53. Falkland Islands. Port William. $51^{\circ} 40' S$. — $57^{\circ} 47' W$. Sand and gravel. 12 m. 3. 9. 1902.
Number of species found: 65; Number of specimens found: 372.
- St. 54. Falkland Islands. Stanley Harbour. $51^{\circ} 42' S$. — $57^{\circ} 50' W$. Ooze with shells. 10 m. 3. 9. 1902.
Number of species found: 2; Number of specimens found: 7.
- St. 55. Falkland Islands. Port Albemarle. $52^{\circ} 11' S$. — $60^{\circ} 26' W$. Sandy bottom with algae. 40 m. 8. 9. 1902.
Number of species found: 33; Number of specimens found: 113.
- St. 56. Falkland Islands. Port Albemarle. Albemarle Harbour. $52^{\circ} 9' S$. — $60^{\circ} 33' W$. Sandy bottom with algae. 15 m. 8. 9. 1902.
Number of species found: 15; Number of specimens found: 40.
- St. 57. Falkland Islands. Port Albemarle. Albemarle Harbour. $52^{\circ} 8' S$. — $60^{\circ} 33' W$. Sand. 18—30 m. 11. 9. 1902.
Number of species found: 21; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland. $52^{\circ} 29' S$. — $60^{\circ} 36' W$. Sand and gravel. 197 m. 11. 9. 1902.
Number of species found: 23; Number of specimens found: 93.
- St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank. $53^{\circ} 45' S$. — $61^{\circ} 10' W$.
Crushed shells with stones 137—150 m. 12. 9. 1902.
Number of species found: 20; Number of specimens found: 70.
- St. 62. Fuegian Archipelago. Beagle-Channel. $54^{\circ} 53' S$. — $67^{\circ} 56' W$. Sand-mixed clay. 140 m. 16. 9. 1902.
Number of species found: 12; Number of specimens found: 63.
- St. 64. Fuegian Archipelago. North side of the Beagle Channel between Ushuaia and Lapataia. $54^{\circ} 52' S$. — $68^{\circ} 25' W$. Shells and algae. 35 m. 13. 10. 1902.
Number of species found: 33; Number of specimens found: 192.
- St. 67. Fuegian Archipelago. Ushuaia. $54^{\circ} 49' S$. — $68^{\circ} 18' W$. Ooze. 6 m. 16. 10. 1902.
Number of species found: 9; Number of specimens found: 42.