

Eurystomatina tenuicaudatum ALLGÉN

ALLGÉN 1932, p. 120—122, fig. 15 a—c.

Localities and material. — Coast of Uruguay, St. 1: 1♀; Fuegian Archipelago, St. 64: 1♀; Falkland Islands, St. 15: 1♂, 1 juv.; St. 41: 1♂, 6♀♀, 1 juv.; St. 42: 6♂♂, 1♀; St. 42a: 1♂; St. 47: 1♂; St. 53: 1♂; South Georgia, St. 21: 1♂; St. 23: 2♂♂, 2♀♀, 1 juv.; St. 23b: 1♂, 1♀; St. 25: 3 ♂♂, 1 juv.; St. 26: 1♂; St. 28: 1♀; St. 30: 13 ♂♂, 12 ♀♀, 4 juv.; St. 33: 1♂, 1♀; St. 34: 2♂♂, 1♀, 1 juv.; Graham Land, St. 6: 1♂.

From the above localities, numerous specimens of an *Enchelidiid* have been gathered, which in their external shape seem to be very similar and closely related to *Eurystomatina tenuicaudatum*, described by the present author from the Campbell Island. They seem to differ only in their tail being stronger thinned in its posterior half.

Body about evenly thick and tapering only in the oesophageal region.

Head rounded with strongly elongated slender bristles, longer than the head diameter at their level.

Buccal cavity consisting of 2 parts, separated from each other by a ring of minute bodies; the anterior part shorter and broader than the posterior one and bordered by walls, strongly bent outwards. Posterior part deeper and bordered by straight walls, only posteriorly bent inwards.

Tooth sharply pointed and more elongated than in the type specimen. Its point situated at the limit between both parts.

Oesophagus elongated, posteriorly gradually thickened.

Tail, as already mentioned, in its posterior half stronger thinned than in the Campbell specimens, tapering in its anterior half gradually, in its posterior half about uniformly thinned, the thickness of this part making only 1/8 of the anal body diameter. In front of the anus there are 3 small, pyriform, caudal glands, the anterior ventral, the 2 other more dorsal.

Spicules and supplementary organs in shape and structure quite typical.

Vulva and the female organs too, of a shape characteristic of this species.

Geographical distribution. — Campbell Island (ALLGÉN 1932).

List of Localities

- St. 1. Off the Coast of Uruguay. Black-grey clay. $33^{\circ} 0' S$. — $51^{\circ} 10' W$. 80 m. 12. 12. 1901.
Number of species found: 8; Number of specimens found: 21.
- St. 2. Off the Coast of North Argentina. $37^{\circ} 15' S$. — $56^{\circ} 8' W$. Sand-mixed gravel. 100 m. 23. 12. 1901.
Number of species found: 12; Number of specimens found: 26.
- St. 3. Fuegian Archipelago. $54^{\circ} 43' S$. — $64^{\circ} 8' W$. Rubble stones and gravel. 36 m. 6. 1. 1902.
Number of species found: 20; Number of specimens found: 37.
- St. 5. Graham Region. S. East of the Seymour Sound. $64^{\circ} 20' S$. — $56^{\circ} 38' W$. Sand and gravel. 150—
200 m. 16. 1. 1902.
Number of species found: 1; Number of specimens found: 10.
- St. 6. Graham Region. $64^{\circ} 36' S$. — $57^{\circ} 42' W$. Stones and gravel. Mud-sample. 125 m. 20. 1. 1902.
Number of species found: 29; Number of specimens found: 40.
- St. 7. Graham Region. $65^{\circ} 56' S$. — $54^{\circ} 35' W$. Stone-mixed mud. 920 m. 22. 1. 1902.
Number of species found: 3; Number of specimens: 7.
- St. 8. Graham Region. Position of the station as well as depth uncertain. $64^{\circ} 5' S$. — $56^{\circ} 37' W$. Loose
clay. 360 m. 11. 2. 1902.
Number of species: 9; Number of specimens: 33.
- St. 11. Graham Region. $65^{\circ} 19' S$. — $56^{\circ} 48' W$. Gravel-mixed clay. 400 m. 18. 2. 1902.
Number of species: 31; Number of specimens: 68.
- St. 15. Falkland Islands. Port William. $51^{\circ} 40' S$. — $57^{\circ} 49' W$. Macrocystis-Formation. 10 m. 31. 3. 1902.
Number of species found: 44; Number of specimens: 150.
- St. 18. South Georgia. Mouth of the Westford, Cumberland Bay. $54^{\circ} 15' S$. — $36^{\circ} 25' W$. Loose clay. 250
m. Bottom temp. + 1,2 C. 22. 4. 1902.
Number of species: 13; Number of specimens found: 125.
- St. 21. South Georgia. Mouth of the Possession-Bay. $54^{\circ} 8' S$. — $37^{\circ} 3' W$. Clay. 200 m. 9. 5. 1902. Bottom
temp. + 1,5 C.
Number of species found: 15; Number of specimens found: 79.
- St. 22. South Georgia. Off the May-Bay. $54^{\circ} 17' S$. — $36^{\circ} 28' W$. Clay with some algae. 75 m. Bottom temp.
+ 1,5 C. 14. 5. 1902.
Number of species found: 26; Number of specimens found: 85.

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.
Number of species found: 63; Number of specimens found 247.

- St. 49. Falkland Islands. Berkeley Sound. 51° 35' S. — 57° 56' W. Shells and stones. 25—30 m. 10. 8. 1902.
Number of species found: 27; Number of specimens found: 58.
- St. 51. Falkland Islands. Port William. 51° 40' S. — 57° 42' W. Sand. 22 m. 3. 9. 1902.
Number of species: 45; Number of specimens found: 245.
- St. 53. Falkland Islands. Port William. 51° 40' S. — 57° 47' W. Sand and gravel. 12 m. 3. 9. 1902.
Number of species found: 65; Number of specimens found: 372.
- St. 54. Falkland Islands. Stanley Harbour. 51° 42' S. — 57° 50' W. Ooze with shells. 10 m. 3. 9. 1902.
Number of species found: 2; Number of specimens found: 7.
- St. 55. Falkland Islands. Port Albemarle. 52° 11' S. — 60° 26' W. Sandy bottom with algae. 40 m. 8. 9. 1902.
Number of species found: 33; Number of specimens found: 113.
- St. 56. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 9' S. — 60° 33' W. Sandy bottom with algae. 15 m. 8. 9. 1902.
Number of species found: 15; Number of specimens found: 40.
- St. 57. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 8' S. — 60 ° 33' W. Sand. 18—30 m. 11. 9. 1902.
Number of species found: 21; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland. 52° 29' S. — 60° 36' W. Sand and gravel. 197 m. 11. 9. 1902.
Number of species found: 23; Number of specimens found: 93.
- St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank. 53° 45' S. — 61° 10' W.
Crushed shells with stones 137—150 m. 12. 9. 1902.
Number of species found: 20; Number of specimens found: 70.
- St. 62. Fuegian Archipelago. Beagle-Channel. 54° 53' S. — 67° 56' W. Sand-mixed clay. 140 m. 16. 9. 1902.
Number of species found: 12; Number of specimens found: 63.
- St. 64. Fuegian Archipelago. North side of the Beagle Channel between Ushuaia and Lapataia. 54° 52' S. — 68° 25' W. Shells and algae. 35 m. 13. 10. 1902.
Number of species found: 33; Number of specimens found: 192.
- St. 67. Fuegian Archipelago. Ushuaia. 54° 49' S. — 68° 18' W. Ooze. 6 m. 16. 10. 1902.
Number of species found: 9; Number of specimens found: 42.