

***Neochromadora poecilosoma* (DE MAN)**

Chromadora poecilosoma DE MAN 1893, p. 98, pl. VI, fig. 7 a—g; *Neochromadora poecilosoma* (DE MAN) MICOLETZKY 1924, p. 157.

Localities and material. — Falkland Islands, St. 41: 2♀♀; St. 42a: 1♀; St. 46: 1♀; South Georgia, St. 28: 1♂, 4♀♀

The few specimens recorded from the above localities are in every respect typical representatives of this old North Sea species.

Geographical distribution. — *Norway*: Oslofjord (ALLGÉN 1931), Herdla (ALLGÉN 1932), Bud (ALLGÉN 1939), Trondheimsfjord (ALLGÉN 1933), Tarva (ALLGÉN 1934) Rörvik (ALLGÉN 1946), Vega, Rodøy, Gåsnes (ALLGÉN 1940, 1943), *Sweden*: West Coast, vicinity of the Zool. Stat. Kristineberg (ALLGÉN 1929), South Kattegatt (ALLGÉN 1934), the Sound (ALLGÉN 1935), *Denmark*: Little Belt (DITLEVSEN 1919), *North Sea and Channel* (DE MAN 1893), *Holland*: Coast of Walcheren (DE MAN 1893), Zeeland (DE MAN 1906, 1907), *Zuider Sea* (DE MAN 1922), *Belgium* (SCHUURM. STEKH. and DE CONINCK 1933).

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.
Number of species found: 63; Number of specimens found 247.