

Paralinhomoeus pellucidus n. sp.

Fig. 228 a, b

Locality and material. — South Georgia, St. 28: 1♀

Dimensions: L = 2,815 mm., $\alpha = 59,89$, $\beta = 18,77$, $\gamma = 22,99$, V = 51,50 %

Body slender, elongated, in its whole length of about even thickness. Cuticle smooth. Head rounded, light-yellow-gray coloured, with small, slender, submedian bristles. Buccal cavity typical. Oesophagus short, slender, and increasing posteriorly to a bulb, only very weakly demarcated from the remaining oesophagus, in a short distance (14μ) before its middle surrounded by the nerve ring. Lateral organs situated far anteriorly, only 13,6—17 μ behind the front end, circular, in section 1/3 of the corresponding head diameter. Tail short, slender, conical, tapering gradually, posteriorly evidently swollen, its length $4,4 \times$ the anal body diameter.

Vulva about in the middle of the body. Female organs paired-symmetrical, with straight, not reflexed ovaries.

St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. $54^{\circ} 22' S$. — $36^{\circ} 28' W$. Sand and algae. 12—15 m. 24. 5. 1902.

Number of species found: 58; Number of specimens found: 338.

