

4. *C. PUNCTATUS*, n. sp. (Plate XIII. figs. 217, 218.)



Male, length $\frac{1}{12}$ " , breadth $\frac{1}{263}$ " .

External Characters.—Body powerful, tapering very slightly anteriorly, but gradually to a point posteriorly. Head obtusely rounded, provided with a circlet of 6–8 small spreading setæ; other small setæ scattered over posterior part of body. Integument having rows of dots arranged in transverse series $\frac{1}{15000}$ " apart; two lateral convex prominences of integument in cervical region nearly opposite base of pharyngeal cavity, $\frac{1}{3333}$ " in diameter.

Pharyngeal cavity cup-shaped. *Œsophagus* about $\frac{1}{10}$ th of total length. *Intestine* sparsely and irregularly covered with large brownish-coloured fat-particles. *Anus* $\frac{1}{117}$ " from posterior extremity. *Spicules* $\frac{1}{333}$ " long. *Ocelli* two, greenish brown, large, $\frac{1}{1428}$ " from anterior extremity. *Subcutaneous gland-cells* wanting.

Female, not seen.

Hab. Marine surface-mud from estuary, Falmouth.



217. *Cyatholaimus punctatus*, n. sp.; anterior extremity of male.

218. Posterior extremity of male.

p. Integument.

p'. Cervical markings of same.