

**Viscosia antarctica** n. sp.

Fig. 59 a, b

Localities and material. — South Georgia, St. 22b: 1♂; St. 23: 2♂♂; St. 23b: 1♂; St. 24: 1♂; St. 26: 2♂♂, 1♀; St. 30: 1♂; St. 33: 1♂; St. 34: 1♂; Falklands, St. 58: 1♂

Dimensions: ♂ L = 5,0 mm.,  $a = 65,88$ ,  $\beta = 8,0$ ,  $\gamma = 43,75$   
♂ L = 6,069 mm.,  $a = 52,77$ ,  $\beta = 7,88$ ,  $\gamma = 41,86$   
♀ L = 4,850 mm.,  $a = 52,72$ ,  $\beta = 6,93$ ,  $\gamma = 31,7$

This species, recorded as males mostly from the above localities in South Georgia, is also in its external shape similar to the old *V. langrunensis*, but differs in the armature of its buccal cavity as well as in the shape of its tail and its spicules.

Head clearly rounded, with rather slender, submedian bristles.

Buccal cavity about twice deeper than broad. *Left subventral tooth larger than the right tooth.* Dorsal tooth immediately in front of the middle of the buccal cavity. All three teeth sharply pointed. Oesophagus in relation to the body length rather short, slender, strongly muscular, posteriorly thickened.

Tail tapering gradually in the anterior third and in the 2 posterior thirds rather strongly and uniformly thinned and at the end provided with 2 slender hairs. Anterior part of tail 50  $\mu$  in length, the posterior part 98  $\mu$  in length. Thickness of the lastnamed part 1/6 of the anal body diameter. Spicules absolutely straight, at the proximal end button-shaped, distally sharply pointed, in length 65,6  $\mu$ .




Fig. 59. *Viscosia antarctica* n. sp. a. Anterior end,  $\times 450$ , b. Posterior part of body,  $\times 770$

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.  
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.  
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.  
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.  
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.  
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.  
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.  
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.  
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.  
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.  
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.  
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.  
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.  
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.  
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.  
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.  
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.  
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.  
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.  
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.  
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.  
Number of species found: 63; Number of specimens found 247.

- St. 49. Falkland Islands. Berkeley Sound. 51° 35' S. — 57° 56' W. Shells and stones. 25—30 m. 10. 8. 1902.  
Number of species found: 27; Number of specimens found: 58.
- St. 51. Falkland Islands. Port William. 51° 40' S. — 57° 42' W. Sand. 22 m. 3. 9. 1902.  
Number of species: 45; Number of specimens found: 245.
- St. 53. Falkland Islands. Port William. 51° 40' S. — 57° 47' W. Sand and gravel. 12 m. 3. 9. 1902.  
Number of species found: 65; Number of specimens found: 372.
- St. 54. Falkland Islands. Stanley Harbour. 51° 42' S. — 57° 50' W. Ooze with shells. 10 m. 3. 9. 1902.  
Number of species found: 2; Number of specimens found: 7.
- St. 55. Falkland Islands. Port Albemarle. 52° 11' S. — 60° 26' W. Sandy bottom with algae. 40 m. 8. 9. 1902.  
Number of species found: 33; Number of specimens found: 113.
- St. 56. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 9' S. — 60° 33' W. Sandy bottom with algae. 15 m. 8. 9. 1902.  
Number of species found: 15; Number of specimens found: 40.
- St. 57. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 8' S. — 60° 33' W. Sand. 18—30 m. 11. 9. 1902.  
Number of species found: 21; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland. 52° 29' S. — 60° 36' W. Sand and gravel. 197 m. 11. 9. 1902.  
Number of species found: 23; Number of specimens found: 93.
- St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank. 53° 45' S. — 61° 10' W.  
Crushed shells with stones 137—150 m. 12. 9. 1902.  
Number of species found: 20; Number of specimens found: 70.
- St. 62. Fuegian Archipelago. Beagle-Channel. 54° 53' S. — 67° 56' W. Sand-mixed clay. 140 m. 16. 9. 1902.  
Number of species found: 12; Number of specimens found: 63.
- St. 64. Fuegian Archipelago. North side of the Beagle Channel between Ushuaia and Lapataia. 54° 52' S. — 68° 25' W. Shells and algae. 35 m. 13. 10. 1902.  
Number of species found: 33; Number of specimens found: 192.
- St. 67. Fuegian Archipelago. Ushuaia. 54° 49' S. — 68° 18' W. Ooze. 6 m. 16. 10. 1902.  
Number of species found: 9; Number of specimens found: 42.