

At present genus *Viscosia* consists of the following seven species:

- 1) *V. viscosa* (Bastian 1865)—*Oncholaimus* v. Bastian 1865: Tr. Linn. Soc. 25, p. 136, Table 11, Figures 131—133; Bütschli 1874: Abh. Senck. Ges. 9, p. 39, Table 9, Figure 38; *Oncholaimus* (*Viscosia*) v. de Man 1890: Mém. Soc. Zool. France 3, p. 184, Table 4, Figure 7. Holland, France (de Man), Falmouth, mud from the harbor (Bastian), Kiel Firth, oyster bank at Sylt—Norway (Bütschli).
- 2) *V. minor* n.sp., see p. 88.
- 3) *V. cobbi* n.sp., see p. 86.
- 4) *V. langrunensis* (de Man 1890); *Oncholaimus* (V.) l. de Man: Mém. Soc. Zool. 3, p. 186, Table 4, Figure 8.
- 5) *V. glabra* (Bastian 1865), see p. 127.
- 6) *V. linstowi* (de Man 1904); *Oncholaimus* (V.) l. Rés. voy. Belgica. Nématodes libres p. 15, Table 3, Figure 5. On shell of crab: Orange Bay, Tierra del Fuego.
- 7) *V. pellucida* (Cobb 1898); *Oncholaimus* p. Cobb: Proc. Linn. Soc. NSW 23, p. 39. Sand near tidemark. Port Jackson NSW.