

(11)

Lernaeopodoida.

Lernaeopodidae.

Brachiella, valid.

Cuvier, 1830.

B. thynnica.<sup>Ⓟ</sup>

Syn. Brachiella Cuvier, 1830, vol. 3, p. 257,

Brachiella M-Edwards, 1840, p. 511,

Brachiella Wilson, 1915, p. 698,

Brachiella T. & A. Scott, 1913, p. 203.

Diminutive of βραχύς = small.

Cephalothorax markedly thin & elongated,  
worm-like. First mxpds. placed close  
behind the mouth; second mostly long and  
arm-like, generally divided to a very tip  
where is found a organ of attachment.

Genital segment oval or quadrilateral -  
abdomen simple.

Male a pigmy, attached to body, neck, or mxpds.  
of ♀. Cephalothorax distinctly divided from the  
long segmented abdomen by a marked con-  
striction: both pairs of mxpds large, uncinata.

Brachiella

(2)

proposal to suppress Parabrachiella & transfer all  
of its species to Brachiella; Kabata, 1970,  
p. 2165.

- , ACLOQUE, 1899: 204-205, key to genera.
- CRESSEY ET AL. 1983: 232 -
- KABATA 1992: 149-150, DIAGNOSIS

Brachiella albida (Rangnetar)

- SEE NEOBRACHIELLA

— —, Pillai, 1967, p. 1666, fig. 260



Brachiella annulata Markevitch, 1940 - see Neobrachiella

Brachiella anserina Wilson, 1908 - see Neobrachiella

Lernaeopodidae. Lernaeopodidae

Brachiella antarctica Quidor - ♀♂

— —. Quidor, 1906, p. <sup>1, 14-17,</sup> 30: pl. 3, figs. 45-48,

ANTARCTIC - N.SP.

Referred to genus Eubrachiella.

(Wilson, 1915, p. 699).

Host. Gills of Dissostichus eleginoides.

Antarctic Ocean.

Brachiella appendiculata, Stp. & Totk. ♂ & ♀.

— — —, Stp. & Totk., 1861, p. 419: pl. XV, fig. 35.

B. appendiculosa Steenstrup & Lutken, 1861;

Rangnekar, 1961, p. 203; B. appendiculata p. 204 transfer to Thysanote.

= B. appendiculosa B-Smith  
A synonym for Thysanote a — — — (g.v.)  
(Wilson, 1915, p. 700).

Brachiella appendiculata (Krøyer, 1863) - see Neobrachiella

Lernaeopodidae

Brachiella appendiculosa, B-Smith, ♂, ♀,

— —, B-Smith, 1898.a, p. 14; pl. VI, figs. 1-3.

--, Rangnekar, 1961, p. 203-4, transfer to Thysanote;  
= B. appendiculata Stp. & Ltk.

A synonym for Thysanote appendiculata (g. v.)  
(Wilson, 1915, p. 700).

Brachiella appendiculosa (Krøyer, 1863) - see Neobrachiella

Brachiella australis (Wilson, 1923)

<sup>from</sup>  
~~see~~ Parabrachiella australis, Kabata, 1970, p.  
2165.



Brachiella bera Yamaguti, 1939 - see Neobrachiella

Brachiella bispinosa Nordmann, 1832 - see Neobrachiella

Brachiella chavesii Van Beneden - see Neobrachiella

Brachiella cheveuxii Van Beneden - see Neobrachiella

Brachiella cirrata Heegaard - SEE NEOBRACHIELLA

-- --, Heegaard 1962. pp. 191-192, fig. 237-245

--, Kabata, 1968, p. 505

NEOBRACHIELLA -, CASTRO + BAEZA 1987:22

Brachiella cirrocauda Heegaard - SEE NEOBRACHIELLA

-- --, Heegaard 1962. pp. 189-190, fig. 233-236  
-- , Kabata, 1968, p. 505

NEOBRACHIELLA -, CASTRO + BAEZA 1987: 22

Brachiella concura Wilson ♀

— —, Wilson, 1913, p. 262: pl. 51, f. 284-288.

— —, Wilson, 1915, p. 703 (key).

— —, Wilson, 1932, p. 521: fig. 309.

--, Yamaguti & Yamasu, 1959, p. 137

Host. Gills of Dasyatis lastata.

--, LEWIS, 1967: 186

West Indies.

Lernaeopodoida Lernaeopodidae

Brachiella coryphaenae Pearse

--- Pearse, A.S. 1952, pp. 35-36, figs. 129-135

A synonym of Charopinus quaternius C.B. Wilson T&B.

-- --, Pillai 1962. pp. 85-86, fig. 18



Brachiella cybii Pillai, et al.

— Pillai, Prabha, & Balaraman, 1982: 11-13.  
figs. 1c-g.

Brachiella dentata, n. sp.

— —. Wilson, 1912. p. 97: pl. IX.

Changed to genus Charopinus. g. v.  
(Wilson, 1915, p. 700).

Host. Gills of Raja binoculata.

Nanaimo, B.C.

Brachiella dentici Nunes Ruivo

-- -- Nunes Ruivo, 1962. pp. 83-85, fig. 7-8 —

SEE NEOBRACHIELLA

Brachiella elegans, Richiardi, ♀, ♂.

- — Richiardi, 1880, p. 151, (no description).  
 — — Carus, 1885, p. 377.  
 — — Brian, 1899.b, p. 8, fig. 4 (text).  
 — — Brian, 1899.c, p. 204.

Host. Gills of Lichia glauca and L. amia.

Mediterranean.

— —, Miculicich, 1905, a, p. 603.

— —, Wilson, 1915, p. 707; pl. 54, figs. 226-233;  
pl. 55, fig. 234.

— —, Wilson, 1932, p. 522, :fig. 311

--, Yamaguti & Yamasu, 1960, p. 139

Brachiella elegans

card 2

--, Ho, 1977, p. 11, fig. 16

-- Castro Romero & Baeza Kuroki 1983: 3, 8-12, Figs.  
14-27, ♀♂; host Seriola mazatlana, Chile; as  
Neobrachiella elegans.

Branchiella exigua Brian, 1906 - see Neobrachiella

Brachiella exilis (Shiino, 1956) - see Neobrachiella



Brachiella fasciculata (Teidy), ♀ <sup>Leptaeopodidae.</sup>

Anchorella, Teidy, 1889, p. 138, 1 text figure.

Host Throat of Red Snapper, Neomacris aya

Florida.

Brachiella fimbriata, Heller, ♀.

— — Heller, 1865, p. 240: pl. XXIV, fig. 2.

--, Rangnekar, 1961, p. 203<sup>204</sup> transfer to Thysanote.

A synonym for Thysanote fimbriata (g.v.),  
(Wilson, 1915, p. 700).

Brachiella gaini Quidor ♀.

— — Quidor, 1912, p. 211: pl. 2, fig. 24: pl. 3,  
fig. 27.

Transferred to new genus Eubrachiella 1914

Host. Gills + gill cavity of Trematodus sp.

Port-Lockroy -  
Antarctic Ocean.

Brachiella genypteri Capart, 1959 - see Neobrachiella

- SEE NEOBRACHIELLA

Brachiella gracilis Wilson, 1908

~~card 2~~

Neobrachiella gracilis, Castro Romero & Baeza Kuroki,  
1983: 11.

Brachiella gulosa Wilson, 1915 - see Neobrachiella

Brachiella hoplognathi (Yamaguti, 1939 ) see  
Neobrachiella

Brachiella hostilis (Heller, 1865) - see Neobrachiella



Brachiella impudica Nordmann 1832 - see Neobrachiella

Brachiella impudica parva Bassett-Smith, 1896

---, Kabata, 1973, p. 7, see B. parva

Lernaeopodidae. Lernaeopodoidea.

Brachiella inconcinna Richiardi, ♀.

— — Richiardi, 1880, p. 151, nomen nudum.

— — Carus, 1885, p. 377.

— — Miculicich, 1905, a, p. 603.

— — Wilson, 1915, p. 700.

Host. Raja maculata.

Mediterranean.

Brachiella incurva (Shiino, 1956) - see Neobrachiella

Brachiella indica Pillai, 1968

n. sp. Pillai, 1968, pp. 123-129, figs. 3-6

-- Kabata & Tareen, 1987: 137, 141, 142 a syn of  
Neobrachiella pillaii nom. nov. Kuwait

Brachiella insidiosus, Heller, ♂, ♀.

—, Heller, 1865, p. 239; pl. XXIV, fig. 1.

—, —, 1866, p. 756.

—, Carus, 1885, p. 376.

—, B-Smith, 1896, p. 14, pl. VI, fig. 2, ♂.

Host, Godus spp. : Merluccius vulgaris

M. esculentus.

Mediterranean. [Wilson, 1915, p. 70]

Made type of new genus Parabrachiella.

- —, Brian, 1898, p. 25, pl. IV, fig. 27.  
— —, ———, 1899, z, p. 7.  
— —, B-Smith, 1899, p. 502.  
— —, T. Scott, 1900, p. 175, pl. VIII, figs. 40, 41.  
— —, A. Scott, 1901, a, p. 352.  
— —. Richiardi, 1880, p. 7.  
— —. Valle, 1880, p. 77. Stossich, 1880, p. 267.  
— —, ———, 1882, p. 247.  
— —, A. Scott, 1904, b, p. 124; 1904, a, p. 44.  
— —, Miculicich, 1905<sup>a</sup>, p. 603.  
— —, T. Scott, 1906, b, p. 373.

Brachiella insidiosa Heller - SEE NEOBRACHIELLA

--, Brian, 1939, pp.4,6



Brachiella intermedia Bere, 1936 - see Neobrachiella

BRACHIELLA LAGENIFORMIS - SEE

NEOBRACHIELLA INSIDIOSA LAGENIFORMIS

Brachiella lithognathae Kensley & Grindley 1973

-- Kensley & Grindley 1973: 110-112, fig. 29a-g, n. sp.

♀♂, from Lithognathus lithognathus and L. aureti,

Cape, S. Africa & Rocky Point, S.W.A.; p. 127.

--Kabata & Tareen, 1987: 137, 141 renamed Sparidicola -

Lernaeopodidae.

Lernaeopodoidea.

Brachiella lobiventris Heller. ♀,

— — Heller, 1865, p. 241; pl. 24, f. 3,

--, Rangnekar, 1961, p. 203-4, transfer to Thysanote.

A synonym of Thysanote lobiventris,  
(Wilson, 1915, p. 700).

Lernaeopodoida.

Brachiella lophii Milne Edwards, ♀

- — Milne Edwards, 1840, vol. 3, p. 514: pl. 41, g. 4,  
— — Vogt, 1877, p. 426: Carus, 1885, p. 376.  
— — B-Smith, 1899, p. 501: Wilson, 1915, p. 700  
— —, ACLOQUE, 1899: 205.

Host. Lophius piscatorius.

Naples.

Brochiella macrura, New Species, ♀.

— Wilson, 1920, p. 7: pl. 3, f. 23-28.

— Capart, 1959: 117-118, fig. 35a-d

Host Gills of African snapper, Neomacris  
fulgens Cuv. & Val.  
Banana, Mouth of Congo River -  
Africa.

Brachiella magna Kabata, 1968

n. sp., Kabata, 1968, p. 505, 508-11, fig. 2 A-J

--, Kabata, 1973, p. 10

--, CRESSEY & CRESSEY 1980: 45; Figs. 95b, 108.

Brachiella malabarica Pillai et al. 1982 - see

Neobrachiella



Brachiella malleus, Rudolphi, ♂, ♀,

Dirhynchus m-, Rudolphi, 1817, (manuscript)

Brachiella m-, Nordmann, 1832, <sup>II,</sup> p. 95.

— —, Kröyer, 1837, p. 194.

— —, Vogt, 1877, b, p. 46, pl. III, figs. 1-8 ♂,  
pl. IV, fig. 1-♀. (over)

Host. Mouth of Torpedo marmorata.

Mediterranean.

—, Carus. 1885, p. 376.

—, Brian, 1899, C, p. 204.

—, B. Smilk, 1899, p. 502.

—, Valle, 1880, p. 77.

—, Miculicich, 1905<sup>a</sup>, p. 600.

—, Stossich, 1880, p. 267.

Changed to genus Charopinus (1912)  
(Wilson, 1915, p. 700).

Brachiella merlucci Bassett-Smith, 1896) - see  
Neobrachiella

Lernaeopodoida  
Brachiella minuta Richiardi ♀.

- — Richiardi, 1880, p. 151 nomen nudum,
- — Carus, 1885, p. 377; Miculicich, 1905, a. p. 603,
- — Wilson, 1915, p. 700

Host, Pagellus erythrinus.

Mediterranean.

Brachiella mitrata C. B. Wilson - see Neobrachiella

Brachiella multifimbriata B-Smith, ♂.♀.

— — B-Smith, 1898, b, p. 96: pl. 6, fig. 2, a-g.

— — B-Smith, 1899, p. 503.

— — Wilson, 1915, pp. 652, 700

(made type of new genus Thysanotella)

—, Rangnekar, 1961, p. 203 †transfer to Thysanote.

Host, Operculum of Serranus sp.

Musc. cat.

Brachiella neglecta Richiardi ♀  
hermatopoda

- —. Richiardi, 1880, p. 151. nomen nudum  
— —, Carus, 1885, p. 377.  
— —, Brian, 1898, p. 24: pl. 4, fig 29.  
— —, Miculicich, 1905a, p. 603.

Host, Gills of Sciaeva aquila

Mediterranean.

Brachiella nitida C. B. Wilson, 1915 - see Neobrachiella



Brachiella obesa (Krøyer), ♀. ①

Lernaeopoda — Krøyer, 1837, p. 270: pl. 3, f. 13, a-c.

— — Milne Edwards, 1840, p. 516.

— — T. J. v. Beneden, 1870, p. 10: B-Smith, 1899, p. 500.

Brachiella — Olsson, 1869, p. 42.

Host, Squalus acanthias; Trigla gurnardus.

TRIGLA CUCULUS ♂.

Europe.



(2)

BRACHIELLA OBESA (KROYER)

--, ACHOUX, 1899: 205 (as LERNAEPODA OBESA)

Brachiella obesa <sup>Terrestrialopodidae</sup> Richardi, ♀

— —, Richardi, 1880, p. 157; Carus, 1885, p. 377.  
never described or figured, so cannot be absolutely  
located, but probably identical with Kröyer's  
Terrestrialopoda obesa, q. v.

Host, Gills of Trigla corax

Mediterranean.

- , Kabata, Raibaut & Ben Hassine, 1971, p. 88
- , Nunes - Ruivo, 1957, p. 91-95, fig. 1.

Brachiella oblonga, Valle, ♀.

—, Valle, 1880, p. 76. (no description or figs).

—, Miculicich, 1905<sup>a</sup>, p. 603.

—, Stossich, 1880, p. 267.

(never described or figured)

—, Nunes-Ruivo, 1957, p. 95

Host. Mugil cephalus & M. saliens  
on the pectoral fins.

Mediterranean.

---, Wilson, 1915, p. 701.

--, Kabata, Raibaut & Ben Hassine, 1971, p. 88,  
nomen nudum

Brachiella otolithi Pillai, 1962 - see Neobrachiella



Brachiella ovalis (Beneden) ♀ <sup>hermaphroditic.</sup>

Anchorella, ~~Kröyer, 1837, p. 289, pl. 3, fig. 6~~  
— —, P. J. van Beneden, 1870, p. 31: pl. 2, fig. 8.  
Brach. —, T. Scott, 1901, p. 133: pl. 7, figs. 30-35.  
— —, A. Scott, 1901a, p. 352.

Host, Gill rakers of common gurnard  
Trigla gurnardus.

—, A. Scott, 1904, p. 124; 1904, a. p. 44.  
—, T. & A. Scott, 1913, p. 210; pl. 60, fig. 7; pl. 63, figs. 26-30.

A synonym of Brachiella obesa

Scott's reference to Kröyer's Anach. ovalis is wrong - see Clavellisa ovalis. Beneden refers his species to Kröyer's A. ovalis, but he is also wrong. He should have referred it to Kröyer's Ternaeopoda obesa.

—, Wilson, 1915, p. 701.

—, Brian, 1924, p. 49.

Brachiella ovalis

card 2

--, Heegaard, 1959, p. 231

--, Boer, 1970-2, tbl. 1

Brachiella parkeri, Thomson. ♀.

— —. Thomson, 1889, p. 374: pl. 28, fig. 8, arb.

— —. B. Smith, 1899, p. 502.

Made type of new genus Thomsonella (g.v.), 1914.

Host. Raja nasuta and Trygon, sp.

New Zealand.

Brachiella parkeri Thomson, 1889

card 2

-- Kabata 1988: 56, B. p. of Bainbridge (1909) ?=

Charopinus dubius

Brachiella parva (Nunes-Ruivo)

--, Kabata, 1973, p. 7, should be suppressed, see  
B. impudica.

--, Pillai, 1968, p. 129

--, Nunes-Ruivo, 1957, p. 96-99, fig. 2.

Lernaeopodidae. Lernaeoidea.

Brachiella pastinacae P. J. van Beneden, ♀.

— — Beneden, 1857, a. p. 118: pl. 4, figs. 8-9.

A synonym of Charopinus bicaudatus,  
(Wilson, 1915, p. 656).

-- RAIBAUT + MAAMOURI 1975: 1042, 1044, SYN. OF  
PSEUDOCHAROPINUS MALLEUS

*Terusaepodidae.*  
Brachiella pernetta Blainville. ♀

"A Teruseau" - Pernetti, 1770, Vol. I. pl. I, figs. 546  
Teruseomyzon pernettaia, Blainville, 1822, p. 439 -  
Brachiella —, Cuvier, 1830, p. 257 (foot-note)  
— —, Kröyer, 1837, p. 194.

Host, A Tunny.

Europe -



--, Wilson, 1915, p. 701,

Brachiella pinguis C. B. Wilson - see Neobrachiella

Brachiella pteroplateae Yamaguti & Yamasu, 1959

n. sp. Yamaguti & Yamasu, 1959, p. 136-7, pl. XIV,  
figs. 287-293

--, Ho, 1975, p. 368

--, Lewis, 1967: 186

Brachiella ramosa, Richardi <sup>Lernaeopodidae.</sup> ♀.

— — . Richardi, 1880, a, p. 26 - no description or fig.

— — . " " , 1880, b, p. 151. " " " "

— — . Goode, 1883, p. 346. " " " "

— — . Carus, 1885, p. 376.

Host. Gills of Swordfish, Xiphias gladius

Mediterranean

- —. Miculicich, 1905, p. 603.
- —. M. J. Rathbun, 1905, p. 102.
- —. Wilson, 1916, p. 576.
- —. Wilson, 1915, p. 700

Brachiella regia Lewis — SEE NEOBRACHIELLA

--, Lewis, 1967, pp. 180,181,182,184,185,186

Brachiella richiardi Ben Hassine & Raibaut, 1978 - see

Neobrachiella

Brachiella robusta (Wilson, 1912) - see Neobrachiella



Brachiella rostrata Krøyer 1837 - see Neobrachiella

Brachiella rotunda Pearse 1952 - see Neobrachiella

Brachiella sciaenophila (Heller) ♀.

Anchorella — Heller, 1865, p. 243: pl. 24, fig. 8.

Brachiella —, Wilson, 1915, pp. 670 & 703.

Brachiella sp. (? Anchorella) Brian, 1902, p. 43: text-fig.

Clavella sciaenae Brian, 1906, p. 119: pl. 10, fig. 2: pl. 20,

fig. 13-14

Host. Gills of Sciaena sp

Mediterranean: Indian Ocean

— — Wilson, 1915, p. 670: 703,

— — B-Smith, 1899, p. 506,

Brachiella sciaenae, Brian, p. 49.

— —, Causey, 1960, p. 325. From croaker (Roncador stearnsi?), Ensenada, <sup>B.C.</sup>

Brachiella septicauda (Shiino, 1956) - see Neobrachiella

Brachiella seriolae Yamaguti & Yamasu, 1960

--, Izawa, 1969, p. 128

n. sp. Yamaguti & Yamasu, 1960, p. 138-139, pl. 9:  
figs. 7-14

Neobrachiella seriolae, Castro Romero & Baeza Kuroki,  
1983: 11-12.

Brachiella spinicephala (Ringuelet, 1945) - see

Neobrachiella

*Brachiella squali* Wilson

Wilson 1944, pp. 547-548, pl. 31, figs. 163, 164 ♀  
Host: shark (*Squalus mitsukurii*). China



Brachiella stellifera Heegaard, 1962 - see Neobrachiella

Brachiella superba Leigh-Sharpe, 1934 - see Neobrachiella

Brachiella supplicans Barnard - SEE NEOBRACHIELLA

-- Kensley & Grindley 1973: 127

Brachiella tetrici Kabata, 1968 - SEE NEOBRACHIELLA

n. sp., Kabata, 1968, pp. 505, 511-13, fig. 3 A-D

-, Kabata, 1970, p. 875

NEOBRACHIELLA -, CASTRO + BAEZA 1987: 22

Brachiella thynni, Cuvier. ♀, ♂.

- —, Cuvier, 1830, III, p. 257; pl. XV, fig. 5.  
 — —, Nordmann, 1832, p. 90.  
 — —, Kröyer, 1837, p. 194.  
 — —, M. Edwards, 1840, p. 512.

Host. Thynnus thynnus: Sciaena aquila:  
Th. vulgaris, Gills + pectoral fins.

Europe.

- , Gould, 1841, p. 341,
- , van Beneden, 1851, a, p. 128: 1861, a, p. 153.
- , Stp. & Ltk, 1861, p. 420, pl. XV, fig. 36, ♂
- , Heller, 1866, p. 756.
- , van Beneden, 1870, p. 37: pl. II, fig. 10.
- , E. van Beneden, 1870, b, p. 224 (embryology)
- , Vogt, 1877, p. 57, pl. III, fig. 9, ♂
- , Carus, 1885, p. 376.
- , Brian, 1898, p. 24: 1899, b, p. 7.
- , B-Smitk, 1899, p. 502: 1896a, p. 162.
- , Brian, 1901, pp. 1-3 - text fig. a-f.
- , Richards, 1880, p. 7.
- , Valle, 1880, p. 77
- , Brian, 1905, p. 8, Thompson & Scott, 1903, p. 294

Brachiella thynni, Cuvier.

*Thynnicola ziegleri*, Miculicich, 1904, p. 47.

*Brachiella thynni*, Stenta, 1904, p. 345.

— —, Miculicich, 1905<sup>a</sup>, p. 601.

— —, —————, 1905<sup>b</sup>, p. 733.

— —, Stossich, 1880, p. 268.

— —, M. J. Rathbun, 1905, p. 102.

*Lernaeomyzon elongata*, Blainville, 1822, p. 439

pl. unum. figs. 9, 9a.

*Brachiella thynni*, Willson, 1915, p. 703: pl. 25.  
fig. c: pl. 53, figs. 209-215.

- —, Nunes - Kuivo, 1956, p. 31. On Parathunnus obesus  
(Lowe). Angola. Near Benguela.
- —, Delamare-Deboutville + Nunes Kuivo, 1954, p. 217. Mediterranean.
- —, Kir tisinghe, 1935, p. 342, figs. 40-41. Thunnus (Germo) macropterus. Ceylon.
- —, Bere, 1936, p. 613. Pomatomus saltatrix. Gulf of Mexico.
- —, Causey, 1953, p. 15. Scomberomorus cavalla. Gulf of Mexico.
- —, Shiino, 1956, pp. 283-287, fig. 8-9. Acanthocybium solandri  
(C. + V.). Off Owase, Mie Pref., Japan.
- --, Shiino, 1960, pp. 539-540.
- --, Shiino 1963. p. 346 <sup>Shiino</sup>
- --, Pillai 1962. pp. 81-83, fig. 15-16
- , Kabata, 1973, p. 10
- , Kazachenko, 1975, p. 216



Brachiella thynni CUVIER

--, Boer, 1970-2, tbl. 1

--, Lewis, Dean, & Gilfillan, 1969, p. 427, fig. 9,  
tbls. 17, 18, 19

--, Lewis, 1967, pp. <sup>1</sup>174, 175, 176, 177, 178, <sup>186</sup>figs. 66,  
67

--, CRESSEY & CRESSEY 1980: 45; Figs. 95a, 108.

--, Pillai, 1967, p. 1665, fig. 259

--, ACLOQUE, 1899: 204-5, pl. 96

-- CRESSEY ET AL. 1983: 228-, FIG. 1j

-- JONES, J. B. 1991: 420-, S. PACIFIC, EX THYNNUS  
ALALUNGA

Böehiella thynni

--, Boer, 1970-2, tbl.1

-- ZWEARNER & LAWLER 1972: 85, OFF VIRGINIA,  
HOST ACANTHOCYBIUM SOLANDERI

-- KABATA 1992: 150-, FIG. 34C-F, SYNONYMY, HOST LIST

-- BONNET 1948: 7, HAWAII, THYNNUS SP.

Brachiella trichiuri Gnanamuthu, 1951 - see

Neobrachiella trichiuri indica

Brachiella trichiuri (Yamaguti, 1939) - see

Neobrachiella trichiuri trichiuri

Brachiella triglae Claus, 1860 - see Neobrachiella