

in

Candaciidae-Calanoida.

Calamoecia

Brady - 1906.

C. lucasi.^(M.)

Syn. Calamoecia Brady, 1906, p. 696.

Káλαμος = a reed and οἴχεω = to dwell.

CENTROPAGIDAE

CALAMOECIA BRADY, 1906

CARD 2

- SARS 1908b: 11-, DIAGNOSIS
- TOLLY, 1955: 280-281; SYN: BRUNELLA
- BAYLY 1974: 87-88, CONTRASTS TO HEMIBOECKELLA
- BAYLY 1961, 1962 - REVISED GENUS
- DASSART 1986: 64, SHORT DIAGNOSIS
- LEWIS, M. 1986: 123
- BREHM 1960b: 86-, STATUS; NOT = BRUNELLA
- BAYLY 1992c: 5-, KEY, GENERIC DIAGNOSIS

Calamoecia ampulla (Searle) 1911

(1)

Brunella ampulla Searle, 1911, pp. 177-8

Calamoecia ampulla (Searle)--Bayly 1961. pp. 78-80,
fig. 1e, 4e-f, 5d, 9a-d.

--, TOLLY, 1966: 731, 732, 734; TAB. 4

--, TOLLY, 1955: 281

--, LINDQVIST, 1961: 12

--, BAYLY 1984: 151

--, BAYLY 1985: 82, NEAR ESPERANCE, AUSTRALIA;
83, N.Z.

--, BAYLY 1979: 124-5, DISTRIBUTION; TAB. 4

--, BAYLY 1991: 2829-, DISTRIBUTION

Calamoecia ampulla (Searle, 1911)

card 2

-- Bayly 1992c: 7-, fig. 7, southern Australia

Calamoecia attenuata (Fairbridge)

Brunella attenuata Fairbridge, 1945b, pp. 67-8

Calamoecia attenuata (Fairbridge)--Bayly 1961. pp. 81-82, fig. 3a-d, 11a-b

--, TOLLY, 1955: 281

--, LINDQVIST, 1961: 12; FIG. 2

--, BAYLY 1984: 148, 149, 151

-- BAYLY 1979: 124-5, DISTRIBUTION; TAB. 4

-- MALY 1991: 2829-, DISTRIBUTION

-- BAYLY 1992c: 7, FIG. 13, WESTERN AUSTRALIA

Calamoecia australica G.O. Sars, 1908

CARD 1

- Sars 1908b:12-14, Pl. II figs. 5-19, n. sp. female + male, St. Arnaud, southern Australia
- Jolly 1955:279, 281
- Bayly 1961:63
- Lindquist 1961:12
- Bayly 1962:252-254, fig. 1
- Bayly 1979:112, 124-125, distribution
- Bayly 1984:148
- Bayly 1991: 2829-, DISTRIBUTION

Calamoecia australica G.O. Sars, 1908

card 2

-- Bayly 1992c: 7-, fig. 10, western Victoria, Australia

-- HANER ET AL. 1996: 91

Calamoecia australis (Searle)

CARD 1

---, Bayly 1961. pp. 75-78, fig. 4h, 8a-d

Brunella australis Searle, 1911, pp. 176-7

Brunella viridis Searle, 1911, pp. 177

Brunella expansa Sars, 1912, pp. 9-10

B. -- , Brehm, 1953a, pp. 44-5

Calamoecia australis (Searle)--Bayly 1962. pp. 254-6,
fig. 2

--, TOLLEY, 1955: 281

--, LINDQUIST, 1961: 62

--, BAYLY 1962c: 7; FIG. 6, VICTORIA, S. AUSTRALIA & TASMANIA;
SYN. C. EXPANSA NEW SYNONYMY

SYN. OF *C. AUSTRALIS*

Calamoecia expansa (G. O. Sars 1912)

CARD 2

Brunella -, G. O. Sars 1912: 8, pl. 8 figs. 1-5, n. sp.,

South Australia

B. ♂, Brehm 1953: 44-45

C. ♂, Jolly 1955: 281

C. ♂, Lindquist 1961: 12

C. ♂, Bayly 1979: 113, 124-5, DISTRIBUTION

C. ♂, Mary 1991: 2823; DISTRIBUTION

Calamoecia canberra Bayly

--- --- Bayly 1962. pp. 258-260, fig. 4

-- Bayly 1984: 148, SOUTH AUSTRALIA; 151, 153-
154, RECORDS

-- Timms & Morton 1988: 162-, N. AUSTRALIA

-- Bayly 1979: 124-5, DISTRIBUTION, TAB. 4

-- Bayly 1991: 2829-, DISTRIBUTION

-- Bayly 1992c: 7-, FIG. II, AUSTRALIA

Calamoecia clitellata Bayly

--- --- Bayly 1962. pp. 260-263, fig. 5

-- BAYLY 1984: 148

-- BAYLY 1979: 124-5, DISTRIBUTION

-- BAYLY 1992c: 7-, FIG. 7, S. AUSTRALIA

-- W. D. WILLIAMS 1985: 89-, FIG. 3-, AUSTRALIA

-- M. C. GEDDES 1976: 1-, VICTORIA, W. AUSTRALIA

Calamoecia elongata Bayly 1979

- Bayly 1979: 107-109, FIGS. 2A-E, N.sp. ♂♀, W. AUSTRALIA;
-- Bayly 1984: 149, Western Australia; 151 124, 125
- MALY 1991: 2829-, DISTRIBUTION
- Bayly 1992c: 7-, FIG. 12, AUSTRALIA
- HAMER ET AL. 1996: 91

Calamoecia expansa - see C. australis

Calamoecia gibbosa (Brehm)

-3

Brunella gibbosa Brehm, 1950, pp. 252-3; 1953a, pp. 41

Calamoecia gibbosa--Bayly 1961. pp. 82-3, fig. 5e-k, 12

C. -, BAYLY 1967: 98-99, TASMANIA

C. -, BAYLY 1984: 148, 152-153, DISTRIBUTION

C. -, BAYLY 1979: 110-111, DISTRIBUTION; 118,

C. -, BAYLY 1979: 124-5, DISTRIBUTION...

TASMANIA; 124-5, DISTRIBUTION

C. -, BAYLY 1991: 2829-, DISTRIBUTION

C. -, BAYLY 1992c: 10-, FIG. 14, TASMANIA, S. AUSTRALIA

C. -, BAYLY 1992c: 10-, FIG. 14, TASMANIA, S. AUSTRALIA

Calamoecia gibbosa newmanensis Bayly 1979

-- BAYLY 1979: 109-111, FIGS. 3A-G, N.S.P. ♀[♂],

WESTERN AUSTRALIA

-- Bayly 1984: 152

CALAMOECIA LONGICORNIS

--, TOLLY, 1955: 281

--, LINDQVIST, 1961: 12

Centropagidae - Calamocia
Calamocia lucasi Brady, ♀, ♂.

— — Brady, 1906, p. 696; pl. 49, fig. 1-10.

— — Bayly, 1961. pp. 63-67, fig. 1A-D, 4A-D, 6A-F

— , Evans, 1970. 86

— , Chapman-Burns, 1976: 137; FIG. 5; TAB. 1, 2.

— , Tolley, 1966: 732, 734; TAB. 4

— , Tolley, 1967: 351

Habitat. New Zealand.

— , Tolley, 1955: 279, 280, 281-283; PL. I; SYN: STEELI

BRUNELLA
STEELI

— , Tolley, 1955: 279, 280, 281-283; PL. I; SYN: STEELI

--, LINDQVIST, 1961: 12, 13, 14, 84; FIG. 2;
SYN: Brunella
Lucasi
STEELI, Limnocalanus

-- UYE 1984a: 45.

— Shiel & Walker, 198 — : 267-268 Australia

— Shiel, 1985: 2138 "

-- BAYLY 1984: 147, 149-153, FIGS. 4-6, AUSTRALIA

-- FORSYTH & JAMES 1984: 803-810, LAKE
OKARO, NEW ZEALAND

Calamoecia lucasi Brady, 1906

card 2

-- Timms & Morton 1988: 162-, N. Australia

-- Bayly 1979: 111, 124-5, DISTRIBUTION

-- Herrzig 1983: 70 -

-- Kawabata 1991: 32

- Epp + Lewis 1979: 129 -

- Jars 1908b: 11 -

- Bayly 1991: 2829-, DISTRIBUTION

- Bayly 1992c: 10; FIG. 15, AUSTRALIA, NEW ZEALAND

Calamoecia salina (Nicholls) 1944

- Brunella salina Nicholls, 1944, pp. 6-8
- Calamoecia salina (Nicholls)--Bayly 1961, pp. 80-81,
fig. 2a-f, 4g, 10a-e.
- C. salina Bayly — Williams, et.al., 1984: 1210
tbl. 2
- C. -, BAYLY 1984: 148

C. -, BAYLY 1979: 124-5, DISTRIBUTION

C. -, BAYLY 1992c: 5-, FIG. 5, S. AUSTRALIA

C. -, W.D. WILLIAMS 1985: 89-, FIG. 3-, AUSTRALIA

C. -, M.C. GEDDES 1976: 1-, VICTORIA, AUSTRALIA

CALAMORECIA SALMA

--, FARMER, 1980: 359

CALAMOECIA SUBATTENUATA - SEE C. TASMANICA
SUBATTENUATA

= CALAMOECIA Centropagidae - Calanocida. ①
Brunella tasmanica Smith ♂, ♀.

(TYPE SP. OF BRYNELLA N.G.)

— Smith, 1909, p. 87: pl. 18, figs. 6-16.

Bruhn, 1953, pp. 43-44

--, = CALAMOECIA I., TOLLEY, 1955: 279

Habitat. Tasmania.

Calamoecia tasmanica (Smith) G.W., 1909 (2)

-- --, Bayly 1961. pp. 68-73, fig. 2g-k, 3e-j, 4i-j,
5a-c, 7a-d

--, TOLLY, 1955: 281; 54N: Brunella T.

--, LINDQVIST, 1961: 12

-- BAYLY 1963: 216, LOCALITY / ECOL. DATA

-- BAYLY 1984: 149, WESTERN AUSTRALIA

-- BAYLY 1979: 111, 120, TASMANIA; 122 - 125,
FIGS. 8A-B; WESTERN AUSTRALIA; 124-5

-- MALLY 1991: 2829; DISTRIBUTION

Calamoecia tasmanica tasmanica (Smith) ③

---, Bayly 1961. pp. 73-75, fig. 2g-k, 4i-j, 5a-b, 7a-d

--- Bayly 1984: 152-153

--- Bayly 1992c: 127, FIG. 17, AUSTRALIA

Calamoecia tasmanica subattenuata (Fairbridge)

Brunella subattenuata Fairbridge, 1945b, pp. 69-70

Calamoecia tasmanica subattenuata (Fairbridge)--
Bayly 1961. pp. 75, fig. 3e-j, 5c

C--, Bayly 1984: 152-153

C--, Bayly 1979: 109, 122, WESTERN AUSTRALIA;

C--, Bayly 1979: 109, 122,
Also as C. subattenuata, 122-123

C--, Bayly 1992c: 12-

C. subattenuata, Jolly 1955: 281

C. subattenuata, Lindquist 1961: 12, 13

Calamoecia trifida Bayly

-- -- Bayly 1961. pp. 83-87, fig. 1f-h, 4k, 13a-f.

-- -- TIMMS & MORTON 1988: 162-, N. AUSTRALIA

-- -- BAYLY 1979: 124-5, DISTRIBUTION

-- -- BAYLY 1991: 2829-, DISTRIBUTION

-- -- BAYLY 1992c: 10-, FIG. 16, QUEENSLAND, N. AUSTRALIA

Calamoecia ultima (Brehm, 1960)

Brunella -, Brehm 1960b:84-87, figs. 1-5, n. sp. ♀♂,
New Guinea

C. -, Bayly 1962:256-258, fig. 3

C. -, Bayly 1979:124-5, distribution

C. -, Bayly 1980:9-10

C. -, Timms & Morton 1988:162-, N. Australia

C. -, MARY 1991: 2829-, DISTRIBUTION

C. -, BAYLY 1992C: 12-, FIG. 18, N. AUSTRALIA, NEW GUINEA

CALAMOECIA VIRIDIS

--, TOLLY, 1955: 381

--, LINDQVIST, 1961: 12

Calamoecia zeidleri Bayly 1984

-- Bayly 1984: 147-149, Figs. 1, 2, n. sp. ♂♀, South
Australia

- Bayly 1991: 2829-, DISTRIBUTION
-- Bayly 1992c: 7, FIG. 9, S. AUSTRALIA
-- Hamer et al. 1996: 91