

Harpacticoida - Canthocamptidae.

Cancerincola valid

Ameiridae Lang  
Cancerinoidae

Wilson, 1913

C. jamaicensis (M)

Syn. Cancerincola Wilson, 1913, p. 264.

Has no synonyms.

Cancer, a crab and incola, inhabitant

Body 10 seg<sup>s</sup>/5, cylindrical, diminishing slightly  
from in front backwards - 5° and 6° seg<sup>s</sup>/5. more  
distinctly separated than in Canthocamptus.  
10 anten, short, thickset, 8-jointed, strongly  
hinged in male - 2° anten, short, outer ramus  
1-jointed - mand, palp also 1-jointed, 1° max.  
with distinct endo + exopods, the latter armed  
with claw-like spine - 2° max, with only one  
digitiform lobe inside the claw-bearing joint.  
mxpds, normally developed - 1° swim. legs with  
2-jointed endopod, much longer than exopod  
other legs with exopod longer than endopod.  
both rami 3-jointed - only one ovisack.

Cancrincola Wilson, 1913

card 2

- Fiers 1990b:70-72, type of Cancrincolidae n. fam.,  
and key to spp.

*Cancrincola abbreviata* Humes

1957--- Humes, A.G., pp. 185-190, pl. 3 figs. 30-38.

W. coast of Africa, in gill chambers of Sesarma huyardi (Desmarest),  
+ Sesarma angolense Brito Capello

--, PETKOVSKI, 1978: 100, 102

- FIEBS 1930b: 72, 80

Harpacticoida - Canthocamptidae

Canthocamptus jamaicensis Wilson ♂, ♀.

— —, Wilson, 1913, p. 264: pl. 50, fig. 281;  
pl. 51, fig. 282-283; pl. 52, fig. 289-302.

— —, LANG, 1948, p. 843.

— —, Humes, 1957, pp. 180-183, pl. 1, fig. 1-13, pl. 2, fig. 14-17.

Gill chamber of <sup>Sesarma huzardi (Desmarest)</sup>  
Cardisoma armatum Hirklots, w. coast of Africa.

Host. Gills of white land crab - Cardisoma  
granulatum -

East Indies.

Cancrincola jamaicensis Wilson

card 2

--, Petkovski, 1973, pl. 125

--, Schmitt, 1954, p. 440.

--, PETKOVSKI, 1978: 99, 100, 101, 102, 103-108 ♀; FIG. 1-9

- FIEBS 1990b: 70-80, FIGS. 1-6, ♀♂ COPEPODIDS;  
ARUBA, BONAIRE, ZAIRE

*Cancrincola longiseta* Humes

1957--- Humes, A.G., pp. 183-185, pl. 2, figs. 18-22,  
pl. 3. Gill chamber of Goniopsis cruentata (Latreille),  
west coast of Africa, also Sesarma curvatum (M. Edw.),  
+ S. huzardi (Desmarest).

--, PETKOVSKI, 1978: 100, 103

- FIERS 1990b: 72, 80-86, FIGS. 7-11 ♀♂ COPEPODIDS;  
ZAIRE-BANANA.

*Cancrinicola plumipes*, Humes

\_\_\_, Humes, 1941, p. 370, Host-gill-chamber Sesarma,  
Louisiana, 1 fig.

--, Schmitt, 1954, p. 440.

--, PETKOVSKI, 1978: 100, 102

- FIEBS 1990b: 72, 86

CANCRIINCOLA

WILSONI

PEARSE

— —, LANG, 1948. P. 843.

--, HTO, 1971: 320

HABITAT: JAPAN: ABURATSUBO.