

Ascidicolidae - Notodelphyoidea -
Ceratrichodes = synonym. Cyclopoidea

Hesse, 1866, c.

C. albidus (M)

Syn. Ceratrichodes Hesse, 1866, c, p. 75.

A synonym of Botrillophilus, Hesse
(Schellenberg, 1921^b, p. 281)

Κέρασ, horn, antenna, τριχάδης, hairy

Body narrow and elongate; thorax 6-jointed, not including the cephalothorax. Abdomen 8-jointed, very short and imbricated, armed with stout hairs and ridges. Abdomen appendages terminating in long slender hairs. First maxillae long and slender, armed with a claw, the others stouter and larger. Swimming legs biramous, armed with spines and hairs. Genital openings bordered with a denticulate ridge. Antennae with a flat appendage at the base, covered with hairs

Female pear-shaped, thorax 7-jointed, abdomen 3-jointed; abdomen appendages flat, armed with 4 curved claws; no eye.

CERATRICHODES HESSE

CARD 2

-, ILLG + DUDLEY 1980:141, - OF SCHELLENBERG, 1922

NOT SYNONYM OF BOTRYLLOPHILUS.

Asterocheridae - Cyclopoidea.

Ceratrichodes albidus Hesse, ♂.♀.

Hesse, 1866, c, p. 75: pl. 4, f. A, 1-11.

Host. A red social Ascidian.

French Coast.

Asiderocheridae - Cyclopoida.
Ceratrichodes flavus Hesse, ♂, ♀.

— Hesse, 1871, p. 26;

Host. A social ascidian.

France.