

m) Brancheletia Arguloida - Argulidae.

Dipteropeltis - valid -

W. T. Calman 1912.

D. hirundo (M)

Syn. Dipteropeltis Calman, 1912, p. 34.

Dipteropeltis Calman, 1912<sup>b</sup>, p. 84.

Talaus Moreira, 1913, p. 147.

Moreiriella Leitaõ, 1913, p.

Diptera = 2-winged and peltis = carapace.

alluding to the greatly elongated lateral wings of the carapace.

Differing from Argulus in having no spine on the preoral papilla; in having the antennules and antennae very minute and imperfectly segmented; in having no large spines or hooks on the under surface of the carapace, body, or appendages; in having no furcal rami on the abdomen, and in having the lateral wings of the carapace greatly elongated.

Dipteropeltis CALMAN, 1912

CARD 2

- , Hessler, 1969, p. 205
- FRYER 1978b: 368-373
- CRESSEY 1978:1
- CRESSEY 1983:252
- FRYER 1956b: 338-
- THATCHER 1991: 494, DIAGNOSIS

Dipteropeltis hirundo. Calman - ♀. Argulidae

— —, Calman, 1912a, p. 34, May 21st. <sup>Paraguay</sup> R.

— —, Calman, 1912b, p. 763, pl. 84, Dec.

— —, Moreira, 1915, p. 210.

— — Ringuélet 1943 p. 97, pl. 13. figs. 1-5 ♀  
p. 89

Host "Dorado" (Salminus sp.)

— — Carvalho, 1941, p. 267-272, pls. 19-20. On

Tetragonopterus areolaris ~~Paraguay River.~~

+ T. rutilus, tributary to Parana R., S. Paulo, Brazil.

--, Kabata, 1970, p. 12- fig. 3

-- THATCHER 1991: 494, FIG. 9-27, HOST LIST

Argulidae

Dipteropeltis hirundo. Calman - ♀.

— —, Calman, 1912a, p. 34, May 21st. <sup>Paraguay</sup> R.

— —, Calman, 1912b, p. 763, pl. 84, Dec.

— —, Moreira, 1915, p. 210.

— — Ringuélet 1943 p. 97, pl. 13 figs. 1-5 ♀  
p. 89

Host "Vorado" (Salminus sp.)

— — Carvalho, 1941, p. 267-272, pls. 19-20. On

Tetragonopterus curassavicus ~~Paraguay River.~~

+ T. rutilus, tributary to Parana R., S. Paulo, Brazil.

--, Kabata, 1970, p. 12- fig. 3

-- THATCHER 1991: 494, FIG. 9-27, HOST LIST