

#10
Doropygidae - Notodelphyoidea
Doropygus, valid. Notodelphyidae: Cyclopoidea
Thorell, 1859. D. pulex.

Syn. Doropygus Thorell, 1859, p. 43.
Doropygus Brady, 1878, p. 133.
Doropygus Sars, 1921, p. 42.
Doropygus Schellenberg, 1921^b, p. 238.
Δορός = a leather bag & πυγή = the rump.

Body wider than in Notodelphys; cephalothorax a little compressed (♀) or subterete (♂): anal laminae furnished with curved spines at v tips.

1st antennae 8-10 jointed: 2nd pair with an immovable hook at v tip. Basal part "palp twice as wide as the inner ramus - endopod 2-jointed, exopod one-jointed. Maxillae compressed; 1st pair armed with 5 setae, hairs, or spines: 2nd pair sub-triangular, 5-jointed: 3rd pair oblong: both pairs armed on v inner margin with setae or spines.

Thorax feet compressed, with non-plumose or sparsely plumose setae. Endopod of 4th pair with 3 or 2 joints - 5th pair rudimentary & 4 ovaries, joined in pairs anteriorly!

Live in the respiration sac of Ascidians + Cyntheia.

Notodelphyidae: Cyclopoida

Doropygus Thorell 1859

(2)

- Stock 1967e: 20-21

DOROPHYGUS, KOGA 1968: 19, EGGS

- ILLG + DUDLEY 1980: 7, 24

- IZAWA 1987: 160-

- GOTTO 1993: 32, DIAGNOSIS

Doropygidae - Notodelphyoidea -

Doropygus acutus Hesse, ♀.

— Hesse, 1866, c. p. 64.

Host. A small yellow Ascidian.

French Coast.

Doropygidae - Notodelphyoidea

Doropygus albidus Hesse, ♀

— Hesse, 1866, c. p. 61.

Host, Ascidia intestinalis.

French Coast.

Doropygidae - Notodelphyoidea.

Doropygus antarcticus Schellenberg, ♀.

— — Schellenberg, 1921^b, p. 243;
1922?

-- Stock 1967e: 70

Host. Paramolgula gigantea.

Falkland Islands.

Doropygus apicatus Stock 1967

- Stock 1967e: 3, 23-27, figs. 25-30, n. sp. ♀♂, from Ascidia nigra, Red Sea.
- Illg 1970: 182
- Jones, J.B. 1973: 262
- Jones 1979: 540

Doropygus arcticus Aurivillius ♀

- — Aurivillius, 1885, p. 233: pl. 8. figs. 1-11.
- — Wilson, 1920, p. 14 T.
- — ILLG 1958: 587, INCERTA SEDIS

Host. Branchial sac of Chelyosoma mac-
leanum Sow. & Brod.
arctic Ocean.

Notodelphyidae

Doropygus auritus Thorell. ♂♀.

- —, Thorell, 1859, p. 50: pl. 7: pl. 8, fig. 10.
— —, Hesse, 1866, c, pp. 54 & 64.
— —, Aurivillius, 1882, p. 55: pl. 6, fig. 13:
p. 61; pl. 13, fig. 12: pl. 15, figs. 1-2.

Host. Phallusia obliqua: T. mentula:
P. venosa:

Mediterranean
N. Atlantic.

- —, Aurivillius, 1883, p. 25: pl. 2, fig. 13.
p. 57: pl. 4, fig. 12: pl. 6, figs. 1-2.
- — Brady, 1878, p. 135: pl. 29, figs. 1-11.
- — Carus, 1885, p. 343.
- — Norman, 1869, p. 299.
- — T. Scott, 1907, p. 364.
- — Thorell, 1859, a, p. 339.
- — transferred to genus Notopterophorus, Sars, 1921,

p. 53

Doropygus auritus
var. elongatus (Buchholz), ♀

Notopterophorus veranyi Teuckart, 1859, p. 241.
" " " elongatus Buchholz, 1869, p. 127.
Doropygus — —, Aurivillius, p. 57: pl. 6, fig. 3.

Host. Phallusia mentula: P. mamillata

Mediterranean

— — *canis*, 1005, p. 343.

— — var. e — Aurivillius, 1882, no. 8, p. 61: pl. 15, f. 3.

Doropygus bayeri Illg

1958--- Illg, P.L., pp. 544-550, fig. 6, WASHINGTON
1966 -- Dudley, P., pp. 105; STATE

-- IZAWA 1987: 155-

Dogorygus brevipes Ho

→ Ho, 1984b: 31-37, fig. 5-8

-- Ooishi 1987: 95, JAPAN

Doropygiidae - Notoaelephyoidea.

Doropygus callipygus Hesse, ♀.

-- Hesse, 1866, c. p. 58,

Host. Ascidia venosa.

French Coast.

Doropygus catalai Illg, 1970

n. sp. Illg, 1970, pp. 177-183, figs. 3-5, NOUMEA,
NEW CALEDONIA

--Jones, J.B., 1973, p. 262.

--, Jones, 1979: 540

Doropygidae - Notodelphyoidae

Doropygus coccineus Hesse, ♀.

— — Hesse, 1866, c. p. 67.

Host. A small yellow Ascidian.

French Coast.

Doropygidae - Notodelphyoidea

Doropygus conicus Hesse ♀.

--- Hesse, 1866, c. p. 57.

Host, Cynthia microcosmus on the valves
of a Pecten maximus.

French Coast.

Notodelphyoidea.

Doropygus cristatus. Hesse, ♀.

— Hesse, 1871, p. 21; pl. 2, fig. 18.

Transferred to genus Notopterophorus (Wilson, 1935)

Host. Ascidia canina

Mediterranean.

Doropygidae - Notodelphyoidea.

Doropygus curculio Hesse, ♀.

— Hesse, 1866, c, p. 54.

Host, "A simple green Ascidian"

French Coast.

Doropygus curvatus Gray, 1938

-- Dudley & Illg 1991:6-, Figs. 9, 17, 25; review records

DOROPYGUS CURVIPES GOTTO, 1975

--, Gotto, 1975, p. 169-171, fig. 3, AUSTRALIA

--, JONES, 1979: 540; TAB. 1

Notodactylidae - Notodelphyoidea.

Doropygus cylindriciformis Schellenberg-

— Schellenberg, 1921^b, p. 241: text-figs. 21-22.

-- Stock 1967 e: 26

Host. Ascidia canelata.

Gulf of Suez.

Doropygus dakarensis Monniot, 1966

— Monniot, 1966: 651-655, fig. 1-2.

-- Jones, J.B., 1973, p. 262.

--, JONES, 1979: 540

--, Gotto, 1975, p. 169

Doropygidae - Notodelphyoidea

Doropygus deflexus Hesse, ♀.

— — Hesse, 1866, c. p. 58

Host. Cortical envelope of a zoophite -

French Coast.

Doropygus demissus ^{Nolodelphyon} Aurivillius, ♀ ①

— Aurivillius, 1885, p. 230: pl. 7, figs 14-24.

— Wilson, 1920, p. 142.

-- Jones, J.B., 1973, p. 272.

-- ILLG & DUDLEY 1980: 123

Host. Respiratory sac of Cynthia echinata.

Arctic Ocean.

Doropygus demissus Aurivillius, 1885

card 2

-- Prefontaine & Brunel 1962:253-254, St. Lawrence,
host Boltenia ovifera

-- DUDLEY & ILLG 1991: 7-, FIGS. 3, 4, REVIEW RECORDS;
MAINE, CAPE COD, MASS., NOVA SCOTIA, NEW BRUNSWICK

Doropygus depressus Stock, 1967

-- Stock, 1967e: 3, 27-28, figs. 31-33, n. sp. ♀, from

Herdmania momus, Red Sea.

-- Jones, J.B. 1973: 262, 266

Doropygus elegans Ooishi, 1963

--Jones, J.B., 1973 p. 262.

--, JONES, 1979: 534

-- OOSHII 1987: 95, JAPAN

Doropygus elongatus Giesbrecht,
var. maculatus Gourret,

— — Gourret, 1888, p. 52; pls. 2, 3, 4.

— — Gourret, 1887, p. 186.

A synonym of Notopterophorus e-m-.

Doropygus fernaldi Illg

1958---Illg, P.L. pp. 551-557, fig. 7, WASHINGTON SOUND,
WASHINGTON

1966 -- Dudley, P., pp. 110;

-- IZAWA 1987: 155 -

DOROPYGUS FLEXUS GOTTO, 1975

--, Gotto, 1975, p. 167-169, fig. 2

--, Jones, 1979 : 540 ; TAB. 1

Doropygus gibber Thorell, ♂. ♀.

- —, Thorell, 1859, p. 52: pl. 8, fig. 11.
— —, Hesse, 1866, c, p. 63.
— —, Buchholz, 1869, p. 120: pl. 7, fig. 4, A. B. C.
— —, Aurivillius, 1882, p. 56: pl. 6, figs. 11-12.

Host. Phallusia rustica: Ascidia
canina Heller: P. obliqua: P. patula
Ciona intes tinalis: Clavellina sp.
Mediterranean - N. Atlantic.

- —, Aurivillius, 1883, p. 26: pl. 2, figs. 11-12
— —, Carus, 1885, p. 343.
— —, Brian, 1898, p. 9.
— —, T. Scott, 1907, p. 364.
— —, Thorell, 1859, a, p. 339.
— —, Stephensen, 1913, p. 349.
— —, Wilson, 1920, p. 14 L.
— —, Thompson, 1893, p. 190: pl. 18. f. 2, a-b.
Notopterophorus — Pesta, 1909, p. 258: pl. 1, figs. 1-6
— —, Schimkewitsch, 1896, p. 342: pls. 15 & 16.

DOROPYGUS GIBBER THORELL

CARD 2

--, ILLG + DUDLEY 1980:150, D.G. OF VANHÖFFEN 1897

= SCHIZOPROCTUS INFLATUS

-- IZAWA 1987:154-

Doropygidae - Notodelphyoidea.

Doropygus gibbosus Hesse, ♀,

— Hesse, 1866, c, p. 62.

Host. Ascidia intestinalis,

French Coast.

Doropygidae - Notodierpygidae
Doropygus globosipherus Hesse, ♀.

— Hesse, 1870, p. 307.

Habitat. Compound Ascidian.

French Coast.

Doropygus globosus J.B. Jones, 1974

-- Jones, J.B. 1974: 256, 262, 268-273, figs. 9-12, n.sp.
female only, ex Corella eumyota, Wellington Harbor,
New Zealand

-- Jones, J.B., 1979: 533

Doropygus humilis Stock, 1967

-- Stock 1967e: 21-27, figs. 21-24, n. sp. ♀, from tunicates (Pyuridea), Red Sea.

-- Jones, J. B. 1973: 262.

-- Jones 1979: 540, Tab. 1

Doropygus hummi Illg

1958---Illg, P.L. pp. 557-562, fig. 8.

DOROPYEUS KERGUELENSIS SCHELLENBERG, 1922

--, TONES, 1979: 534, 538

Doropygus lamellipes Schellenberg. ♀. ♂.
Notodelphyidae - Notodelphyoidea

— — Schellenberg, 1921^b, p. 242; text-fig. 23.
(1922?)

— — Stock 1967e: 20

Host Styela paessleri.

Falkland Islands.

Doropygidae - Notodelphyoidea.
Doropygus laticornis, n.sp. ♂, ♀.

— Wilson, 1931.

— Wilson, 1932, p 388: pls 19; 24.

-- DUDLEY, P. 1986: 18-, FIGS. 5, 7i, j

-- ZWERNER + LAWLER 1972: 83, CHESAPEAKE BAY, MOLGULA

Host. Branchial cavity Molgula manhattanensis, Woods Hole.

-- DUDLEY + LEEG 1991: 6- , FIGS. 2G, H, 22; REVIEW RECORDS; CAPE COD, MASS., GEORGIA

Doropygus longicauda Aurivillius, ♀.

- —, Aurivillius, 1882, no. 3, p. 48: pl. 7, f. 1-12
— — " " , 1882, no. 8, p. 60: pl. 13, f. 11.
— — " " , 1883, p. 18: pl. 3, figs. 1-12: ^{p. 50}
— — " " , 1886, p. 43: pl. 1, f. 1-6.

— —, ~~Host~~, ~~1882~~, ~~no. 3~~
Host. Phallusia (Ascidia) obliqua and
P. mentula.

Bohuslan-

made type, of n. gen. Doropygopsis. Sars 1921, p. 47.
Doropygopsis — Hansen, 1923, p. 22.

~~Stephensen, 1940, p. 2~~

Doropygidae - Notodelphyoidea.

Doropygus longimatrix Schellenberg, ♀.

— — Schellenberg, 1921^b, p. 245; text-figs. 24-5.

--, Illg, 1970, p. 182

--, JONES, 1979: 540

-- STOCK 1967 e 126

Host. Pyura momus.

Gulf of Suez.

DOROPYGUS LOUISAE JONES, 1979

-- JONES, 1979: 538-540 ♀; FIGS. 4, 5; EX PYURA
CANCELLATA (ASCIDIAN); TAB. 1

Doropygidae - Notodelphyoidea.

Doropygus macroon Hesse, ♀.

— — Hesse, 1866, c. p. 66.

Host, "A green Ascidian".

French Coast.

Doropygus mirabilis McKinnon

— McKinnon, 1984: 1-5 , fig 1-3

Doropygus mohri Illg

1958---Illg, P.L., pp. 562-568, fig. 9.

1966 -- Dudley, P., pp. 108;

-- IZAWA 1987: 155-

Doropygus molgulensis Pearse

--- Pearse, A.S., 1952, pp. 188-189, fig. 1, N. SP.,

HOST MOLGULA OCCIDENTALIS, FLORIDA GULF
COAST (ALLIGATOR HARBOR)

--- ILLG 1958:588, "INDETERMINABLE SPECIES"

Doropygidae — Notodelphyidae

Doropygus normani Brady, ♀

— — Brady, 1878, p. 136: pl. 32, figs. 1-14

— — Carus, 1885, p. 342.

— — Gouret, 1888, p. 1.

— — T. Scott, 1902, b, p. 455.

Host. Cynthia papillosa: "Ascidians".

Mediterranean: Indian Ocean.

— — T. Scott, 1906, b. p. 363:

— — T. Scott, 1907, p. 364.

— — Thompson & Scott, 1903, p. 255.

— — GOTTO 1957: —, ? SYNONYM OF D. PSYLLUS

Doropygidae - Notodelphyoidea.

Doropygus novae seelandicus. Schellenberg, ♂, ♀.

— — Schellenberg, 1921^b, p. 244.

-- Jones, J.B., 1973, pp. 255-257.

--, JONES, 1979: 533

-- Stock 1967e: 20.

Host. Pyura pulla: Paramolgula filholi:
Cnemidocarpa novae seelandiae

New Zealand.

Doropygidae - Notodelphyoidea

Doropygus novemsetiferus (Schellenberg,)[♀]

- 1922?

— Schellenberg, 1921^b, p. 244:

-- Stock 1967e: 28-29, FIGS. 34-37, FROM ASCIDIANS
(PYURIDAE), RED SEA.

Host. Styela canopus; Pyura gangelion.

Gulf of Suez.

Doropygidae - Notodelphyoidea

Doropygus oblongus Hesse, ♀.

— Hesse, 1866, c, p. 59.

Host. Polyclinium stellatum (mollusk).

French Coast.

Doropygidae - Nordaefpnygidae

Doropygus papilio massiliensis Gourret

— — — Gourret, 1887, p. 185.

— — — Gourret, 1888, a. p. 36: pls. 2, 3, 4.

Host. Phallus sp.

Doropygus pinguis (Ooishi)

—, Ooishi, 1962:

—, Ho, 1984b: 24-31, figs 1-4.

-- Ooishi 1987: 95, JAPAN

See
Doropygus
pulex
pinguis
Ooishi 1962

Doropygus platythorax

--Jones, J.B., pp. 256, 262-267, figs. 5-8.

--, JONES, 1979: 533-534 ♂; FIG. 1; EX CNEMIDOCARPA
DISIOTIS

Doropygidae — Notodelphyocidae

Doropygus porcicauda Brady, ♀.

figs. 14-16

— — Brady, 1878, p. 138: pl. 27, figs. 1-9: pl. 33, n

— — Norman & Brady, 1909, p. 154.

— — T. Scott, 1906. b. p. 363.

— — T. Scott, 1907, p. 364.

Host. Ascidia parallelogramma.

French Coast;

- Sars, 1921, p. 45: pl. 22.
- Thompson, 1893, p. 190: pl. 18, f. 1, a-b.
- GOTTO 1957: 282

Doropygus profundus Illg

1958---Illg, P.L., pp. 569574, fig. 10.

Doropygidae-Notodelphyoidea

Doropygus postremoglobosus Hesse, ♀.

— Hesse, 1871, p. 23; pl. 3. fig. 12.

Host. Ascidia canina

French Coast.

Doropygidae - Notodelphyoidea

Doropygus propinquus Hesse, ♀.

— Hesse, 1866, c. p. 56.

Host, A simple Ascidian.

French Coast.

Doropygus psyllus Thorell, ♀

p. 339

- —, Thorell, 1859, p. 49: pl. 7, fig. 9: 1859, a,
 — —, Hesse, 1866, c, pp. 54 & 64.
 — —, Aurivillius, 1882, No. 3, p. 54.
 — —, Aurivillius, 1883, p. 24.

Host. Phallusia virginia: P. patula.

Mediterranean.

— — Carus, 1885, p. 342

— — Gouret, 1888, p. 1.

— — T. Scott, 1907, p. 364.

— — Sars, 1921, p. 44: pl. 21.

— — Brian, 1924, p. 7.

— — GOTTO, 1957a: —, HOST ASCIDIA CONCHILEGA,
U. K., NEW HOST RECORD

Doropygus pulex Thorell, ♀ Notodelphyoidea

- —, Thorell, 1859, p. 46: pl. 6. fig. 8.
- —, Hesse, 1866, c, p. 55.
- —, Aurivillius, 1882, No. 3, p. 54.
- —, Aurivillius, 1883, p. 24. [European Seas]

Host. Phallusia obliqua : P. mantula :
P. patula : P. virginia : Corella
parallelogramma : Styela gyrosa.
P. conchilegia : Cynthia loricata : C. rustica.

- — Brady, 1878, p. 133: p. 20, 11
 — — Carno, 1885, p. 342: Canu, 1892, p.
 — — Kerschner, 1879, p.
 — — Brian, 1905, a, p. 2: pl. 4, figs. 10-12.
 — pullus Buchholz, 1869, p. 116: pl. 6, §. 3: pl. 7, §. 3.
 — pulex Norman & Brady, 1909, p. 153.
 — — T. Scott, 1907, p. 363.
 — — Thompson & Scott, 1903, p. 255.
 — — Thorell, 1859, a, p. 339.
 — — Sars, 1921, p. 42: pl. 20
 — — Schellenberg, 1921^b, p. 246: text fig. 26
 — — Thompson, 1893, p. 189: pl. 17, f. 8, a-b.
 — — ~~1118~~ & DUDLEY, 1965, pp. 379-385

Doropygus pulex

D. pulex

(2)

— — Stephensen, 1929, p. 6. Faroes.

— — Wilson, 1932, p. 319: fig. 239.

-- --, Illg & Dudley, 1961. pp. 39-45, fig. 13-14

--, Illg, 1970, p. 182

--, Gotto, 1966, p. 192

--, Lejeune & Monniot, 1965, p. 615

--, Wailes, 1935, p. 6

-- Jones, J.B., 1973, pp. 255-256.

-- Hamond, 1973, p. 349

--, Jones, 1979: 533, 540

--, Gotto, 1975, p. 165

--, Stock, 1969, p. 220-221

- Monniot, 1961: 97-~~100~~, fig. 1 b, tab. 1
- Monniot, 1966: 441
- Stock 1967e: 22-23; POSSIBLY A COLLECTIVE SPECIES?; 26

DORYPYGUS -, KENSLEY & GRINDLEY 1973: 122

-- GOTTO 1957: 282

-- HUMES 1987b: 106

Dorpygus pulex

card 3

-- PREFONTAINE & BRUNEL 1962: 254, 5T.
LAWRENCE, HOST BOLTENIA OUIFERA; DUBIOUS IDP

- Holmes and Gotto, 1987:341, Host: Polycarpa rustica and Molgula complanata (Ascidians)
Co Galway

-- GOTTO 1957:--

-- DUDLEY & LEE 1991: 23; FIG. 20; REVIEW RECORDS

-- C.B. WILSON 1935: 779, TUNICATE, CORONADEL
MAN, CALIFORNIA

-- GOTTO 1993: 32-, FIG. 5 F-I

-- RAIBAUT 1985: 266, FIG. 19B

Doropygus pulex pingius Ooishi, 1962

--Jones, J.B., 1973 pp. 262.

Doropygus pulex pinguis

- Jones, 1979: 540; Tab I
- Stock, 1967e: 23

Notodrilus pullus

Doropygus pullus Buchholz, ♂. ♀.

— — Buchholz, 1869, p. 116: pl. 6: fig. 3: pl. 7, ^{fig. 3}

— — Brian, 1905, a, p. 2 - made a synonym
of Doropygus pulex.

Host, unknown.

Mediterranean.

Doropygus reductus Stock, 1970

--Jones, J.B., 1973, p. 262.

Doropygidae - Notodierpny...

Doropygus reflexus Hesse, ♀,

— Hesse, 1866, c. p. 65.

Host. Ascidia venosa.

Mediterranean

Doropygus rigidus Ooishi, 1962

--Jones, J.B., 1973, pp. 262,267.

--, JONES, 1979: 534

-- Stock 1967e: 28

-- OOSHII 1987: 95, JAPAN

Notodelphyoida- Doropygidae

Doropygus robustus Pearse, 1952

1952--- Pearse, A.S., pp. 189- 191, figs. 2-8, n. sp.,

♀ ONLY, HOST STYELIA PLICATA, ALLIGATOR
HARBOR, FLORIDA

Doropygidae - Notodelphyoidea,

Doropygus rotundus Hesse, ♀,

— Hesse, 1866, c, p. 60.

Host. Ascidia aspera.

Mediterranean.

Doropygidae - Notodelphyoidea.

Doropygus rufescens Hesse, ♀.

— — Hesse, 1866, c. p. 67.

Host. A reddish Eucoelium (mollusk).

French Coast.

Doropygus schellenbergi Illg

1958---Illg, P.L., pp. 574-579, fig. 11.

- DUDLEY + ILLG 1991: 23 -, FIG. 21; REVIEW RECORDS

Doropygus seclusus Illg

1

1958---Illg, P.L., pp. 579-586, fig. 12.

1966 --- Dudley, P., pp. 65;

--, ILLG 1968: 7-42; X Pls.

-- DO ET AL. 1984: 25-26

-- OISHI 1980: 286.

-- IZAWA 1987: 155 -

-- DUDLEY, P. 1986: 15-, FIG. 3A, B

-- HIRFAU - JACQUOTTE 1984: 65, "METAMORPHIC ♂" ? =

~~DORO~~

Doropygus seclusus Illg

card 2

-- Dudley & Illg 1991:15

-- RAIBAUT 1985: 257-, FIG. 15

-- GOTTO 1990: 308

Doropygidae - Notodelphyoidae

Doropygus sphaerispherus Hesse. ♀.

— — Hesse, 1870, p. 305.

Habitat. Compound Ascidian.

French Coast.

Doropygidae - Notoxipha
Doropygus spiniferus Schellenberg, ♀.

— — Schellenberg, 1921^b, p. 248;

Host. Molgula nodosa.

S. W. Australia.

DOROPYGUS SPINOSUS JONES, 1979

-- JONES, 1979: 534-538 ♀; FIGS. 2, 3; EX

ASTEROCARPA CERA (ASCIDIAN)

Doropygidae - Notodelphyidae

Doropygus thorelli Aurivillius

14, figs. 1-14

— Aurivillius, 1882, p. 49: pl. 13, fig. 10: pl.

— " " " , 1883, p. 45: pl. 4, f. 10: pl. 5, f. 1-14.

made type of n. gen. Doropygella.
Sars, 1921, p. 49.

Hosts, Phallusia mentula.

Bohuslän.

Doropygidae - Notodactylidae.

Doropygus trisetosus Schellenberg, ♀.

— — Schellenberg, 1921^b, p. 249:

— — Stephensen, 1927, p. 379: fig. 33, 1-4, text.

-- Jones, J.B., 1973, pp. 255-256.

--, Jones, 1979: 533, 534, 538

Host. Polycarpa pegasi.

New Zealand.

Doropygidae - Notodelphyoida

Doropygus trisetosus;

var. kerquelensis Schellenberg, ♀.

— — Schellenberg, 1921^b, p. 249: text-fig. 28.

Host. Molgula sp.

Kerguelen.

Doropygidae - Notodelphyoidea.

Doropygus tumefactus Hesse, ♀.

— Hesse, 1866, c, p. 63.

Host, "A brown Ascidian".

French Coast.

Doropygidae - Notodierpnyocidae,

Doropygus verrucosus Hesse, ♀.

— Hesse, 1866, c. p. 60.

Host, Ascidia venosa.

French Coast,

Doropygidae - Notodelphyoidea

Doropygus viridis Hesse, ♀.

— Hesse, 1866, c, p. 61,

Host. Cynthia microcosmus.

French Coast.