

Cletodidae-Harpacticoida

Ilyophilus, synonym -

Lilljeborg, 1902.

= I l i o p h i l u s

Syn Ilyophilus Lilljeborg, 1902, p. 61-62

Ilyophilus Sars, 1909, p. 47.

Not
"y"

A synonym of Nannopus Brady 1880.

Body cylindrical: cephalic segt large: rostrum
plate rounded, not defined at base.

Urosome attenuated behind: gen segt divided:
caudal rami not lamellar, one apical seta spi-
niform and stout. First antennae in ♀ 5-
segt, thickly clothed with spiny setae, those
of the ♂ hinged. Endopod of 2nd antennae
strong, end segt armed with claw-like spines: ex-
opod 1-segt. Mandibular palp 1-segt: mx. and mxpds
short and stout. First legs not prehensile: endopod
of 1st, 2nd, and 3rd legs 2-segt and smaller than the
exopod, of 4th legs 1-segt and rudimentary: 3rd
legs transformed in ♂, 5th legs small, distal segt
sometimes distinct, sometimes fused with basal segt:
basal expansion broad but not projecting -
One ovisac.

Clerodidae - Mariposa
Ilyophilus flexibilis Tulljeborg, ♀.

— — Tulljeborg, 1902a: 62 —

A synonym of Nannopus palustris.
(Sars, 1909, p. 307).

Cletodidae Harpacticoida

Ilyophilus perplexus Sars. ♀.

— — Sars, 1909, p. 47: pl. 13, figs. 93-104.
—, Kiefer, 1956, p. 265

Habitat. Lake Tanganyika.