

Canthocamptidae-Harpacticoida

ⁱⁿ⁾
Leimia

Willey, 1923.

L. vaga ^(M)

Syn. Leimia, Willey, 1923, p. 314.

1° ant. 6-jointed - 2° ant. 2-jointed, exopod
1-jointed but large and conspicuous.
Md. palp. simple, bipartite. Mx. like that
of *Canthocamptus*. 2° mx. "with 2 digitiform
lobules inside the unguiferous joint."
Mxp. with simple hand and delicate claws.
10 leg non-prehensile - all 4 pairs have 3-
jointed ex. and 2-jointed en. rami equal
in 1° leg - en. much shorter in 2°, 3°, 4°.
5° leg 2-jointed, basal lobe produced outside
with long seta - inner expansion with 5 setae
distal lobe with 4 setae.
Furca longer than anal segm.

Leimia Willey, 1923

card 2

- Por 1986: 423, transferred from Cletodidae to
Canthocampidae incertae sedis.

- Hays 1992: 71-

LEIMIA DUBIA WELLS - SEE BORFOLIMELLA

Canthocomptidae e. Harpacticoid

Leimia vaga Willey, ♀

313, 314-317, 320

— — Willey, 1923, p. 314; figs. 8-12, text.

— —, LANG, 1948, p. 1320, pl. 534:1.

— —, Coffin, 1978-79, p. 593

— — SHIH ET AL. 1971: 51, CANADA-ATLANTIC

Habitat. Scotsman Bay, Nova Scotia.
+ NEW BRUNSWICK.