

~~Asterocauleridae - Cyclopoida.~~

ENTOMOLEPIDIDAE - SIPHONOSTOMATOIDA

Lepeopsyllus, valid

Thompson & Scott, 1903.

L. typicus.

Syn. Lepeopsyllus, T. & S., 1903, p. 290.

- McKinnon 1988: 996 -

Λέπος, a scale and ψύλλος, a flea.

Body oval, thin, 4-jointed. Abdomen 3-jointed, covered by last thorax segt. Furca partly or completely covered by same segt. Margin of the carapace lined with papilla-like prolongations of irregular length. 1st antennae 13- to 15-jointed. Exopod of 2nd. antennae 4-jointed, endopod long and blade-like. Mandible rudimentary palp 2-jointed. Both rami of first 3 pairs of legs 3-jointed; exopod of 4th. pair 3-jointed, endopod a minute knob. 5th. pair each a long, curved, hairy appendage.

Entomolepidae ~~Asterocheridae~~ - Cyclopoidea,

Lepeopsyllus ovalis Thompson & Scott, ♀.

I.C.

-- Thompson & Scott, 1903, p. 291: pl. 19, f. 30-33.

-- , Eiselt, 1959, p. 659

-- Stock 1992a: 67

-- McKinnon 1988: 996-997

-- Johnson 2000: 119, in key

Habitat, General washings, Ceylon.

Entomolepidae ~~Asterocheridae~~ - Cyclopoidea,

Lepeopsyllus typicus Thompson & Scott, ♀.

In C.

A.

-- Thompson & Scott, 1903, p. 290: pl. 19, fig. 21-29.

-- , Eiselt, 1959, p. 658.

-- Stock 1992a: 67

-- McKinnon 1988: 996-1997

-- Johnson 2000: 119, in key

Habitat, Pearl oyster washings, Ceylon.