

Lucicutia. valid  
Giesbrecht, 1898

L. flavigornis<sup>♂</sup>.

Lucicutia Giesbrecht, 1898, p. 110.

Leuckartia Claus, 1863, p. 182.

Leuckartia Brady, 1883, p. 50 (pars).

Leuckartia Giesbrecht, 1892, p. 62, 358.

Lucicutia Steuer, 1904, p. 596.

Lucicutia Wolfenden, 1904, p. 121.

Lucicutia Esterly, 1905, p. 179

Lucicutia Pesta, 1921, p. 529.

Lucicutia Giesbrecht, 1898      LUCICUTIIDAE

Sewell 1932, p. 285, revision

—, Gardner & Szabo, 1982, p27, 355

— MAUCHLINE 1988: 708-, FIG. 6e, MORE SIG.

— VYSHKVERTZELA 1989b: 14-

Lucicutia anisofurcata M. Heptner, 1971

n. sp., Heptner, 1971, p. 116-8, fig. 21

HUCICUTIA

ANOMALA

Brodsky, 1950

— — — 1950, p. 333, fig. 230.

— — —, 1966, Huelsemann, p. 705;

— — —, Heptner, 1971, p. 118, fig. 22

— — —, Vidal, 1971, pp. 24, 26, 27-, fig. 4.

Lucicutia atlantica Wolfenden, ♀.  
Lucicutiidae - Calanoida

- — Wolfenden, 1904, p. 121  
— — Sars, 1925, p. 217: pl. 59, f. 1-5  
— — Jespersen, 1934, p. 105. W. Greenland, SW Iceland.  
— — Jespersen, 1940, p. 51. S. of Iceland.

Habitat. English Seas; Mediterranean.  
—, (= L. MAGNA), TANAKA, 1963: 36-38  
— GISLASON & ASTTHOR-  
OFF ICELAND SSON, 1992: 221, TAB. I.

LUCICUTIA AURITA CLEVE - SEE L. BICORNUTA



Iucicutia aurita G.O.Sars, ♀,  
Iucicutidae - Calanoida

— Sars, 1905, b, p, 8

Name preoccupied by Cleve in 1904.

Habitat. Station 1768, Monaco (1 specimen).

Lucicutia bella, 1966, Huelsemann;

—Deevey & Brooks, 1977:264

Lucicutia bicornuta Wolfenden, ♂  
Lucicutiidae - Calanoida

l. sp. Wolfenden, 1905, p. 24: pl. 2.

— — Sars, 1907, a, p. 3.

— — A. Scott, 1909, p. 126: pl. 39, figs. 1-11.

— aurita Sars, 1905, b, p. 8.

Habitat. Tropical Pacific: Atlantic.

--; TANAKA, 1963: 26, 27, 51-52 ♀ ♂, FIG. 173

--; OWREY FOY, 1964b: 366

--; OWREY FOY, 1972: TAB. 3

— — Wolfenden, 1911, p. 321: Text-fig. 65, a-c.

— — Sars, 1925, p. 214: pl. 58, f. 1-3.

— — Wilson 1942, p. 192

— — Sewell 1932, p. 290, Indian O.

— —, Brodsky, 1950, P. 339, fig. 237.

— —, Sewell, 1941, P. 113, S. ARABIAN SEA AREA.

— —, 1966, Huelsemann, p. 707;

— —; Park, 1970, p. 477

— , Vervoort, 1965: 115-116 SYNONYM: L. AURITA

— , Gardner & Szabo, 1982, p. 27, 356-357, fig. 22.1.1,  
Syn.: L. aurita.

— , Grice & Huelsemann, 1968: 4b) 2

— , Deepdy & Brooks, 1977: 264

Lucicutia bicornuta Wolfenden, 1905

card 2

-- Mauchline 1988: 708-

-- Soldevilla, et al. 1988: 59 Canary Islands

L. AURITA, HERRING 1988: 184 -

-- COPLEY & WIEBE 1990: 6, FIG. 18a-b, GUAYMAS BASIN

-- RAZOUKS 1994: 10

- *L. BICORNUTA*

Calanoida.

# Lucicutia aurita Cleve.

(3)

L. a. Cleve, 1904<sup>5</sup>, pp. 192, 202-203, figs.?

L. a. Stebbing, 1910, p. 537,  
not — Sars, 1905, p. 8.

L. a., 1966, Huelsemann, p. 705;

L. a. Grie & Huelsemann, 1968: tabl. 2

## Habitat. So. Atlantic.

L.a. CLEVE, 1905: 202; 203, figs.?, n. sp.

L.a. (= Lucicutia BICORNUTA) VERVOORT, 1965: 115-116

Lucicutia biuncata M. Heptner, 1971

n. sp., Heptner, 1971, p. 120-1, fig. 23

Tucicutia bradyana Cleve, ♀. Calanoida.

- Cleve, 1904, pp. 192, 204; pl. 6, figs. 33-34  
— Stebbing, 1910, p. 537.  
— CLEVE 1905: 204-206, FIGS. 33-34, N.sp.

Habitat. So. Atlantic.

*Lucicutia challengereri* Sewell

- — — Sewell 1932, p. 290, fig. 95, ♀♂, Indian O.  
includes *L. flavicornis* Brady 1883, p. 50  
— — SEWELL, 1947, P. 174, ARABIAN SEA N.  
--, VERVOORT, 1965: 112  
--, TANAKA, 1963: 26

Lucicutia cinerea M. Heptner, 1971

n. sp., Heptner, 1971, p. 122-3, fig. 24

Lucicutiidae - Calanoida.

Lucicutia clausii (Giesbrecht), ♀, ♂.  
orig. spelling 2 is

Leuckartia — Giesbrecht, 1889, p. 812

Leucicutia clausi Giesbrecht & Schmeil, 1898, p. III.

— clausii Cleve, 1904, p. 192.

— — Stebbing, 1910, p. 536.

— —, Owre, 1962

Habitat. So. Atlantic.

--, Heinrich, 1960, p. 4

--, OWREY FO40, 1964b: 366

--, OWREY FO40, 1972: TAB. 3

- — Steuer, 1910, p. 1029 (25).  
— clausi Pesta, 1921, p. 529: textfig. D 43.  
— — A. Scott, 1909, p. 126  
— — Farrar, 1929, p. 263: Sars, 1925, p. 208.  
— — Wilson, 1942, p. 192, fig. 54  
— — i, Sewell 1932, p. 289, Indian O.  
— — —, Cervigon, 1962. p. 185  
— — —, 1966, Huelsemann; p. 709;  
— — — Park, 1968, pp. 527-572.  
— — —, Shmeleva, 1963, p. 141  
— — —; Park, 1970, pp. 477, 517, figs. 211-219.  
— — — Bradford, 1970, p. 346

Lucicutia clausi (Giesbrecht, 1889)

card 2

--- ---, Hure & Scotto di Carlo, 1968, p. 58.

---, Wolfenden, p. 23, 1905

---, Vives, 1972, p. 219-220.

---, Bjornberg, 1963: 9, 53, 104 (Table 7), Tables 8 & 10, 121

---, TANAKA, 1963: 26, 38-39 ♀♂ FIG. 166, SYN: LEUCKARTIA L.

---, DEEVY, 1971: 227; FIG. 6; TAB. 3

---, BINET & DESJER, 1971: 435, 436, 455

---, BINET, ET AL., 1972: 133, 136

---, SCOTTO DI CARLO, ET AL., 1975: TAB. I

Lucicutia clausi (Giesbrecht), 1889

card 3

- , Roe, 1972a, pp. 297-
- , Shmeleva & Zaika, 1973, p. 874
- Gorice & Hart 1962b: TABLE 3, w. NORTH AMERICA
- Björnberg, 1981: 646, fig. 217 #2
- MADHU PRATAP & HARIDAS, 1986: 107, 108, 112 - INDIAN OCEAN
- Deevay & Brooks, 1977: 264
- DeDecker, 1984: 317
- Huse, 1955: 32
- Huse, 1961: 24
- Vives, 1978: 266, 280

Lucicutia clausi (Giesbrecht, 1889)

card 4

-- Mauchline 1988: 708-

-- STEPHEN 1984: 168-, ARABIAN SEA

-- Soldevilla et al 1988: 79 Canary Islands

-- CUMMINGS 1982: 33-, GULF OF MEXICO

-- HERRING 1988: 184-

-- ROE 1974: 105-, N. ATLANTIC

-- SUAREZ, E. & GASCA 1989a: 198-, YUCATAN CHANNEL

-- COPLEY ET AL. 1989: 13-, NW ATLANTIC

-- LAKKIS 1984: 290-, E. MEDITERRANEAN

Lucicutia clausi

card 5

- Wishner & Allison, 1986 : 714, fig. 9 + bl. 2
- CLAUSII, SUÁREZ, E. ET AL. 1990: 126; W. GULF OF MEXICO
- ALI-KHAN & ALI-KHAN 1982: 265, OFF PAKISTAN
- ALI-KHAN & ALI-KHAN 1982: FIGS. 1-4
- CLAUSII, SUÁREZ M., E. 1992: 142; W. GULF OF MEXICO
- CLAUSII, SUÁREZ & GASCA 1991: 596; GULF OF MEXICO
- RAZOUKS 1994: 149; PL. 72, ANTARCTIC; 10



Lucicutiidae - Calanoida.  
Lucicutia curta Farran, ♀

- — Farran, 1904, p. 603  
— — Farran, 1929, p. 264.  
— — Wilson, 1932, p. 129 : fig. 87. 1942, p. 192, f. 85  
— — JESPERSEN, 1934, BAFFIN Bay (1 SPEC.) ??  
— — Furuhashi, 1961. p. 8

Habitat. North Atlantic - Antarctic.

- — Hure & Scotto di Carlo, 1968, p. 56.

- --, 1966, Huelsemann, p. 709;  
-- -- Grainger, 1965, pp. 543-564.  
-- --; Park, 1970, p. 477  
-- --; Vervoort, 1957, p. 128-129, figs. 114-117.  
--, Heptner, 1971, p. 123, fig. 25, syn.: L. Longiserrata,  
L. ellipsoidalis
- , Roe, 1972a, pp. 297-  
--, Roe, 1972d, pp. 1023-1025.  
--, Vives, 1972, p. 220.  
--, Furuhashi, 1966: 303, 313:  
--, Bjornberg, 1963: 104 (table 7)  
--, VEROORT, 1965: 114-115  
--, TANAKA, 1963: 27, 34, 41-43  
SARS, 1925
- ♀ ♂ FIG. 168 54N: L. LONGISERRATA,

LUCICUTIA CORTA

CARD 2

- RAZOULS 1994: 151; PL. 73, ANTARCTIC; 10
- , WHEELER, 1970: 6; TAB. 4
- , FLASHAM, ET AL, 1974: 104, 109; FIG. 4; TAB. 1, 8, 9
- , SCOTTO DI CARLO, ET AL, 1975: 183, 185; FIG. 4; TAB. 1
- Devery & Brooks 1977: 264
- Vives, 1978: 266, 280
- Soldevilla, et al. 1988: 59 Canary Islands
- CUMMINGS 1982: 94; GULF OF MEXICO
- ALI-KHAN & ALI-KHAN 1982: 265-266, INDIAN OCEAN OFF PAKISTAN; FIGS 5-8

Lucicutia curvifurcata M. Heptner, 1971

n. sp., 1971, p. 127-9, fig. 26

HUCICUTIA ELLIPSODALIS

BRODSKY

— — BRODSKY, 1950, p. 333, fig. 234.

--, Heptner, 1971, p. 123, a syn. of L. curta

-- MOTODA + MINODA 1974: 232

Lucicutiidae-Calanoida.

Lucicutia glavicornis (Claus), ♀, ♂.

Leuckartia — Claus, 1863, p. 186; pl. 32, fig. 17,

Lucicutia — Giesbrecht & Schmeid., 1898, p. III.

— Cleve, 1904, p. 192; 1903, p. 364.

— Stebbing, 1910, p. 536

— Wolfenden, 1906: 1012.

Habitat, So. Atlantic: Burmese Coast:

Gulf of Mannar.

— Thompson, 1900: 281

- — Steuer, 1910, p. 1029 (25).
- — Thompson & Scott, 1903, p. 249.
- Leuckartia — Giesbrecht, 1892, p. 358: pl. 5, fig. 4:  
pl. 19, figs. 2, 3, 15-17, 21, 23, 29, 38: pl. 38, fig. 38, 40
- Lucicutia — Esterly, 1905; text-fig. 36, a-6
- Not Leuckartia — Brady, 1883, p. 50
- Lucicutia — Pesta, 1921, p. 530: textfig. D 42.
- — A. Scott, 1902, p. 405: 1909, p. 125.
- — Wolfenden, 1905, p. 1012.
- — Sewell, 1912, p. 366: 1914, p. 228. :  
reidentified as L. ovalis, 1932, p. 290  
— — Matoda and Anraku 1955 p. 212, Idzū Is.
- — , Anraku, 1954, pp. 1-8.. NW Pacific
- — , Mori, 1937, p. 72, JAPAN.
- — , Brodsky, 1950, p. 327, FIG. 226.

*Lucicutiidae - Calanoida*

*Lucicutia flavigornis* (Claus) ♀

*Leuckartia* f - Claus, 1863, p. 183: pl. 32, figs. 1-7.

— — Giesbrecht, 1892, p. 358: pls. 5, 19, 38.

*Lucicutia* f - Giesbrecht & Schmeil, 1898, p. 3

— — Cleve, 1904, p. 192 : T. Scott, 1912, p. 535.

— — A. Scott, 1902: 405

Habitat. Mediterranean : N. & S. Atlantic.

Indian Ocean.

- — A. Scott, 1909, p. 125; Sars, 1925, p. 207  
— — Wolfenden, 1911, p. 323.  
— — Farran, 1929, p. 262.  
— — Rose, 1929, p. 33  
— — Wilson 1942, p. 192 (does not mention sex)  
— — Sewell, 1932, p. 294, Indian O.  
— —, SEWELL, 1947, p. 174, Arabian Sea N.  
— —, Chiba, Tsuruta, + Maeda, 1955. Bikini.  
— —, Deerey. 1964, p. 590  
— —, OWREN FOY, 1964 b: 360, 361, 366  
— —, OWREN FOY, 1972: 501 ; TAB. 3, 6

Lucicutia flavigornis

(2)

- --, Minoda, 1972, p. 325-327.  
-- --, Cervigon, 1962, p. 186  
-- --, A.G. Fish, 1962. p. 22  
-- --, Furuhashi, 1961. p. 7  
-- --, Honjo et al, 1957. pp. 120-128  
-- --, Grice, 1962. pp. 220-222, pl. 24, fig. 1-4  
-- --, Heinrich, 1961. p. 87 (vertical distribution)  
-- --, Brodsky, 1962. p. 132, fig. 33  
-- --, Tanaka, 1960. p. 52  
-- --, Owre, 1962  
-- --, 1966, Huelsemann, p. 711; L. flavigornis  
-- -- Park, 1968, pp. 527-572.  
-- --, Heinrich, 1960, p. 4

-- --, Vilela, 1968, pp. 25-26, Est. VII fig, 3

-- --, Saraswathy, 1967, p. 80

-- --, Shmeleva, 1963, p. 141

-- --, Gilat, Kane & Martin, 1965, pp. 13-20

-- --; Park, 1970, p. 477

-- --; Bradford, 1970, p. 347

-- --, Hure & Scotto di Carlo, 1968, p. 57.

--, Chiba, 1956, p. 11

L. f. 3, Bradford, 1972, p. 46, fig. 12.4-12.6

--, Furuhashi, 1966: 300 309, 313

--, WHEELER, 1970: 5; TAB. 4

Lucicutia flavicornis Giesbrecht, 1898 card 3

--, Corral, 1970, pp. 185-6 ↙

--, Heinrich, 1960, p. 10

--, Berdugo & Kimor, 1968, p. 448

--, Roehr & Moore, 1965, p. 566-

--, Sato, 1913, no. 23, pl. 6

--, Ehrhardt, 1967, p. 740

--, Wolfenden, 1905, p. 23

--, Kolesnikov, 1966, p. 154

--, Krylov, 1967?, p. 248

--, Bowman, 1971, p. 6-

--, Vervoort, 1965: 111-113 SYNONYMS LEUCKARTIA FLAVICORNIS;  
RUDOLFIA FLAVICORNIS

- , Arcos, 1975, pp.21-22, figs. 65-67 *L. flavivornis*
- , Hue, 1967, p. 15
- , Roe, 1972a, pp. 281-
- , Roe, 1972d, pp. 1025-6, 1040, 1023
- , Vives, 1972, pp. 219-220.
- , Bowman, 1977, pp. 695-701, figs. 1-2, tables 1-2
- , Furuhashi, 1966: 299-301, 305, 307.
- , Furuhashi, 1961: 73, 107
- , Bjornberg, 1963: 9, 51, 52, 95-97, 104 (3 tables), tables 8-10  
(Table 10 misspelling Lacistema), 114, 120, 123, table 17, 125
- , TANAKA, 1963: 26, 39-41, 88 FIG. 167 SYN: LEUCKARTIA F.
- , ~~Roe~~
- , RAINBRIDGE, 1972: 69, 72
- , Mori, 1942: 554

LUCICUNA FLAVICORNIS

CARD 4

--, DEEVY, 1971: 225, 227; TAB. 6; TAB. 3

--, BINET & DESSIER, 1971: 435, 436, 455

--, BINET, ET AL, 1970: 133

--, ITOH, 1970: TAB. 1

--, FAOHAM, ET AL, 1974: TAB. 1

--, JAWED, 1973: 22, 25, 26, 31-35, 38-41, 43-46, 54, 55,  
57, 58, 60-62, 66, 77-80, 84, 85, 90, 93, 94, 96, 98-100, 102

--, SCONTO DI CARLO, ET AL, 1975: TAB. 1

--, EVANS, 1961: 196; TAB. III

--, CALEFF & GRICE (UNPUBL.) in KETCHUM & RYTHE  
1966 (UNPUBL.): APPENDIX, p. 35

--, Cervigon & Marcano 1965: 280; Fig. 14.  
--, Chiba & Hirakawa 1972: 69, 78; Tab. 2.

Lucicutia flavigornis

card 5

--, Grindley & Wooldridge, 1974, p. 204

--, Gardner & Szabo, 1982, p. 27, 358-359, fig. 22.1.2

--, Marques, 1953: 117-118

- Chen, Zhang, Chen, 19--: 65.

- Chen, Zhang, Chen, 19--: 82

- Motoda & Marumo, 1963: Pl. 1

- Marikova & Campos, 1967: 67

- Marques, 1958: 216

- Ramirez, 1969: 76-77, fig. 118, 119, 122

LUCICUTIA FLAVICORNIS (CLAUS)

CARD 6

--, PETERSON + MILLER, 1977: 718

--, VAISSIÈRE + SEGUNN, 1984: 7, TABS. 1-3

-- MASSUTI + NAVARRO 1950: 20.

-- TRANTER 1977: 583-625.

-- SANDER + MOORE 1978: 233.

-- YOUNG-BLUTH 1982: 120, BAHAMAS

-- YOUNG-BLUTH 1979: 22, CARIBBEAN

-- MICHEL + FOYO 1976: 90, 131, 136, 137, 140-141, 159

(FIG. 17A), 191-197 (TAB. 39)

- Björnberg, 1981: 648, fig. 217<sup>#1</sup>

Lucicutia flavigornis (Claus)

card 8

-- Carli 1971b: 372-3, Ligurian Sea

-- CARLI + CRISAFI 1983: 56, PL. 8 FIG. 1, ITALY

-- MAUCHLINE 1988: 708-

-- KIORTSIS ET AL. 1969: 37-, AEGEAN

-- STEPHEN 1989: 165-, ARABIAN SEA

-- CAMPANER 1986: 262-, S. BRAZIL SHELF

-- SCHULZ 1986b: 459- 465, JARGASSO SEA

-- Soldevilla, et al. 1988: 59 Canary Islands

-- DESSIER + DENIGUY 1987: 14, 397-, EASTERN  
EQUATORIAL PACIFIC

Lucicutia flavigornis (Claus)

card 7

- Goswami 1983: 252, India
- GRICE & HART 1962a: 297, GULF STREAM
- GRICE & HART 1962b: TABLE 3
- Dakin & Coletaux, 1940: 92, fig. 130a-b.
- Tanaka, 1965a: 27, fig. 7a-c
- Binet & Dessier, 1968: 9, OFF MADAGASCAR
- MADHUPRATAP & HARIDAS, 1986: 108, 107, 110-112,  
INDIAN OCEAN
- Dawson & Knatz, 1980: 71, pl. 58, fig. a-g.
- Kimmerer, et al., 1985: table 4 Australia

- Deevay & Brooks, 1977: 264
- DeDecker, 1984: 317
- MICHEL ET AL. 1986: 8, 20-21, 35 - ARABIAN GULF
- Hure, 1955: 31-32
- Hure, 1961: 24
- Heberer, 1932: 265, passim, fig. 86, 87
- Moryakova & Kampos, 1966: 131, passim Cuba
- Vives, 1978: 266, 280
- REGNER 1987: 373-, ADRIATIC
- CARLI, 1968: 100-, LIGURIAN SEA
- CARLI 1971b: 372-, "

Lucicutia flavigornis (Claus)

card 9

- Rodriguez & Vives 1984: 242-, Malaga Harbor
- Cummins 1982: 31-, GULF OF MEXICO
- Herring 1988: 183-
- Roe 1974: 104-, N. ATLANTIC
- Jimenez-Perez & Lara-Lara 1988: 126-,  
GULF OF CALIFORNIA
- Suarez, F. & Gasca 1989a: 198-, YUCATAN CHANNEL
- Valentim: 1989: 30-, CABO FRIO, BRAZIL
- Suarez M. & Gasca 1989c: 94-, COSTA RICA DOME

- Lucicutia flavigornis (Claus) card 10
- Suarez, E. & Gasca 1990:62, Quintana Roo, Mexico
- LAKKIS 1984: 290-, E. MEDITERRANEAN
- LICIAUTIA -, SHNELEVA + KOVALEV 1968: 66, FIG 10b
- ALVARINO 1989: 69-, OFF S. CALIFORNIA
- Wishner et Allison, 1986: 714-715, fig. 5, pl. 2
- SUÁREZ, E. ET AL. 1990: 126-, W. GULF OF MEXICO
- REID 1990g: 188
- ROBICHAX ET AL. 1981: 82-, SAN SALVADOR, BAHAMAS
- BOWNAN 1978a: 114-

Lucicutia flavicornis (Claus)

card 11

-- Ali-Khan & Ali-Khan 1982: 266-267, figs. 9-13, Indian  
Ocean off Pakistan

-- SUÁREZ H., F. 1992: 142-, W. GULF OF MEXICO

-- SUÁREZ Y GASCA 1991: 596-, GULF OF MEXICO

-- RAZOUKS 1994: 10

-- YAMASHITA Y MOREIRA 1981: 41

-- OKEMWA & REVIS 1986: 31, TAB. I, OFF KENYA

-- CARMAN & DOBBS 1997: 121, TAB. I

-- ROSE 1927: 145, 153, 154, ALGERIA

LUCICUTIA FLAVICORNIS (CLAUS)  
-- ROSE 19~~2~~<sup>3</sup>: 54-81, ALGERIA

CARD 12

*Lucicutia formosa*, 1966, Huelsemann, p. 713;

--, GRCE, 1969: 454

11. 2 450 1

*Lucicutia frigida* Wolfenden, ♀.


— — Wolfenden, 1911, p. 320: Text-fig. 62, a-b.  
— — ; Vervoort, 1957, p. 126-128, figs. 111-114.

— , Vervoort, 1965: 114

— MOTODA + MINODA 1974: 232

— HULSEMANN 1989: 97

Habitat. Antarctic.



Lucicutia gaussae Grice, 1963 ♂

-- --, Heptner 1965 p. 1175

L. ovalis Wolfenden, 1906

L. gaussae; Park, 1970, p. 477.

--, Corral, 1970, pp. 187-8, pl. 49

--, Bowman, 1971, p. 6-

--, Roe, 1972a, pp. 297-

--, Roe, 1972d, pp. 1026, 1040.

--, Vives, 1972, p. 220.

-- Björnberg, 1981: 648, fig. 217 #5

--, Wheeler, 1970: 4

--, Bowman, 1977, pp. 695-701, figs. 1-2, tables 1-2

--, Deevey, 1971: TAB. 3

--, Binet & Desseix, 1971: 436, 455

--, Calef & Grice (unpubl.) in Ketchum & Ryther  
1966 (unpubl.): TAB. 2 ; APPENDIX

—Deevey & Brooks, 1977: 264

—DeJacker, 1984: 317

- Soldevilla, et al. 1988: 59 Canary Islands  
-- HULSENANN 1989: 98  
-- GAUSSAE, COMUMINOS 1982: 32, GULF OF MEXICO  
-- ALI-KHAN & ALI-KHAN 1982: 267-268, FIGS.  
14-18, INDIAN OCEAN OFF PAKISTAN  
-- RAZOUKS 1994: 155  
-- CARMAN & DOBBINS 1997: 121, TAB. I

= L. GAUSSAE GRICE Lucicutiidae - Calanoida

Lucicutia ovalis Wolfenden, ♀.

(3)

n. sp., Wolfenden, 1906, p. 28-29, pl. 9

— — Wolfenden, 1911, p. 319: pl. 35, fig. 6:

Text-fig. 61, a-c. ♀ only

— — Harran, 1929, p. 263: Text-fig. 25, (descr. of ♂)

— — Sewell 1932, p. 290, Indian O. (includes, in  
part, *L. flavicornis*, Sewell 1912, p. 366)

Habitat. Atlantic, Indian Ocean

— — Rose, 1933, p. 197, fig. 233 (after Wolfenden)

— — Wilson, 1950, p. 256

- — Matoda and Anraku 1955. p. 212, Idzu Is.
- —, Mori, 1937, P. 72, JAPAN.
- —, Chiba, Tsuruta, Maeda, 1955. Bikini.
- —, Furuhashi, 1961. p. 7
- —, Honjo et al, 1957. pp. 120-128
- —, Grice, 1962. p. 222, pl. 24, fig. 5-9
- —, Bowman, 1977, p. 700
- —, Hure & Scotto di Carlo, 1968, p. 57.
- , Corral, 1970, p. 187
- . Berdugo & Kimor, 1968, p. 448

= L. GAUSSAE GRICE

Lucicutia ovalis Wolfenden, 1906

card 4

-- Kiortsis et al. 1969: 37-, Aegean

-- Soldevilla, et al 1988: 59 Canary Islands

renamed L. gaussae Grice 1963: 498.

— Wishner & Allison, 1986: 715, Fig. 5, Tab. 2

-- Razouk 1994: 155

*lucicutiidae*-Calanoida

*Lucicutia gemina* Farran, ♀♂.

— — Farran, 1926, p. 275: pl. 9, figs. 4-8.

— — Farran, 1929, p. 263

— —, 1966, Huelsemann, p. 715;

— —, Shmeleva, 1963, p. 141

— —; Park, 1970, p. 477

Habitat. Bay of Biscay, 0-1250 fath.

New Zealand

— ; WHEELER, 1970: 4

— , SCOTTO DI CARLO, ET AL., 1975: TAB. I

(see back)

- ---, Hure & Scotto di Carlo, 1968, p. 58.  
---, Roe, 1972d, p. 1025.  
---, VERVOORT, 1965 : 113  
---, TANAKA, 1963 : 41  
---, BINET & DESSIER, 1971 : 435-436, 455  
---, BINET, ET AL., 1972 : 133  
---, CALEF & GREE (UNPUBL.) IN KETCHUM & RYTHM,  
1966 (UNPUBL.) : TAB. 2; APPENDIX  
---, ARTIONKIN ET AL. 1966: 6.  
— Bjornberg, 1981 : 648, fig. 217 #6  
— Deevney & Brooks, 1977 : 264  
— Deevney & Brooks, 1980 : 327

Lucicutia gemina Farran, 1926

card 2

- Schulz 1986b: 463, 464, Sargasso Sea
- CAMMINOS 1982: 33; GULF OF MEXICO
- HERRING 1988: 184-
- ROE 1974: 103-, N. ATLANTIC
- SUAREZ M. & GASCA 1989c: 94-, COSTA RICA DOME  
— Wishner & Allison, 1986: fig. 2
- ARTEMKIN ET AL. 1966(1973): 179-, CARIBBEAN

Lucicutia gigantissima M. Heptner, 1971

n. sp., Heptner, 1971, p. 129-30, fig 27

Lucicutiidae - Calanoida

Lucicutia gracilis G.O. Sars. ♀

— — Sars, 1905, p. 8.

--, (= L. MAGNA), TANAKA, 1963: 36-38

Habitat. Station 1834, Monaco.

Calanidae - Calanoida

Lucicutia grandis (Giesbrecht), ♂. ♀

Leuckartia — Giesbrecht, 1895, p. 258: pl. 4, f. 4.

Lucicutia — Giesbrecht & Schmeil, 1898, p. 111

— (♀) Wolfenden, 1904, p. 121: pl. 9, f. 37-38.

— — Wolfenden, 1911, p. 315: text-fig. 58, a-b.

Habitat. Pacific Ocean. North Atlantic  
Antarctic.

- — Parran, 1929, p. 264.
- — Sars, 1925, p. 208; pl. 56, f. 1-14
- — Wilson, 1932, p. 128; fig. 86. 1942, p. 192, fig. 87
- —, Brodsky, 1950, p. 328, fig. 227.
- —, Jespersen, 1934, p. 104, W. GREENLAND WATERS (DEEP).
- —, Jespersen, 1940, p. 51, Iceland.
- — —, Clarke, et. al., 1962. pp. 541-564
- — —, 1966, Huelsemann, p. 717.
- — —; Vervoort, 1957, ~~p.~~ p. 131.
- part = L. wolfendeni; Vervoort, 1957, p. 130-131, <sup>figs. 120-121.</sup>
- , Stormer, 1929, p. 48
- , Heptner, 1971, p. 146, a syn. of L. wolfendeni

Lucicutia grandis

card 2

--, Wolfenden, 1905-1906, p. 8, pl. 2, p. 22-23

--, TANAKA, 1963: 57

-- MOTODA + MINODA 1974: 232 (as L. GRADIS)

-- MAUCHLINE 1988: 708-

-- MARUKAWA 1921: 8-, JAPAN

-- HERRING 1988: 184-

-- ALI-KHAN + ALI-KHAN 1982: 268-269,  
FIGS. 19-22, INDIAN OCEAN OFF PAKISTAN

-- RAZOUKS 1999: 157; 10

A ♀ 6.5 mm. long - v largest known species of  
Tricicotia from v deep waters of Atlantic: The  
anterior antennae about 4 joints longer than  
v whole body: v gen. seg't. asymmetrical: v 1<sup>o</sup> pair  
of feet with tubular process on v 1<sup>o</sup> basal; all  
swimming feet, including v 5<sup>o</sup> pair, with 3-  
jointed ex. 4 endopodites (pl. IX. fig. 37-38) Very  
pigmented, with deep orange pigment about 6  
in bulk, and all organs smooth and i feet color  
ed a shade of burnt sienna. This may be v  
♀ species described by Giesbrecht from Pacific  
T. grandis, of which, however, he only knew v 8-

Published August 16, 1923.

CHALMERS, LORD

Thomas William Rhys Davids, 1843-1922: Memoir by Lord  
CHALMERS. 1923. Royal 8vo (9 $\frac{3}{4}$ x6 $\frac{1}{4}$ ), pp. 6. Paper 35c.

*From the Proceedings of The British Academy. Vol. X.*

A note on the life-work of this great student of Buddhism and  
Founder of the PALI TEXT Society.

OXFORD UNIVERSITY PRESS AMERICAN BRANCH, NEW YORK

Iucicutiidae - Calanoida.

Lucicutia intermedia G.O.Sars. ♀

— — Sars, 1905b, p. 8

— — Sars, 1925, p. 215; pl. 58, f. 4-8.

Temperate Atlantic.

— —, 1966, Huelsemann, p. 721;

— —; Park, 1970, p. 477

Habitat. Station 1768, 1851, Monaco.

- Deevey & Brooks, 1977: 264

--, Heptner, 1971, p. 131, fig. 28, syn.: L:tenuicauda,  
L. phlyra

--, TANAKA, 1963: 32

--, WHEELER, 1970: TAB. 4

?- MAUCHLINE 1988: 708, AS Lueicutta sp.

Calanidae - Calanoida.

Lucicutia longicornis (Giesbrecht), ♂. ♀

Leuckartia - Giesbrecht, 1889, p. 812. Tg. 39

— — Giesbrecht, 1892, p. 359: pl. 19, f. 7, 30: pl. 38,

Lucicutia - Giesb. & Schmeil, 1898, p. 111

— — Wolfenden, 1911, p. 323.

— — Wilson, 1942, p. 192

Habitat. Pacific Ocean: Tropical  
Atlantic.

-- JESPERSEN, 1934, P. 104.

W. GREENLAND WATERS.

= Matikova & Camara 1967: 17

— —, 1966, Huelsemann, p. 721;

—, Roe, 1972a, pp. 297-

— Bjornberg, 1981:648, fig. 217#7

— Deevey & Brooks, 1977:264

— Moryakova & Kampos, 1966: 131, passim

Cuba

Lucicutia longicornis (Giesbercht, 1889) card 2

-- Soldevilla, et al. 1988: 59 Canary Islands

-- ALI-KHAN & ALI-KHAN 1982: 268, 269,  
FIGS. 23-26, INDIAN OCEAN OFF PAKISTAN

HUCICUTIA HONGIFURCA BRODSKY

— — 1950, p. 336, fig. 233.

— — , 1966, Huelsemann, p. 723;

— , Heptner, 1971, p. 132, fig. 29

Lucicutia longiserrata (Giesbercht 1892) card 3

-- Soldevilla et al. 1988: 59 Canary Islands

LUCIDUNIA LONGISERRATA (GIESBRECHT) CARD 3

-- CARLI + SERTORIO 1964: 207-, LIGURIAN SEA

--, TANAKA, 1963: 26, 27, 32-34 ♀♂, FIG. 163, SYN:

LEUCKARNA L.

--, (SARS, 1905; FARRAN, 1926, 1929; WILSON, 1932, 1950;

JESPERSON, 1934 = L. CURTA), TANAKA, 1963: 41-43

--, WHEELER, 1970: TAB. 4

--, FASHAM, ET AL, 1974: TAB. 1

--, SCOTTO DI CARLO, ET AL, 1975: 183, 184, 185; FIG. 3, 4;

TAB. 1

—Deevey & Brooks, 1977: 264

Calanidae - Calanoida  
Lucicutia longiserrata (Giesbrecht), ♀.

Leuckartia - Giesbrecht, 1889, p. 813. fig. 36

— — Giesbrecht, 1892, p. 359: pl. 19, f. 18, 25: pl. 38,

Lucicutia - Giesbrecht & Schmeil, 1898, p. 112

— — Thompson, 1903, p. 25.

Habitat. Pacific Ocean: Atlantic

- — Farran, 1908, p. 64: pl. 6, figs. 21-22.  
— — v. Breemen, 1908, p. 116, text-fig. 139.  
— — A. Scott, 1909, p. 127.  
— — Wolfenden, 1911, p. 323.  
— — Sars, 1925, p. 220: pl. 59, f. 12-15.  
— — JESPERSEN, 1934, P. 105, W. GREENLAND.-DAVIS STR.  
— —, 1966, Huelsemann, p. 723;  
— —; Park, , 1970, pp 477.  
— —, Heptner, 1971, p. 123, a syn. of L. curta  
— —, Roe, 1972a, pp. 297-  
— —, Roe, 1972d, pp. 102~~3~~-1025.  
— —, Furuhashi, 1966: 309

LUCICUNA

LONGISPINA

TANAKA, 1963

-- TANAKA, 1963: 30 - 32 ♀ FIG. 162

--, VIVES, 1972: 220

Lucicutiidae - Calanoida

- Lucicutia lucida Farran, ♀♂. 120  
— — Farran, 1909 (1906), p. 62: pl. 3, f. 22: pl. 4, f. 16  
— pera, A. Scott, 1909, p. 129: pl. 40, f. 1-9  
— lucida Sars, 1925, p. 219: pl. 59, f. 6-11 = <sup>L. PERA</sup>  
<sub>(TANAKA,  
1963)</sub>  
— —, Jespersen, 1940, p. 52. SE Iceland.  
— —, 1966, Huelsemann, p. 725;

Habitat. Atlantic: Malay Archipelago:  
Mediterranean.

- —, Hure & Scotto di Carlo, 1968, p. 57.  
—, Roe, 1972a, pp. 297-  
—, Roe, 1972d, pp. 1022-1024.

--, Vives, 1972, pp. 219-220.

--, TANAKA, 1963: 27, 43-45 8, 46

--, BINET & DESSIER, 1971: 436

--, FASHAM, ET AL., 1974: TAB. I

--, SCOTTO DI CARLO, ET AL., 1975: TAB. I

-- MASSUTÍ & NAVARRO 1950: 20.

—Deevey & Brooks, 1977: 264

Tucutiidae - Calanocera

Tucutia macrocera G.O.Sars. ♀

— Sars, 1920, p. 10.

— Sars, 1925, p. 213: pl. 57, f. 12-15.

— JESPERSEN, 1934, p. 105, W. GREENLAND WATERS.

— " , 1940, p. 52. S. of Iceland.

Habitat. Temperate Atlantic

— Deevney & Brooks, 1977 : 264

— TANAKA, 1963 : 26, 32, 34-36 ♀, FIG. 164

— RAZOUKS 1994: 10

— —, 1966, Huelsemahn, p. 725;  
— —; Vervoort, 1957, p. 130, figs. 118-119.

—, Bradford, 1971, p. 24, fig. 89

—, Heptner, 1971, p. 134, fig. 30, 31

— Razouks 1994: 153-, Pl. 74, ANTARCTIC

Calanidae - Calanoidea.  
Lucicutia magna Wolfenden, ♂.

- — Wolfenden, 1904, p. 121: pl. 9, f. 35, 35a.
- — Wolfenden, 1911, p. 316: text-fig. 59, a-b.
- — Farran, 1929, p. 264.
- — Wilson, 1932, p. 130 : fig. 88.

Habitat. Faroe Channel: Atlantic:  
Antarctic:

--; OWEN FORO, 1964a: 343

- — Fowler (Wolfenden), 1903, p. 124.
- — , Anraku, 1954, pp. 123-136.. Aleutian Is.
- — , SEWELL, 1947, P. 174, Maldives area,
- — , 1966, Huelsmann, p. 727;
- — ; Park, 1970, p. 477
- — . Bradford, 1970 p. 347
- , Roe, 1972a, pp. 297-
- , Roe, 1972d, pp. 1022-1023.
- , Vives, 1972, p. 220.
- , Furuhashi, 1966: ~~310~~ 309
- , TANAKA, 1963: 26, 36-38 ♀♂, 43, FIG. 165. SYN: L. GRACILIS; L. ATLANTICA
- , WHEELER, 1970: TAB. 4

LUCICUTIA MAGNA

CARD 2

- OWRE & FOYO, 1972: 74B.3
- SANDER & MOORE 1978: 233.
- Deeney & Brooks, 1977: 264
- HERRING 1988: 184-
- SUÁREZ, E. ET AL. 1990: 126-, W. GULF OF MEXICO
- SUÁREZ H., E. 1992: 143-, W. GULF OF MEXICO
- SUÁREZ & GASCA 1991: 596-, GULF OF MEXICO .
- PAZOUIS 1994: 10

*Lucicutiidae-Catarrhini*

*Lucicutia major* Wolfenden, ♀

— Wolfenden, 1911, p. 317.

— —, Grice & Hulsemann, 1967, p. 29, figs. 144-147;

Habitat. Atlantic.

Tucutidae - Calanoida

*Lucicutia maxima* Steuer, J.R.

- — Steuer, 1904, p. 596 : text-fig. 4. (♀)  
— — A. Scott, 1909, p. 127 : pl. 41, figs. 1-10  
— — Wolfenden, 1911, p. 318 : text-fig. 60, a-b.  
— — Sars, 1925, p. 211 : pl. 57, f. 1-6.  
— — Sewell 1932, p. 295, fig. 94, 96, develop. stages

Habitat. Malaysia : Atlantic.

— — Tanaka, and Mori, 1967, pp. 239-260.

--, Grice, 1969 : 454

--, Owre & Foy, 1972 : 493 ; TAB. 3

-- --, 1966, Hulsemann, p. 729;

--, Wolfenden, 1905, p. 22

--, TANAKA, 1963: 26, 27, 49-51 ♂, FIG. 172

--, ~~Wolfenden, 1906.~~

-- Grice & Hulsemann, 1968: Hb. 2

- Binet & Dessier, 1968: 7, OFF MADAGASCAR

-- MAUCHLINE 1988: 708-

lucicutia

OBHONGA

BRODSKY

— — 1950, p. 339, fig. 236.

— —, 1966, Huelsemann, p. 731;

— Minoda, 1972, p. 327-330.

HUCICUTIA ORIENTALIS Brodsky

— — 1950, p. 337, fig. 235.

— — , 1966, Huelsemann, p. 732;

— , Heptner, 1971, p. 137, fig. 32

hucicutia ovaliformis

Brodsky

-- 1950, p. 334, fig. 232.

--, Vinogradov, 1960, p. 502

--, Furuhashi, 1966: 298 (L. ovaliformis of Tanaka, 1963  
x~~xxxxxx~~ = Isochaeta ovalis )

--, VERVOORT, 1965: 114

-- MOTODA + MINODA 1974: 232

-- Gordon, et al, 1985: 91 , table 2

-- HULSEMANN 1989: 97

= Lucicutia

Centropagidae - Calanoida.

Isochaeta ovalis Giesbrecht, ♀.

— Giesbrecht, 1889, c, p. 812.

— Giesbrecht, 1892, p. 367: pl. 29, 38.

— Giesbrecht & Schneid, 1898, p. 112.

—, Brodsky, 1950, P. 341, Fig. 238.

Lucicutia ovalis, Grice, 1963. pp. 497-498

Habitat. Tropical Pacific.

*T. ovalis*

-- --, Furuhashi, 1961. p. 8

--, Furuhashi, 1966: 297, 298  
Tanaka, 1963 = I. ovalis (Lucicutia ovaliformis of  
309, 315), 299, 300, 303, 305, 307,

--, TANAKA, 1963: 53-55 ♀ ♂, FIG. 174

-- HULSEMANN 1989: 97

Lucicutia ovalis

card 2

--, Shmeleva & Zaika, 1973, p. 874

--, Gardner & Szabo, 1982, p. 27, 360-361, fig. 22.1.3  
Syn.: L. frigida, L. ovaliformis

-- GRICE & HART 1962b: TABLE 3, w. NORTH ATLANTIC

— Tanaka, 1965a: 29, fig. 8a-c

— Bjornberg, 1981: 648, fig. 217 #4

-- MADHUPRATAP & HARIDAS, 1986: 108, 112 - INDIAN OCEAN

— Deesey & Brooks, 1977: 264

— Hure, 1955: 32

Lucicutia ovalis (Giesbrecht) 1889.

-- --, Grice, 1963. pp. 497-498

Isochaeta ovalis Giesbrecht, 1889.

L. ovalis, Heinrich, 1961. p. 87 (vertical distribution)

-- --, 1966, Huelsemann, p. 732;

-- --, Vilela, 1968, p. 26 Est. VIII Fig. L

-- --, Heptner 1965, pp 1165-75.

Lucicutia ovaliformis Brodsky, 1950

L. frigida Wolfenden, 1911

-- --: Vaupel-Klein, 1970, p. 37.

L. ovalis; Park, 1970, p. 477

-- --, Heinrich, 1960, p. 4

- , Chiba, 1956, p. 14
- , Bradford, 1971, p. 24, fig. 90-92
- , Heinrich, 1960, p. 3
- , Heptner, 1971, p. 140, ~~figx~~
- , Roe, 1972a, pp. 297-
- , Furuhashi, 1966: 309
- , Furuhashi, 1961: 107
- , Bjornberg, 1963: 53, 9, 104 (3 tables), tables 8ff, 10, 12, 123
- , Vervoort, 1965: 114
- , TANAKA, 1963: 26, 27-28 & 6<sup>7</sup>
- , WHEELER, 1970: TAB. 4

Lucicutia ovalis (Giesbrecht, 1889)

card 4

-- Herring 1988: 184-

-- COPLEY ET AL. 1989: 13; NW ATLANTIC

-- SUÁREZ, E. ET AL. 1990: 126; W. GULF OF MEXICO

-- REID 1990g: 188

-- SUÁREZ, E. 1992: 143; W. GULF OF MEXICO

-- SUÁREZ & GASCA 1991: 596; GULF OF MEXICO

-- RAZOULS 1994: 155; PL. 75; ANTARCTIC; 10

ISOCHARTA - , ROSE 1934: 59-81, ALGERIA

L., ROSE 1934: 54-81, ALGERIA

Lucicutia ovalis Wolfenden - see L. gaussae Grice

*Lucicutia pacifica*, 1950, Brodsky, p. 330, fig. 228;

— —, 1966, Huelsemann, p. 733;

—, TANAKA, 1963: 32

— MOTODA + MINODA 1974: 232.

Lucicutia pallida, 1966, Huelsemann, p. 733;

—Deevey & Brooks, 1977: 264

Lucicutia paraclausi Park, 1970

— — Park, 1970, pp. 477, 517-519, figs. 220-225.

Lucicutia parva Grice & Hulsemann

— — Grice & Hulsemann, 1965, pg. 244, fig. 16 (1) to  
      (s)  
— — ; Park, 1970, p. 477

*Lucicutia pellucida*, 1966, Huelsemann, p. 734;

Lucicutiidae - Calanocidae

Lucicutia peregrina A. Scott, ♀, juv. "

— A. Scott, 1909, p. 129: pl. II, figs. 1-9

— —, 1966, Hußmann, p. 735;

—, Roe, 1972a, pp. 305-

—, TANAKA, 1963: 26, 27, 44, 45-46 ♂, FIG. 160, ♂YR. Lucinda A,  
JARS, 1925

Habitat. Malaysia.

— Deevy & Brooks, 1977: 264

Lucicutia pera A. Scott 1909

card 2

-- Soldevilla, et al. 1988: 59 Canary Islands

*Lucicutiidae - Calanocera*

*Lucicutia philyra* A. Scott, ♀.

— A. Scott, 1909, p. 128: pl. 40, f. 10-18.

L. philyra Scott, 1909, Heptner, 1971, a syn. of  
L. intermedia, p. 131

--, TANAKA, 1963 : 26

Habitat. Malaysia.

hucicutia Polaris Brodsky, 1950

— — 1950, p. 330, fig. 229.

— —, Johnson, 1963. pp. 92, 93, 97

— —, 1966, Huelsemann, p. 735;

—, Heptner, 1971, p. 141, fig. 22 (9-13)

—, Vidal, 1971, pp. 23-, 26, 27-, figs. 3 & 4.

—, TANAKA, 1963: 32

— MOTODA + MINODA 1974: 232

— ALI-KHAN + ALI-KHAN 1982: 269-270, FIGS.

27-29, INDIAN OCEAN OFF PAKISTAN

HUCICUTIA PROFUNDA BRODSKY

— — 1950, p. 336, fig. 234.

— —, 1966, Huelsemann, p. 735;

— , Heptner, 1971, p. 142, fig. 33

Lucicutia pseudopolaris Heptner, 1969

L. polaris Brodsky, 1950 ( $\sigma$  only); Hulsemann, 1966  
( $\sigma$  only).

L. pseudopolaris, Vidal, 1971, pp. 23-25, 26-,  
figs. 2 & 4.

*Lucicutia rara*, 1966, Huelsemann, p. 735;

*Lucicutia sarsi*, 1966, Huelsemann, p. 736;

-- --, Grise & Hulsemann, 1968: fig. 2.

-- HERRING 1988: 184-

Lucicutia scopularis Brady

--, Wolfenden, 1905, p. 23, a syn. of Heterorhabdus

LUCICUTA SEWELLI

TANAKA, 1963

--, TANAKA, 1963: 28-30 ♂ FIG. 161

--, PARK, 1970: 477

— Deevy & Brooks, 1977: 264

Lucicutiidae — Calanoida

Lucicutia simulans G.O.Sars, ♀.♂

— — Sars, 1920, p. 11

— — Sars, 1925, p. 216: pl. 58, f. 9-13.

— , Stormer, 1929, p. 48

-- MAUCHLINE 1988: 708

Habitat. Mediterranean.

*Lucicutiidae* - Calanoida

*Lucicutia tenuicavda* G.O.Sars, ♀♂

— — Sars, 1907, p. 18.

— — Sars, 1925, p. 212: pl. 57, f. 7-11.

— —, Huelsemann, 1966, p. 737;

— —, Soldevilla, et al. 1988: 59 Canary Islands

Habitat. Temperate Atlantic.

--, Heptner, 1971, p. 131, L. tenuicauda Sars: Grice &  
Hulsemann, 1965 a syn. of L. intermedia

*Lucicutia ushakovi* Brodsky

1955---pp. 22-23, fig. 12. (p. 204)

--- ---, 1966, Huelsemann, p. 737;

---, Heptner, 1971, p. 144, fig. 34

①

Lucicutia wolfendeni Sewell, 1932

-- --, Furuhashi, 1961. pl 7

-- --, 1966, Huelsemann, p. 738;

-- --; Vervoort, 1957, p. 130-131, figs. 120-121.

--, Heptner, 1971, p. 146, fig. 30 (7), syn.: L. grandis  
--, Harding, 1974, p. 148.

--, TANAKA, 1963: 46-49 ♀♂ FIG. 171, SVU: L. GRANDIS

--, Grice & Hulsemann, 1968: tabl. 2

-- Bjornberg, 1981: 648, fig. 217 #3

-- Deesey & Brooks, 1977: 264

-- MAUCHLINE 1988: 708 -

-- HERRING 1988: 184 -

Lucicutia wolfendeni Sewell, 1932

card 2

-- Razouls 1994: 157-, Pl. 76, Antarctic; 10

Lucicutia species

--, Stormer, 1929, p. 48

--, Meshcheryakova, 1970, p. 112

--, Bjornberg, 1963: 104 (table 7)

-- Kimmuner et al., 1985: table 4      Australia

-- CAMPANERI 1986: 262-, S. BRAZIL SHELF

LUCICUTIA SP.

--, ANDRONOV & MAIGRET 1980: 71.

--, DAVIS 1950: Tab. V-p. 93.

-- TRANTER 1977: 598.

-- PATRITI 1984a: 157-161, Gulf of FOS-SUR-MER

-- MAUCHLINE 1988: 708, ? ≤. INTERMEDIA

LUCICUTIA sp. 1

-- COPLEY & WIEBE 1990: 6, FIG. 18c-e, GUAYMAS  
BASIN

LUCICUTIA spp.

--, CARRINOS 1982: 33-, GULF OF MEXICO  
spp. 1-3, COPLEY + WIEBE 1990: 17, GUAYMAS BASIN  
— BINET + DESSIER 1968: 9, OFF MADAGASCAR