

in)

Caligidae - Caligoida.

Lütkenia.

Claus, 1864.

L. astrodermi ^(M).

Syn. Lütkenia Claus, 1864, p. 365.

Lütkenia Brian, 1906, p. 48.

Lütkenia T. & A. Scott, 1913, p. 75.

Cecropsina Heller, 1865, p. 209.

not Lütkenia - Duncan - Ophiuroidea - 1878.

To Prof. Christian Lütken.

not Luetkenia Steind - Pisces, 1876.

Cephalothorax divided, 2nd. & 3rd. thorax segts. fused. Two small dorsal plates on 4th. segt. Gen. segt. wide, deeply incised posteriorly: abdomen not segmented, concealed. Anal laminae small with short setae. First legs with 2-jointed rami; endopod small, without setae, exopod elongate. 2nd. & 3rd. legs with 2-jointed rami, set with plumose setae. Fourth pair with foliaceous rami, unsegmented. Egg strings long.

Cecropidae - Siphonostomatoida

Luetkenia Claus, 1864

- Kabata 1992: 38, diagnosis

Cecropidae: Siphonostomidae ^{Caligidae.}

Luettkenia asterodermi, Claus, ♂♀.

— —. Claus, 1864, p. 369; pl. XXXIV, figs. 10-15.

— —, Claus, 1885, p. 362.

asterodermi, B-Smilk. 1899, p. 458.

-- CASTRO ROMERO + BAEZA KUROKI 1983: 3-8, FIGS. 1-13 ♂♂

Host. Gills of Asterodermus. sp. (Coryphaenoides)

-- KABATA 1992: 38, FIG. 7A-C, SYNONYMY, HOST LIST
Mediterranean.

Luetkenia elongata Shiino, 1963

LÜTKENIA -, SHIINO 1963: 349-356, FIGS. 1-2, N.SP.

♀♂, OFF LOWER CALIFORNIA, HOST LUVARUS IMPERIALIS

-- Castro Romero & Baeza Kuroki 1983: 3, 8.

-- KABATA 1992: 38, 40

Luetkenia glabra, Heller, ♂-♀.

Cecropsina glabra, 1865, p. 209, (Heller); pl. XIX. figs. 1+2.

Luetkenia —, Carus, 1885, p. 362.

— —, Brian, 1898, p. 13.

— —, B-Smitk, 1899, p. 458.

Host. Gills of Luvarus imperialis -

(over)
Mediterranean.

Cecropsina g - , Heller, 1866, p. 32.

Lütkenia flava, Richiardi, 1880, p. 3.

— —, Valle, 1880, p. 59.

— —, Brian, 1903, p. 2.

— —. Stossich, 1880, p. 256. On the gills
of Ansonia Cuvieri Ris.

— OF VALLE (1880), CASTRO ROMERO & BAEZA KUROKI
1983: 4, A SYNONYM OF LUETKENIA ASTERODERMI
CLAUS.

Luethkenia integra, Richiardi, ♀.

- —, Richiardi, 1880, p. 148 (no description).
 — —, Carus, 1885, p. 362.
 — —, Brian, 1898, p. 13; pl. II, fig. 7.

Host. Gills of Galeus canis; Mustelus laevis,
 and M. equestris, also in the mouth.

Mediterranean.