

in/ Canthocamptidae - Harpacticoida.

Morariopsis valid

Borutsky, 1931

M. typica^①

Syn. Morariopsis Borutsky, 1931, p. 271.

- BORUTSKY 1931: 102
- BORUTSKY 1960: 147, KEY TO SOME SPP.
- BORUTSKII 1966: 771, cf. EPACTOPHANOIDES, n. gen.

Endopods of 2nd, 3rd, & 4th legs 1-segd. in ♀:
first legs natatory, with setae, without claws:
endopods of 1st legs 2-segd, shorter than the
3-segd exopods: end segt of 5th leg with 3 or
4 setae, basal expansion with 6 unequal setae.
Lake Baikal.

MORARIOPSIS

- LEWIS, M. 1986: 115 -
- BRANCELT 1992 (1991); 199-200, CF. PARAMORARIOPSIS

Morariopsis baicalensis Borutzky & Okuneva

--, Borutzky & Okuneva, 1975: 64-66
fig. 1-16.

MORARIOPSIS INSOLETUS - SEE

CANTHOCAMPTUS

Morariopsis kieferi Petkovski, 1959

—
— Kiefer, 1978: 219
-- Rouch 1986b: 340

Canthocamptidae - Harpacticoida

Morariopsis latifurcata Borutzky ♀

— — Borutzky, 1931, p. 196 : figs. 81, 82.

— — Borutzky, 1952, p. 364, F. 100, 1-12

— —, LANG, 1948. P. 1186.

--, Sterba, 1967, p. 288, 333

--, Jakobi, 1959, p. 277

Habitat. Russia

--, BORUTZKY 1960: 147, FIG. 63(7)

= MORARIOPSIS

Canthocamptidae - Harpacticoida

Moraria scotenophila Kiefer ♀ ①

— — Kiefer, 1930, p. 226: figs. 11 — 14

— —, Borutzky, 1952, p. 366, F. USSR.

--, Brian, 1955, p. 21

--, Klie, 1943, p. 189

Habitat. Karst; St. Canzian

Morariopsis scotenophila (KIEFER, 1930)

CARD 2

--, Jakobi, 1959, p. 277

—Kiefer, 1978:219

-- ROACH 1986b: 340

-- BRANCELJ 1986: 19-, SLOVENIA; FIG. 12

--US, PETKOVSKI + BRANCELJ 1985: 6, SLOVENIA

Morariopsis typica Borutzky ♀

- Borutzky, 1931, p. 195: fig. 80
- , BORUTZKY, 1952, P. 363, F. 99, 100, 13-15. USSR.
- , HANG, 1948, P. 1156.
- , Sterba, 1967, p. 288, 333
- , Jakobi, 1959, p. 277

Habitat Russia

- , BORUTZKY 1960: 147, FIG. 63(6)