

HARPACTICOIDA

NEOBRADYIDAE

- POR 1984: 2

in)

~~Ectinosomidae~~ - Harpacticoida

Neobradya.

T. Scott, 1892.

N. pectinifer.

Syn. Neobradya Scott, 1892^b, p. 249

Neobradya Sara, 1911, p. 439.

Neos new and Bradya.

- Huys, 1987: 133, 134, cf. ANTARCTICOBRADYA, N. GEN.

- Schminke & Dahms 1989: 259, cf. ANTARCTICOBRADYA

Body elongate, cylindrical, without demarkation between anterior and posterior parts. Genital seg't. not subdivided; caudal rami small. First antennae elongate, like those in stenocaris. Exopod of posterior pair fully developed. Mandibles with large biramose palp. Posterior mxpds. not prehensile, 3-jointed. Both rami of 1st. legs 3-jointed; endopod of 3 following pairs 2-jointed. Last legs very small. One ovisac.

Neobradynidae

Neobradya T. Scott, 1892

card 2

- Por 1984:6-

- DAHMS 1992b: 282

Actinosomidae-Harpacticoida.

Neobradya pectinifer T. Scott, ♀. ①

— T. Scott, 1892, p. 249: pl. 13, figs. 19-32.

— T. Scott, 1906, p. 306.

— era Sara, 1911, p. 440: Suppl. pl. 53.

—, LANG, 1948, p. 265, Pl. 136. - NORWAY

--, Jakobi, 1959, p. 145, 279

Habitat. British Seas; Norway Coast.

- PECTINIFERA, POR 1984: 3, FIG. 1

-- HUYS, 1987: 133 - figs. 1-7 tab 1

SARAH... 1989: 249 ... PECTINIFERA 259

Neobradya pectinifer T. Scott, 1892

card 2

- pectinifera, Dahms 1992b: 282