

POECILOSTOMATOIDA

NEREICOLIDAE CLAUS, 1875

- STOCK 1986b: 184

- GOTTO 1993: 136-, DIAGNOSIS, KEY

in) Notodelphysida — Nereiolidae.  
Nereicola, Poecilostomatoida  
Keferstein, 1863, N. ovata <sup>(M)</sup>

Syn. Nereicola Keferstein, 1863, p. 461.

- GOTTO 1993: 136, DIAGNOSIS
- O'REILLY 1995: 55 -

Nereis and cola, inhabiting.

Female. Body (cephalothorax + <sup>thorax</sup> abdomen) broadly ovate, postabdomen (really v abdomen) small, triangular. 2 pairs of antennae many-jointed and armed with setae. Mouth in v form of a sucking beak. Maxillae furnished with hooks. Swimming feet small, 1<sup>st</sup> + 2<sup>nd</sup> pairs biramous 3<sup>rd</sup> rudimentary, papillate. Abdomen with many-jointed, slender caudal appendages: egg-sacs 2, long, cylindrical - male unknown -

(from Claparède, 1870, p. 8).  
(identical with Keferstein's original diagnosis).

Nereicolidae-Notodelphyoidea.

Nereicola concinna T. Scott, ♀.

— T. Scott, 1902, b, p. 455: pl. 25, f. 8-24,

Habitat. On the annelid Eulalia viridis.

English Seas.

Nereididae - Nereididae - Nereididae  
Nereicola ovata Keferstein, ♂. ♀

- — Keferstein, 1863, p. 461: pl. 42, figs. 1-4,  
no name MacIntosh, 1870, p. 39: pl. 5.  
— — Claparède, 1870, p. 18: Carus, 1885, p. 356.  
— — Sars, 1870, p. 113.

Host, Nereis beaucoudragii; N. cultrifera.

- Monod, 1931: 403, fig 2-4  
Mediterranean.

*Nereidicola bipartita* Grube, 1869, p.

*Nereicola ovata* Levinsen, 1878, p. 352.

— — Stenta, 1904, p. 195.

— — Dantan, 1929, p. 175: 20 text-figs.

Nereicola ovatus Keferstein, 1863

(2)

--, Stock, J., 1966, p. 110

--, Stock, 1969, p. 220-221. (ovata)

-- DUDLEY, P. 1986: 19-

NEREIDICOLA OVATA, BRIAN 1932: 58

NEREIDICOLA OVATUS, LAUBIER 1965: 632-637,

FIGS. 1-4, BANYULS - SUR-MER, FRANCE

N. OVATUS, GOTTO 1993: 136, FIG. 32N-P