

Canthocamptidae - Harpacticoida,

Parameira synonym.

Sars, 1907.

P. parva [♂].

Syn. Parameira - Seidlitz, Coleoptera - 1868.

Parameira Sars, 1907, p. 221.

Changed to Sarsameira Wilson.

Ameira (pars) Boeck, Scott,

Παρά and Ameira.

Body short and stout, cylindrical; caudal rami small; eye wholly absent. First antennae short, 8-jointed, some setae on the anterior edge plumose. Exopod of 2nd. antennae 1- or 2-jointed. Mandible palp simple, 2-jointed. Endopod of 1st legs longer than exopod, 3-jointed; last joint in exopod of 3rd. & 4th. and endopod of 3rd. legs with 3 setae inside.

Parameira difficilis Smirnov 1946

-- Smirnov 1946: 238-239, 259, fig. 8, n. sp. ♀ only,
Arctic Ocean

Parameira intermedia (Scott)

--, Shen & Bai, 1956, p. 231

-- SMIRNOV 1946: 238

Canthocamptidae - Harpacticoida.

Parameira major G.O. Sars, ♀.

— — Sars, 1907, p. 223; pl. 147.

--, Shen & Bai, 1956, p. 232

Changed to Sarsameira major.
Wilson,

Habitat, Norway Coast,

Canthocamptidae - Harpacticoida.

Parameira parva (Boeck), ♀.

Ameira — Boeck, 1872, p. 49.

Parameira — Sars, 1907, p. 222: pl. 146.

Ameira longiremis, var., intermedia T. Scott.

Changed to Sarsameira parva Wilson,

Habitat. Scottish Coast (Scott); Norway (Sars).

---. Shen & Bai, 1956, p. 232

— Willey, 1923: 330

SHIBATA 1946: 238 239

Parameira pendula Shen & Bai, 1956

Parameira pendula Shen & Bai, 1956, pp. 211-213,
pl. 12, fig. 90-95, pp. 231-2.

-- CHEN, Q.-C. 1986: 524-

Canthocamptidae - Harpacticoida.

Parameira propinqua (T. Scott), ♀.

Ameira — T. Scott, 1902, a. p. 460: pl. 24. f. 10-18.

Parameira — Sars, 1911, p. 399: Suppl. pl. 28.

Changed to Sarsameira propinqua.
Wilson,

Habitat. Scottish Coast: Norway.

Parameira simplex

--, Jakobi, 1959, p. 139