

Cyclopora da: Notodelphyota - Notodelphyota

Paranotodelphys. valid. ①

Schellenberg, 1921^b.

P. scutiformis

Syn. Paranotodelphys Schellenberg, 1921^b p. 232

Paranotodelphys Schellenberg, 1922

- Stock 1967e: 11, key to ♀♀

Paranotodelphys constricta Illg, 1970

n. sp. Illg, 1970, pp. 172-6, figs. 1, 2

Paranotodelphys engeli Stock 1967

-- Stock 1967e: 3, 11, 12-14, figs. 8-11, n. sp. ♀, from
Pyuridae, Red Sea.

Paranotodelphys furcifera Stock 1967

-- Stock 1967e: 3, 11, 16-18, figs. 16-17, n. sp. ♀, from
Pyuridea, Red Sea.

Notodelphyoidea -

Paranotodelphys gracilis Schellenberg.

--- Schellenberg, 1921^b, p. 235: text-figs. 12-14.
-- STOCK 1967c: 11, 20

Host. Rhodosoma verecundum

Gulf of Suez.

Notodelphyoidea.
Paranotodelphys longicauda Schellenberg.

— Schellenberg, 1921^b, p. 236; text-figs. 15-16.
— STOCK 1967e:11.

Host, Unknown - from 385m. deep.

Paranotodelphys phallusiae (Gurney, 1927)

-- Stock 1967e: 11, 13-14

Paranoto delphys procax Stock & Humes, 1970

--, Stock & Humes, 1970, p. 195-200, figs. 1-3

Paranotodelphys saccata Stock 1967

-- Stock 1967e: 3, 11, 14-16, figs. 12-15, n. sp. ♀♂, from

?Polycarpa or ?Styela, Red Sea

Paranotodelphys scutiformis Schellenberg.
Notodelphyoidea.

— — Schellenberg, 1921^b, p. 233: text figs. 9-11.

Host, Ascidia malaca Travst.; Ascidiella aspersa
(Müller): Ascidiella latesiphonica, and
Ascidia geminata,
S.W. Australia.

Paranotodelphys scutiformis Schellenberg, 1922

-- Stock 1967e: 11, 16

Paranotodelphys villosus 0015#1, 1963

-- Ooishi 1987:95, Japan