

in) Tachidiidae - Harpacticoida.

Psammis,

G. O. Sars, 1910

P. longisetosa (M)

Syn. Psammis Sars, 1910, p. 339.

Psammis, a proper name.

. Body subcylindric, no demarkation between anterior and posterior parts, all seg'ts closely crowded together. Caudal rami moderate. First antennae short thick, hirsute; exopod of second pair 3-jointed. Endopod of first legs 2-jointed. Distal and proximal joints in last legs fused.

PARANANNOPIDAE

PSAMMIS

Sars

2

- , BODIN 1968: pp.32, Key to species - 35.
- GEE 1988a: 52
- GEE 1988b: 186-187, 199- IN KEY
- HUYS + GEE 1992: 32, "BOUNDARY BETWEEN DANIELSSENIA + PSAMMIS SUPERFLUOUS"
- GEE + HUYS 1990: 1549-, IN PARANANNOPIDAE
- GEE + HUYS 1991: 1135-
- GEE + HUYS 1994: 1008-

Psammis borealis Klie

— — Klie, 1939, p. 223. Iceland.

— —, Jespersen, 1940, p. 74. Iceland.

--, Klie, 1941, p. 7-10

--, COULU, 1971: 207

--, BODIN 1968: p. 35 - Key to species.

-- GEE 1988b: 186

-- GEE & HUYS 1990: 1566 -

-- GEE & HUYS 1994: 1008 - , not in PSAMMIS

-- HUYS 1990: 47

Psammis kliei - see Mucrosenia

Psammis longifurca Bodin - see Bathypsammis

PSAMMIS LONGIPES BECKER, 1974

- BECKER, 1974: 2, 3; FIG. 3, 4
- GEE 1988b: 186
- GEE & Huys 1990: 1566-
- GEE & Huys 1994: 1008-

Tachidiidae - Harpacticoida.

Psammis longisetosa G. O. Sars, ♀.

— — Sars, 1910, p. 339: pl. 225,

— —, Handl. 1948, p. 303, Pl. 148 -

-- --, Wells, 1965. p. 3

-- --, Por, 1965, pp. 3-4, figs. 1-8

--, Jakobi, 1959, p. 141, 280

--, Soyer, 1970, p. 269

Habitat. Norway Coast.

--, Klie, 1941, p. 8-10

--, BECKER, 1974: 3

--, COULL, 1971: 207



--, BODIN 1968: p. 35 - Key to species.

-- GEE 1988b: 181-187, FIGS. 1-4, REDESCRIBED -

NORWAY

-- SMIRNOV 1946: 256, 257

-- GEE & HAYS 1990: 1566 -

-- GEE & HAYS 1994: 1008 -

PSAMMIS sp.

-- ECKMAN + THISTLE 1991: 1405-, OFF CALIFORNIA