

(12)

Sarcotretes valid

Caligoida

Lernaeidae.

Jungersen, 1911

S. scopeli ^(M).

Syn. Sarcotretes Jungersen, 1911, p. 10.

Σάρξ = flesh and τρητής = a borer

Sarcotretes

Pennellidae: SIPHONOSTOMATOIDA

2

- Kabata, 1967, p. 866
- HOBANS 1988: 1371-1375, REVIEWS GENUS
- KABATA 1988: 80, GENUS DIAGNOSIS
- CASTRO + BAEZA 1985C: 27

SARCOTRETES, KABATA 1972: 921, IN KEY TO
GENERA OF LERNAEOCERIDAE

Sarcotretes cristaleiformis (Brian) ♀ ^{Uerpaecidae.}

Uerpaecinus — Brian, 1908, p. 13: fig. 5. A & B, text.
— — Brian, 1912, p. 20: pl. 2, fig. 4: pl. 4,
figs. 1 & 2: pl. 8, fig. 2: pl. 9 & 10, fig. 6-14.
Sarcotretes — Wilson, 1917, p. 72.

Host, Bathypterois dubius, an abyssal fish
— Kazatchenko & Avdeev — 1977: 38

North Atlantic

ernaeinus - Brian, 1914, p. 10, fig. 7, text (pl. 10, f. 6)

--- Brian, 1923, p. 20: pl. 2, f. 4: pl. 4, f. 1-2: pl. 8, f. 2: pl. 9;

Sarcotretes - Brian, 1929, p. 22: pl. 2, f. 10: pl. 5, f. 18

-- --, ~~Kabata~~ / Kabata, 1966, p. 173.

-- HOGANS 1988: 1371-1375, REVIEWS RECORDS

-- BOXSHALL 1989: 207

Lernaeidae.

Sarcotretes gempyli (Horst) ♀

Lernaeenicus — Horst, 1878, p. 51: pl. 7.

Sarcotretes — Wilson, 1917, a. p. 71,

S. —, HOGANS 1988: 1371-, POSSIBLE SYNONYM OF S. SCOPELI.

S. —, BOXSHALL 1989: 207

Host, Gempylus serpens

Atlantici.

Sarcotretes inflexus (Stp. & Totk.) ♀ ①

Ternaeicus — Stp. & Totk, 1861, p. 401: pl. 13, fig. 21

— — Richiardi, 1877^c, p. 8 [202].

— — B-Smith, 1899, p. 485.

Sarcotretes — Wilson, 1917, a, p. 71.

Host, Gills of barroecuda, Sphyraena vul-
garis Cur. & Val.

Atlantic.

Sarcotretes inflexus (Stp. & Ltk.)

card 2

-- Hogans 1988: 1371-1375, probable synonym of S. scopeli

S. -, BOXSHALL 1989: 207, NOW IN LERNAEENICKS

Sarcotretes lobatus New Species, ♀.

— Wilson, 1917, a, p. 72; pl. 8, f. 71-79.

— Wilson, 1932, p. 484; fig. 291.

-- HOGANS 1988: 1371-1375, PROBABLE SYN. OF S. SCOPELI

-- BOXSHALL 1989: 207-208, SYN. OF S. SCOPELI

Host Benthosema (Scopelus) mülleri

N. Eng. Coast.

Sarcobates nodicornis (Stp. & Ltk.) ♀

Lernaeicus — Stp. & Ltk, 1861, p. 401: pl. 13, fig. 26.

Lernaeonema — Beneden, 1870, p. 359.

Lernaeicus — Ricciardi, 1877^c, p. 8 [202].

— —, B-Smith, 1899, p. 485.

Host. Coryphaena sp.; Delphinus delphis.
in body muscles.

European Seas.

Sarcotretes — Wilson, 1917, a, p. 71 (key).

S. -, HOGANS 1988: 1371-, POSSIBLE SYNONYM OF S. SCOPELLI

S. -, BOXSHALL 1989: 205, 207, NOW IN LERNAEENICUS

Sarcotretes scopeli Jungerson, 1911

card 2

-- Hogans 1988: 1371-1375, figs. 1-7, Atlantic records, redescription; probable synonyms: S. inflexus, S.

lobatus, S. gempyli, S. nodicornis

-- KABATA 1988: 80-81, FIGS. 36 C, D; HOST BENTHOSOMA

GLACIALIS, CANADA

-- BOXSHALL 1989: 205-208, FIGS. 3, 4, NEW

CALÉDONIA; HOST HYMENOCEPHALUS GRACILIS.

SARCOTRETES SP.

-- PERKINS 1983: 73, RECORD IN TRIPHOTURUS
MEXICANUS

Sarcotretes scopeli Jungersen. Dermatidae.

SCOPELI

- —, Jungersen, 1911, p. 1: pls. 142, fig. 1-27.
— —, Stephensen, 1913, p. 348.
— —, Wilson, 1917, p. 69, 70, 71, (key).
— —, Wilson, 1920, p. 146.

Host. Scopelus glacialis (Benthosema mülleri).

— Kazatchenko & Audeev, 1977: 38-39

Faroes: Greenland.

-- Hansen, 1920 1

-- SPROSTON 1942: 448.

-- PERKINS 1983: 70.