

SIPHONOSTOMATOIDA - LERNAEOPODIDAE

SCHISTOBRACHIA KABATA, 1964

- KABATA 1988: 42, 59-60, DIAGNOSIS, KEY TO CANADIAN SPP.
- KABATA 1988d: 551-, SP. LIST
- KABATA 1992: 142, 144, DIAGNOSIS
- CHANDRAN + BALAKRISHNAN NAIR 1980b: 446

= SCHISTOBRACHIA kernaepodidae - kernaepodocida

Charopinus chimaerae Yamaguti, ♀.

— — Yamaguti, 1939, p. 552: pl. 46, figs. 122-127.

Host. Gills, *Chimaera ogilbyi* Waite.

Nagasaki, Japan.

Schistobranchia chimaerae (Yamaguti, 1939)

-- --, Kabata, 1964, p. 101.

--, Kabata, 1970, p. 867

-- KABATA 1988d: 551-

Schistobrachia pilgrimi Kabata, 1988

-- Kabata 1988d:551-555, figs. 1-12, n. sp. ♀♂, host

Harriotta raleighana, New Zealand

= SCHISTOBRACHIA RAMOSA

Лемнаепоцнне.

Charopinus ramosus, Kröyer, ♂, ♀.

— —, Kröyer, 1863, p. 284; pl. XIV, fig 5a-i.

— —, B. Smilk, 1899, p. 498.

— —, T. Scott, 1901, p. 130; pl. VII, figs. 17-23.

— —, A. Scott, 1904, p. 123; 1904, a, p. 43

Host. Raja clavata. Gylls, R. maculata.

Europe.

- — Wilson, 1915, p. 654 (key),
— — Hansen, 1923, p. 50.
— — Leigh-Sharpe, 1928, p. 276: fig. 4, A-C.
--, Polianskii, 1955, p. 92-
--, Stephensen, 1940, p. 9

Schistobranchia ramosa (Krøyer, 1863)

-- --, Kabata, 1964, p. 99.

--, Kabata, 1970, p. 867

--, Kabata, 1970, fig. 54A

-- KENSLEY & GRINDLEY 1973: 114-15, FIG. 31, ♀ FROM
RAIA BATHS, TABLE BAY, S. AFRICA; p. 128.

-- KABATA 1988: 60, FIG. 28A-D; SYNONYMY.

-- KABATA 1988d: 551-

-- KABATA 1992: 144-, FIG. 32 D-G, SYNONYMY, HOST LIST

Schistobranchia tertia Kabata, 1970

- n. sp., Kabata, 1970, pp. 866-867, fig. 1-16.
-- , Kabata & Cousens, 1972, passim
-- KABATA 1988: 60, FIG. 28E-F
-- KABATA 1988d: 551-