

in)

Tachidiidae - Harpacticoida.

Tachidiopsis.

G. O. Sars, 1911.

T. cyclopoides. ^(M)

Syn. Tachidiopsis Sars, 1911, p. 434.

Tachidius and opsis, likeness.

General form like Tachidius; anterior division wider than posterior. Gen. segt. in ♀ imperfectly divided. First antennae with more joints than in Tachidius; exopod of posterior pair fully developed. A complex masticatory lobe inside the basal joint of 2nd. mxpds. Both rami of 1st. legs 3-jointed, endopod the longer and bent in the middle. Endopod of 2nd. & 3rd. legs 2-jointed, of 4th. legs 3-jointed. Last legs small.

TACHIDIOPSIS

SARS

--, DINET, 1974: 562 - KEY TO SPP.

--, COLL, 1973d: 594-596 - KEY TO FEMALES

--, BODIN 1968: pp. 52, 59. Key to species.

-- KUNZ 1981a: 3-5

TACHYDIOPSIS, COTTARELLI + BALDARI 1987b: 183

TACHIDIOPSIS BOZICI

BODIN, 1968

--, BECKER, 1974: f

--, DINET, 1974: 562

--, COULL, 1973d: 595 & TAB. 1

--, BODIN 1968: pp. 52-57; Pls. XXX - XXXII, ♀+♂;
p. 59 - Key to species.

Tachidiidae - Harpacticoida.

Tachidiopsis cyclooides G.O.Sars, ♀. ①

— — Sars, 1911, p. 435; Suppl. pl. 51.

--, Jakobi, 1959, p. 280

--, BECKER, 1974: 4

--, DINET, 1974: 562

--, COLL, 1973d: 594

Habitat. Norway Coast.

--, BODIN 1968: p. 59 - Key to species.

-- KUNZ 1981a: 5

TACHIDIOPSIS CYCLOPOIDES G.O. JARS, 1911

(2)

-- HUYS 1996: 541

TACHIDIOPSIS IBERICUS (BECKER, 1974)

--BECKER, 1974: 2, 4, 21; FIG. 7, 8

TACHIDIOPSIS LAUBIERI DINET, 1974

562

-- DINET, 1974: 554-558²; FIG. 3, 4

TACHIDIOPSIS PARASIMILIS DINET, 1974

-- DINET, 1974: 558-562 ♀; FIG. 5, 6

TACHIDIOPSIS SARSI

BODIN, 1968

--, BECKER, 1974: 4

--, DINET, 1974: 562

--, COLL, 1973d: 596

--, BODIN 1968: pp. 57-59; Pl. XXXIII, ♀; p. 59

Key to species.

TACHIDIOPSIS SIMILIS (DRZYCIŃSKI, 1968)

--, BECKER, 1974: 4

--, DINET, 1974: 562

--, COLL, 1973d: 594, 596; TAB. 1

TACHIDIOPSIS TYPICA

- BECKER, 1974: 4
- , DINET, 1974: 558, 562
- , COULL, 1973d: 596