

A Systematic Study on the Marine Sponges in Korea 9 Ceractinomorpha

Sim, Chung Ja and Byeon, Hyo Sook

(Department of Biology, Han Nam University, Daejeon 300-791, Republic of Korea)

한국산 해산해면류의 계통분류학적 연구

9. 일축해면류

심 정 자 · 변 효 숙

(한남대학교 생물학과)

적 요

한국산 일축해면류를 연구하기 위해 삼면 연안과 16개 도서지방에서 1987년부터 1988년 8월까지 채집한 표본과 그동안 한남대학교 생물학과에 미해결종으로 보관되어 있던 표본들을 동정, 분류한 결과 12과 21속 41종이 분류되었고 이중 13종은 한국미기록종이고 2종은 신종이었다. 한국미기록종과 신종에 대하여는 특기 또는 기재와 도판을 첨가하였다.

Key words: systematics, marine sponge, Ceractinomorpha, Korea.

INTRODUCTION

On the Systematic study of Korean marine Ceractinomorpha sponges, 66 species were recorded from Korean waters (Kim *et al.*, 1968; Rho *et al.*, 1969; Rho & Sim, 1972a, 1972b, 1976, 1979; Rho & Lee,

This work was supported by grant from the Korea Research Foundation in 1988.




Fig. 1. A map showing the localities where the specimens were collected.

1, Namae; 2, Ullungdo; 3, Mip'o; 4, Samch'ŏnp'o; 5, Mijo-ri; 6, Kŭmodo; 7, Sŏdo; 8, Kŏmundo; 9, Sŏngsanp'o; 10, Sŏgwip'o; 11, Mosŭlp'o; 12, Piyangdo; 13, Ch'ujado; 14, Anmyŏndo; 15, Anhŭng; 16, Chakyakto.

1976; Sim, 1981, 1982, 1985; Rho & Yang, 1983; Sim & Bae, 1987; Sim & Kim, 1988).

The present study is based on the materials which were collected from 16 localities during the period from 1987 to 1988 (Fig. 1), and some unidentified specimens preserved in the Department of Biology, Han Nam University. The identified Ceractinomorpha consist of 41 species, 21 genera and 12 families. Among them, 2 species were new and 13 species were newly recorded in Korea.

LIST OF SPECIES

Phylum Porifera Grant, 1836

Class Demospongia Sollas, 1885

Subclass Ceractinomorpha Levi, 1953

해면동물 문

보통해면강

일축해면 아강

Order 1. Halichondrida Levi, 1953 해변해면 목
Family 1. Halichondriidae Gray, 1867 해변해면 과

1. *Halichondria panicea* (Pallas, 1776) 회색해변해면
2. *Halichondria okadai* (Kadota, 1922) 검정해변해면
3. *Halichondria oshoro* Tanita, 1961 황록해변해면

Family 2. Hymeniacidonidae De Laubenfels 1936 하이메니아시돈 과

4. *Hymeniacidon sinapium* De Laubenfels, 1930 주황해변해면

Order 2. Poecilosclerida Topsent, 1928 다골해면 목
Family 3. Ophlitaspongiae De Laubenfels, 1936 바늘뼈해면 과

5. *Ophlitaspongia noto* Tanita, 1963 바늘뼈해면
6. **Ophlitaspongia pennata californiana* De Laubenfels, 1932 캘리포니아바늘뼈해면 (신칭)
7. *Mycale adhaerence* (Lambe, 1893) 깃해면
8. *Neofolitispa dianchora* (De Laubenfels, 1935) 두닷바늘뼈해면
9. **Desmacella rosea* Fristedt, 1887 장미띠해면 (신칭)
10. *Clathria spinispicula* Tanita, 1968 침유령해면
11. *Clathria toxipraedita* Topsent, 1913 화살유령해면
12. *Clathria terranova* Dendy, 1924 창살유령해면
13. *Clathria madrepora* Dendy, 1921 돌산호유령해면
14. **Clathria mosulpia* sp. nov. 모슬침유령해면 (신칭)
15. *Clathria dayi* Levi, 1963 데이침유령해면 (신칭)
16. *Clathria parva* Levi, 1963 작은침유령해면 (신칭)
17. *Axociella cylindrica* Hallman, 1920 둥근촉털해면
18. *Axociella calla* (De Laubenfels, 1934) 깃촉뼈해면

Family 4. Microcionidae Hentschel, 1923 유령해면 과

19. *Microciona longistyla* Burton, 1959 기둥유령해면

Family 5. Myxillidae Hentschel, 1923 끈적해면 과

20. *Myxilla setoensis* Tanita, 1961 넓적끈적해면
21. *Myxilla incrustans* (Johnston, 1842) 껍질끈적해면
22. *Myxilla productus* Hoshino, 1981 긴끈적해면
23. *Myxilla bivalvia* Tanita, 1967 덩게끈적해면
24. **Myxilla sigmatifera* (Levi, 1963) 시그마끈적해면
25. *Lissodendoryx isodictyalis* (Carter, 1882) 두드럭끈적해면
- Family 6. Tedaniidae Ridley & Dendy, 1887 테다니해면 과
26. *Iotrochota baculifera* Ridley, 1884 보라바귀해면
27. *Tedania brevispiculata* Thiele, 1903 테다니해면
- Family 7. Plocamiidae Topsent, 1928 곱슬해면 과
28. *Lissoplocamia tocushima* Tanita, 1970 미끈이해면
- Order 3. Haplosclerida Topsent, 1928 단골해면 목
- Family 8. Haliclونidae De Laubenfels, 1932 보라해면 과
29. *Haliclona permollis* (Bowerbank, 1866) 보라해면
30. *Haliclona densaspicula* Hoshino, 1981 뽁뽁침보라해면
31. **Haliclona perlucida* (Griessinger, 1971) 진주보라해면 (신칭)
32. **Haliclona ulreungia* sp. nov. 울릉보라해면 (신칭)
- Family 9. Adociidae De Laubenfels, 1936 아도시해면 과
33. *Petrosia ushitsuensis* Tanita, 1963 바위해면
34. **Petrosia nigricans* Lindgren, 1897 검정바위해면 (신칭)
- Family 10. Renieridae Schmidt, 1870 레니에해면 과
35. **Gellius arcoferus* Vosmaer, 1885 판숨덩이해면 (신칭)
36. **Reniera ventilabrum* Fristedt, 1887 판레니에해면 (신칭)
37. **Reniera pigmentifera* Dendy, 1905 보라레니에해면 (신칭)
- Family 11. Callispongiidae De Laubenfels, 1936 예쁜이해면 과
38. *Callispongia elegans* (Thiele, 1899) 예쁜이해면

39. *Callispongia confederata* (Ridley, 1884) 보라예쁜이해면
 40. *Callispongia elongata* (Ridley & dandy, 1886) 길쭉예쁜이해면
 Family 12. *Coelosphaeridae* Hentschel, 1923 강해면 과(신칭)
 41. *Coelosphaera physa* (Schmidt, 1875) 거품강해면(신칭)

The asterisks (*) indicate the species which were newly recorded in Korea and asterisks (**) indicate the species which were new.

SYSTEMATIC ACCOUNT

Family *Halichondriidae* Gray, 1867 해변해면 과

1. *Halichondria panicea* (Pallas, 1766) 회색해변해면
 Material examined: Söngsanp'o, 18/V/1983; Sögwip'o, 19/VII/1987; Södo, 29/VII/1988; Kūmodo, 31/VII/1988.
 2. *Halichondria okadai* (Kadota, 1922) 검정해변해면
 Material examined: Mijo-ri, 13/VII/1983; Söngsanp'o, 8/VII/1985; Sögwip'o, 18/VII/1987; Södo, 29/VII/1988; Kūmodo, 31/VII/1988.
 3. *Halichondria oshoro* Tanita, 1961 황록해변해면
 Material examined: Kūmodo, 31/VII/1988.

Family *Hymeniacidonidae* De Laubenfels, 1936 하이메니아시돈 과

4. *Hymeniacidon sinapium* De Laubenfels, 1930 주황해변해면
 Material examined: Söngsanp'o, 8/VII/1985; Anhūng, 8/V/1987, 14/V/1988; Sögwip'o, 18/VII/1987, 28/VII/1988; Kōmundo, 24/VII/1988; Södo, 29/VII/1988; Kūmodo, 31/VII/1988.

Family *Ophlitaspongiidae* De Laubenfels, 1936 바늘땀해면 과

5. *Ophlitaspongia noto* Tanita, 1963 바늘땀해면
 Material examined: Sögwip'o, 18/VII/1987; Chakyakto, 16/IV/1988; Anhūng, 14/V/1988.
 6. **Ophlitaspongia pennata californiana* De Laubenfels, 1932 캘리포니아바늘땀해면
 (Pl. 1, Figs. 1-4)

Ophlitaspongia pennata (Lambe, 1895) *californiana* De Laubenfels, 1932 (pp. 103-104, text-fig. 62).

Ophlitaspongia pennata californiana: Bakus & Green, 1987(p. 73).

Material examined: Sögwip'o, 1/VII/1984, 5/VIII/1986; Sögwip'o (fish net), 6/VII/1985.

Remarks: This sponge is thin encrusting. Size up to 1.5mm thick spreading laterally indefinitely. It is similar to *Ophlitaspongia noto* in shape, but differs in skeletal arrangement. The texture is firm and the surface of the body is superficially velvety. Colour in life is vermilion, but in spirits is dirty brown.

Spicules: Subtylostyle	338-353 × 15-18μm.
Slender subtylostyle	330-483 × 4-9μm.
Style	266-435 × 13-18μm.
Large toxa	69-140 × 2-4μm.
Small toxa	12-46μm.

Distribution: Korea (Cheju Island), California, Mexico.

7. *Mycale adhaerens* (Lambe, 1893) 깃해면

Material examined: Mosūlp'o, 29/VII/1988; Kōmundo, 29/VII/1988; Namae, 9/VIII/1988.

8. *Neofolitispa dianchora* (De Laubenfels, 1935) 두닷바늘뼈해면

Material examined: Piyangdo, 20/VI/1985; Mosūlp'o, 6/IX/1986, 18/VII/1987.

9. **Desmacella rosea* Fristedt, 1887 장미띠해면 (Pl. 2, Figs. 1-2)

Desmacella rosea Fristedt, 1887 (pp. 439-440, pl. 24, figs. 32-35, pl. 28, fig. 13).

Material examined: Ch'uja-do (100 depth), 6/II/1982.

Remarks: This sponge is flat or leaf-like shape and size, up to 7cm × 6.5cm × 1cm. Texture is firm and surface is smooth. Colour in life is yellow, but in spirits is ivory.

Spicules: Large tylostyle	700-825 × 8-13μm.
Middle tylostyle	401-591 × 7-12μm.
Small tylostyle	181-253 × 2-6μm.
Large sigma	32-41μm.
Small sigma	18-26μm.

Distribution: Korea (Korea Strait), the Atlantic Ocean, Bering Sea, Arctic Ocean.

10. *Clathria spinispicula* Tanita, 1963 침유령해면

Material examined: Mosūlp'o, 8/IV/1987, 19/VII/1987.

11. *Clathria toxipraedita* Topsent, 1913 화살유령해면

Material examined: Mosūlp'o, 18/VII/1987.

12. *Clathria terranova* Dendy, 1924 창살유령해면

Material examined: Mosūlp'o, 8/VI/1987, 18/VII/1987.

13. *Clathria madrepora* Dendy, 1921 돌산호유령해면

Material examined: Sōgwip'o, 15/VII/1982; Mosūlp'o, 18/VII/1987.

14. ***Clathria mosulpia* sp. nov. 모슬침유령해면 (Pl. 3, Figs. 1-5)

Material examined: Three specimens (Por. 9, Por. 9-1, Por. 9-2), Mosūlp'o (fish net), 18/VII/1987, H.S. Byeon & H.Y. Kim.

Holotype: Por. 9, Deposited in Natural History Museum of Han Nam University.

Paratype: Por. 9-1, Por. 9-2, Deposited in the Department of Biology, Han Nam University.

Description: This sponge is flabellate or leaf-like shape and size up to 4cm × 7cm × 0.5cm. Texture is tough and elastic. The surface of the body is a little velvety with the protruding spicules. Pores are

dispersed on the body and honeycomb-like form, but the oscules are not observed. Dermal membrane is covered with spicules which look branch. Colour in life is yellow, but in alcohol is ivory.

Spicules: Style 242-266 × 12-14 μ m.
 Acanthostyle 91-100 × 7 μ m.
 Large spiny toxa 522-768 × 5- 7 μ m.
 Toxa 181-467 × 2- 5 μ m.
 Isochela 12-18 μ m.

Remarks: The new species is similar to *Clathria barleei* De weerd & Van soest 1986 in shape, but differs in spicules. *Clathria barleei* has no spiny toxa, but the new species has large spiny toxa and very rare strongyle.

Distribution: Korea (Cheju-Island).

15. **Clathria dayi* Levi, 1963 데이 침유령해면 (Pl. 4, Figs. 1-2)

Clathria dayi Levi, 1963 (pp. 51-52, pl. 8, fig. B, text-fig. 58).

Material examined: Mosŭlp'o (fish net), 18/VII/1987.

Remarks: This sponge is a branch type, measuring up to 2cm in diameter, 20cm in length. Texture is elastic and surface is uneven. Dermal membrane is thin. Pores and oscles are dispersed on the body. Colour in spirits is pale brown.

Spicules: Thick style 139-260 × 10-17 μ
 Slender style 253-358 × 1- 4 μ m.
 Acanthostyle 100-130 × 4- 6 μ m.
 Spiny toxa 213-346 × 3- 4 μ m.
 Toxa 60-106 × 1- 2 μ m.
 Isochela 15-35 μ m.

Distribution: Korea (Cheju-Island), South Africa.

16. **Clathria parva* Levi, 1963 작은침유령해면 (Pl. 4, Figs. 3-4)

Clathria parva Levi, 1963 (pp. 56-57, pl. 10, fig. D, text-fig. 64).

Material examined: mosŭlp'o (fish net), 18/VII/1987.

Remarks: Sponge covered coral branch, 2mm in diameter, surface of the body is smooth. Colour in spirits is pale brown.

Spicules: Style 157-280 × 2- 6 μ m.
 Spiny style 240-300 × 7-12 μ m.
 Acanthostyle 95-110 × 5- 8 μ m.
 Spiny toxa 100-123 × 2 μ m.
 Toxa 90- 99 × 2 μ m.
 Isochela 12-13 μ m.

Distribution: Korea (Cheju-Island), South Africa.

17. **Axociella cylindrica* (Ridley & Dendy, 1886) 둥근축털해면 (Pl. 5, Figs. 1-2)

Eспериopsis cylindrica Ridley & Dendy, 1886, [cited from Hallmann, 1920 (pp. 780-784)].

Axociella cylindrica: Hallmann, 1920 (pp. 780-784, pl. 37, figs. 1-3, text-fig. 2).

Material examined: Mosŭlp'o (fish net), 18/VII/1987.

Remarks: It is a sparsely and dichotomously ramose sponge with relatively long and slender, flattened branches measuring up to 1.75cm in diameter, 22cm in length. Texture is dense and consistency fairly firm and tough. In alcohol the colour is dark-grey.

Spicules: Large trachy style 450-570 × 20-24μ.
 Trachy style 274-354 × 12-24μm.
 Slender style 415-660 × 2 - 6μm.
 Toxa 45-108 × 2 - 4μm.
 Large isochela 13-23μm.
 Small isochela 10-12μm.

Distribution: Korea (Cheju-Island), Port Jackson.

18. *'Axocielita calla* (De Laubenfels, 1934) 갯축해면 (Pl. 5, Figs. 3-5)

Axociella calla De Laubenfels, 1934 (p.16).

Axocielita calla: De Laubenfels, 1954 (p. 149).

Material examined: Songsanp'o, 30/VI/1984.

Remarks: This sponge is incrusting 2mm-4mm thickness. Surface of the body is a little velvety. Texture is hard. Colour in life is red, but in spirits is dark purple.

Spicules: Subtylostyle 180-250 × 7-12μm.
 Slender style 180-230 × 2- 3μm.
 Large toxa 103-143 × 2- 6μm.
 Small toxa 21-48μm.
 Isochela 12μm.

Distribution: Korea (Cheju-Island), the Atlantic (Caribbean Sea), Puerto Rico.

Family Microcionidae Hentschel, 1923 유령해면 과

19. *Microciona longistyla* Burton, 1959 기둥유령해면

Material examined: Mosŭlp'o, 5/IV/1987.

Family Myxillidae Hentschel 1923 끈적해면 과

20. *Myxilla setoensis* Tanita, 1961 넓적끈적해면

Material examined: Mosŭlp'o, 18/VII/1987; Mipo, 25/X/1987.

21. *Myxilla incrustans* (Johnston, 1842) 꺾질끈적해면

Material examined: Mosŭlp'o, 19/VII/1987.

22. *Myxilla productus* Hoshino, 1981 긴끈적해면

Material examined: Sŏgwip'o, 14/VII/1982; Mosŭlp'o, 18/VII/1987.

23. *Myxilla bivalvia* Tanita, 1967 덮개끈적해면

Material examined: Mosŭlp'o, 18/VII/1987; Mipo, 25/X/1987.

24. **Myxilla sigmatifera* (Levi, 1963) 시그마끈적해면 (Pl. 6, Figs. 1-2)
Burtonanchora sigmatifera Levi, 1963 (pp. 34-35, pl. 6A, text-fig. 37).
Myxilla sigmatifera: Bakus, 1966 (p.512); De Laubenfels, 1936 (p. 94).
 Material examined: Sōgwip'o (fish net), 15/VII/1982; Mosūlp'o (fish net), 18/VII/1987.
 Remarks: Shape is irregular massive. Surface is uneven and texture is fragile. Oscules are observation, size up to 0.5-1mm. Pores are dispersed on the body. Colour in spirits is ivory.
- Spicules: Trachy style 150-302 × 6-16μm.
 Tornote 172-326 × 5-12μm.
 Large sigma 47-104μm.
 Small sigma 12-14μm.
 Large isochela 46-54μm.
 Small isochela 14-18μm.
- Distribution: Korea (Cheju-Island), South Africa.
25. *Lissodendoryx isodictyalis* (Carter, 1882) 두드럭끈적해면
 Material examined: Mosūlp'o, 18/VII/1987.
- Family Tedaniidae Ridley & Dendy, 1887 테다니해면 과
26. *Iotrochota baculifera* Ridley, 1884 보라바퀴해면
 Material examined: Sōgwip'o, 29/VI/1988.
27. *Tedania brevispiculata* Thiele, 1903 테다니해면
 Material examined: Mosūlp'o, 19/VII/1987.
- Family Plocamiidae Topsent, 1928 곱슬해면 과
28. *Lissoplocamia tocushima* Tanita, 1970 미끈이해면
 Material examined: Sōgwip'o, 30/VII/1984.
- Family Haliclونidae De Laubenfels 1923 보라해면 과
29. *Haliclona permollis* (Bowerbank, 1866) 보라해면
 Material examined: Samchōnp'o, 22/VII/1984; Anmyōndo, 24/VII/1984; Kōmundo, 24/VII/1988; Sōdo, 26/VII/1988; Kūmodo, 31/VII/1988.
30. *Haliclona densaspicula* Hoshino, 1981 백백침보라해면
 Material examined: Mosūlp'o, 18/VII/1987.
31. **Haliclona perlucida* (Griessinger, 1971) 진주보라해면 (Pl. 6, Figs. 3-5)
Reniera perlucida Griessinger, 1971 (pp. 127-128, pl. 2, fig. 1, text-fig. 3c, text-figs. 4d and e); Pulitzer-Finali, 1983 [cited from De Weerd & Van Soest, 1986 (p.13)].
Haliclona perlucida: De Weerd & Van Soest, 1986 (pp. 13-16, text-figs. 9-10).
 Material examined: Sōngsanp'o, 30/VI/1984.

Remarks: This sponge is thick massive, size up to 4cm × 30cm × 0.5cm. Oscules are slightly elevated with oscular rims. Surface is smooth and texture is fragile. Colour in life is violet, but in spirits is white yellow. Ectosome is irregular isodictyal reticulum without spongin.

Spicules: Oxea 120-200 × 5-10μm.

Slender oxea 110-180 × 1- 2μm.

Distribution: Korea (Cheju-Island), the Mediterranean Sea, the Canary Islands, the Azores.

32. **Haliclona ulreungia* sp. nov. 울릉보라해면 (Pl. 7, Figs. 1-4)

Material examined: One specimen, Ullungdo (Todong), 11/VII/1984, C.J. Sim.

Holotype: Por. 10, Deposited in Natural History Museum of Han Nam University.

Description: This sponge is irregular massive and size up to 5cm × 3cm × 1cm. Pores are dispersed on the body. Oscules are slightly elevated with oscular rims, and size up to 0.5mm in diameter, Surface is smooth and texture is very soft and fragile. Dermal membrane is specialization. Spicules are irregular reticulation. Colour in spirits is pale ivory.

Spicules: Oxea 103-179 × 3-8μm.

Slender oxea 100-165 × 1-2μm.

Remarks: The new species is similar to *Haliclona permollisimilis* Hoshino, 1981 in spiculation, but differs in sponge surface. The new species is dermal membrane specialization.

Distribution: Korea (Sea of Japan).

Family Adociidae De Laubenfels, 1936 아도시해면 과

33. *Petrosia ushitsuensis* Tanita, 1963 바위해면

Material examined: Sögwip'o, 5/VII/1984.

34. **Petrosia nigricans* Lindgren, 1897 검정바위해면 (Pl. 7, Figs. 5-6)

Petrosia nigricans Lindgren, 1897 [cited from Burton, 1959 (p. 222)].

Petrosia nigricans: Burton, 1959 (p.222).

Petrosia imperforata: Thiele, 1899 (p.20, pl.2, fig.7, pl.5, fig. 12).

Petrosia nigricans var *irregularis*: Hentschel, 1912 (p.405).

Petrosia mammiformis: Dendy, 1921 (p.36, pl.12, fig.6).

Material examined: Sögwip'o (fish net), 14/VII/1982.

Remarks: The sponge is thick-walled, tubular processes, rising from a basal crust and each ending in a wide, circular vent. Oscules are dispersed on the gastral part. Texture is hard, but fragile. Colour in spirits is ivory.

Spicules: Stout oxea 253-346 × 13-17μm.

Slender oxea 253-306 × 3- 6μm.

Micro oxea 86-130μm.

Distribution: Korea (Cheju-Island), North Atlantic, Celebes, Ceylon, Indian Ocean.

Family Renieridae Schmidt, 1870 레니에해면 과

35. **Gellius arcoferus* Vosmaer, 1885 판숨덩이해면 (Pl. 8, Figs. 1-2)

Gellius arcoferus Vosmaer, 1885 (p.29, pl.4, fig. 18, pl.5, figs. 87-90).

Gellius arcoferus: Fristedt, 1887 (pp. 438-439, pl.24, figs. 29-31, pl.28, fig. 16); Lundback, 1902 (pp.62-63, pl. 12, figs. 11a-c).

Material examined: Sögwip'o (fish net), 14/VII/1982.

Remarks: This sponge is flat from, size up to 4cm × 5cm × 0.3cm. Surface is uneven. Texture is hard, but fragile. Dermal membrane is thin. Pores are dispersed on the body and oscules are 0.1-0.2mm in diameter. Colour in alcohol is ivory.

Spicules: Oxea 411-450 × 14-23μm.
Slender oxea 350-370 × 4- 7μm.
Toxa 48-100μm.
Sigma 14-19μm.

Distribution: Korea (Cheju-Island), Ellesmere Island, Davis strait.

36. **Reniera ventilabrum* Fristedt, 1887 판레니에해면 (Pl. 8, Figs. 3-5)

Reniera ventilabrum Fristedt, 1887 (p. 420, pl.24, fig.3, pl.27, fig.8).

Material examined: Mosŭlp'o (fish net), 18/VII/1987.

Remarks: The shape is massive, size up to 10cm × 6cm × 2cm. Surface is smooth. Pores are dispersed on the body and oscules are 0.3mm-0.6cm in diameter. Texture is fragile. Colour in life is red purple, but in spirits is yellow or brown.

Spicules: Oxea 112-224 × 8-14μm.
Slender oxea 112-210 × 1- 3μm.

Distribution: Korea (Cheju-Island), the Atlantic Ocean.

37. **Reniera pigmentifera* Dendy, 1905 보라판레니에해면 (Pl. 9, Figs. 1-4)

Reniera pigmentifera Dendy, 1905 (p. 143, pl.4, fig. 10).

Material examined: Söngsanp'o, 30/VI/1984; Sögwip'o, 2/VIII/1984.

Remarks: This sponge is incrusting on the ston. Pores are dispersed on the body. Surface is a little hair with the protruding spicules. Oscules are 0.1-0.8mm in diameter. Texture is very soft and fragile. Skeleton is irregular isodictyla network. Colour in life is pale purple, but in spirits is ivory.

Spicules: Thick oxea 137-145 × 6-10μm.
Style 106-132 × 5- 6μm.
Strongyle 97-117 × 8- 9μm.

Distribution: Korea (Cheju-Island), South Africa.

Family Callyspongiidae De Laubenfels, 1936 예쁜이해면 과

38. *Callyspongia elegans* (Thiele, 1899) 예쁜이해면

Material examined: Mosŭlp'o, 19/VII/1987.

39. *Callyspongia confederata* (Ridley, 1884) 보라예쁜이해면

Material examined: Sögwip'o, 29/XII/1986.

40. *Callyspongia elongata* (Ridley & Dendy, 1886) 길쭉에쁜이해면

Material examined: Mosŭlp'o, 6/VII/1986, 18/VII/1987, 29/VI/1988; Sŏdo, 28/VII/1988.

Family Coelosphaeridae Hentschel, 1923 강해면 과

41. **Coelosphaera physa* (Schmidt, 1875) 거품강해면 (Pl. 10, Figs. 1-6)

Desmancido physa Schmidt, 1875 [cited from Lundback, 1910 (p. 11)].

Cornulum ascidioides Fristedt, 1887 (p. 495, pl. 25, figs. 1-2, pl. 29, fig. 21).

Histiderma physa: Thiele, 1903 (p. 385, pl. 21, figs. 16a-b); Lundback, 1910 (pp. 11-13, pl. 1, figs. 12-13, pl. 4, fig. 3).

Coelosphaera physa: Hentschel, 1929 [cited from De Laubenfels, 1936 (p. 71)].

Material examined: Anhŭng, 27/X/1984.

Remarks: Collected specimens are thin fragmented. Surface is uneven, and dermal membrane is present. Pores are dispersed on the body, but oscules are not observed. Texture is very soft and fragile. Colour in spirits dermal part is pale grey, and gastral part is dark grey.

Spicules: Tylote 221-400 × 4-10µm.

Isochela 18-54µm.

Raphid 60-100µm.

Distribution: Korea (Yellow Sea), the Atlantic Ocean.

ABSTRACT

The identified Ceratinomorpha consist of 41 species, 21 genera and 12 families. Among them, two species, *Clathria mosulpia* and *Haliclona ulreungia*, were new species and the following species were new to Korea: *Ophlitaspongia pennata californica* De Laubenfels, 1936, *Desmacella rosea* Fristedt, 1887, *Clathria dayi* Levi, 1963, *Clathria parva* Levi, 1963, *Axociella cylindrica* Hallman, 1920, *Axocielita calla* (De Laubenfels, 1934), *Myxilla sigmatifera* (Levi, 1963), *Haliclona perlucida* (Griessinger, 1971), *Petrosia nigricans* Lindgren, 1897, *Gellius arcoferus* Vosmaer, 1885, *Reniera ventillabrum* Fristedt, 1887, *Reniera pigmentifera* Dendy, 1905, and *Coelosphaera physa* (Schmidt, 1875).

REFERENCES

- Bakus, G.J., 1966. Marine poeciloscleridan sponges of the San Juan Archipelago, Washington. J. Zool. Lond. **149**: 415-531.
- Bakus, G.J. & K.D. Green, 1987. The distribution of marine sponges collected from the 1976-1978. Bureau of Land Management Southern California Bight Program. Bull. Southern California Acad. Sci. **86**(2): 57-88.
- Burton, M., 1959. Sponges. The John Murray Expedition 1933-34. Sci. Rep. **10**(5): 150-281.
- De Weerd, W.H. & R.W.N. Van Soest, 1986. Marine shallow-water Haplosclerida (Porifera) from the South Eastern part of the North Atlantic Ocean. Zool. Ver. **225**: 1-49.
- Dendy, A., 1905. Report on the sponges collected by Professor Herdman at Ceylon in 1902. Rep. Pearl Oyster Fish.

- Gulf of Manaar, 3, Suppl. **18**: 57-246.
- Dendy, A., 1921. Report on the sigmatotetraxonida collected by H.M.S. 'Sealark' in the Indian Ocean. Trans. Linn. Soc. London. **18**: 1-164.
- Fristedt, K., 1887. Sponges from the Atlantic and Arctic Oceans and the Behring Sea. Vega-Expeditionens Vetensk. Iakttagelser (Nordenskiöld), **4**: 401-471.
- Griessinger, T.M., 1971. Étude des Réniérides de Méditerranées (Demosponges, Haplosclérides). Bull. Mus. nation. Hist. Nat. Zool. **3**(3): 97-182.
- Hallman, E.F., 1920. New genera of monaxonid sponges related to the genus *Clathria*. Proc. Linn. Soc. New South Wales **44**: 769-792.
- Hentschel, E., 1912. Kiesel-und Hoynschwamme der Aru und Kei-Inseln: Abh. Senckenb. Naturf. Ges. **34**: 291-448.
- Hoshino, T., 1981. Shallow-water demosponges of Western Japan, I.J. Sci. Hiroshima Univ., Ser. B. Div. 1, **29**(1): 47-205.
- Laubenfels, M.W. De., 1932. The Marine and fresh-water sponges of California. Proc. U.S. Nat. Mus., No. 2927 **81**: 1-140.
- Laubenfels, M.W. De., 1934. New sponges from the Puerto Rican Deep. Smithsonian Miscellaneous collection **91**(17): 1-28.
- Laubenfels, M.W. De., 1936. A discussion of the sponges fauna of the Dry Tortugas in particular and the West Indies. With material for a revision of the families and orders of the Porifera. Pub. Carnegie Inst. 467 Pap. Tortugas Lab. **30**: 1-225.
- Laubenfels, M.W. De., 1954. The sponges of the West Central Pacific. Ore. St. Monogr. Zool. **7**: 1-306.
- Levi, C., 1963. Spongiaires Afrique Du Sud (1) Poecilosclerides Trans. roy. Soc. Afr., **37**: 1-72.
- Lundback, W., 1902. Porifera. Part I: Homorhaphidae und Heterorhaphidae. Cappenhagen Bianco Luno, The Danish Ingolf-Exped., **6**: 1-108.
- Lundback, W., 1910. Porifera. Part III: Desmacidonidae (Pars) Danish Ingolf-Exped., **6**: 1-124.
- Kim, H.S., B.J. Rho & C.J. Sim, 1968. Marine sponges in South Korea (1). Korean J. Zool. **11** (2): 1-11.
- Rho, B.J., H.S. Kim & C.J. Sim, 1969. Marine sponges in South Korea (2). J. Korean Res. Inst Better Living, Ewha Woman's Univ. **3**: 153-160.
- Rho, B.J. & C.J. Sim, 1972a. Marine sponges in South Korea(3). Ibid. **8**: 181-192.
- Rho, B.J. & C.J. Sim, 1972b. Faunal studies on the sponges in Korea. R-72-82. Min. Sci. Tech., 121-138.
- Rho, B.J. & C.J. Sim, 1976. On the classification and the distribution of the marine benthic animals in Korea. 4. Sponges. J. Korean Res. Inst. Better Living, Ewha Woman's Univ. **16**: 67-87.
- Rho, B.J. & K.H. Lee, 1976. A survey of marine sponge of Haeundae and Its Adjacent Waters. Ibid. **17**: 93-111.
- Rho, B.J. & C.J. Sim, 1979. A taxonomic study on the Korean sponges. 1. Poecilosclerina. Ibid. **23**: 61-67.
- Rho, B.J. & C.I. Yang, 1983. A systematic study on the marine sponges in Korea. 2. Ceractinomorpha. Ibid. **32**: 25-45.
- Sim, C.J., 1981. A systematic study on the marine sponges in Korea. 1. Ceractinomorpha & Tetractinomorpha. Soong Jun Univ. Eassays and Papers. **11**: 83-105.
- Sim, C.J., 1982. A systematic study on the marine sponges from Jeju Island. Soong Jun Univ. Eassays and Papers. **12**: 187-210.
- Sim, C.J., 1985. A systematic study on the marine sponges from the South Sea and the Yellow Sea of Korea. Korean J. Syst. Zool. **1**(1-2): 1-11.
- Sim, C.J. & J.H. Bae, 1987. On the classification of marine sponges of the Hongdo and its adjacent waters. J. Sci. Res. Inst. Han Nam Univ. **17**: 169-189.
- Sim, C.J. & M.H. Kim, 1988. A systematic study on the marine sponges in Korea. 7. Demospongia and Hexactinellida. Korean J. Syst. Zool. **4**(1): 21-42.
- Thiele, J., 1899. Studien über pazifische spongien. Zoologica. Heft. **24**: 1-33.

Thiele, J., 1903. Kieselschwämme von Ternate: II. Abhandl. Senckenb. naturf. Ges., **25**: 933-968.

Vosmaer, G.C.J., 1885. The sponges of the Willem Barents Expedition 1880 and 1881. Bijdr. Dierk., **12**: 1-47.

RECEIVED: 13 APRIL 1989

ACCEPTED: 4 MAY 1989

EXPLANATION OF PLATES 1-10

Plate 1

Figs. 1-4 *Ophlitaspongia pennata californiana* De Laubenfels.

1. entire animal.
2. surface of skeletal framework (SEM).
3. a seventy times enlarged skeletal framework (SEM).
4. megasclere : A, large subtylostyle; B, small subtylostyle; C-D, slender style.
Microsclere : E, large toxa; F-G, small toxa.

2. megasclere : A, thick style; B, slender style; C, acanthostyle.
microsclere : D, large spiny toxa; E, toxa; F, isochela.

Figs. 3-4 *Clathria parava* Levi.

3. entire animal.
4. megasclere : A, style; B, haed in spiny style; C, acanthostyle.
microsclere : D, spiny toxa; E, isochela.

Plate 2

Figs. 1-2 *Desmacella rosea* Fristedt.

1. entire animal.
2. megasclere : A, large tylostyle; B, middle tylostyle; C, small tylostyle.
Microsclere : D, large sigma; E, small sigma.

Plate 3

Figs. 1-5 *Clathria mosulpia* sp. nov.

1. entire animal.
2. surface of skeletal framework (SEM).
3. two hundred times enlarged skeletal framework (SEM).
4. gastral part of skeletal framework (SEM).
5. megasclere : A-B, subtylostyle; G, acanthostyle.
microsclere : C-D, large spiny toxa; E-F, toxa
H. isochela.

Plate 4

Figs. 1-2 *Clathria dayi* Levi.

1. entire animal.

Plate 5

Figs. 1-2 *Axociella cylindrica* (Ridley & Dendy).

1. Entire animal.
2. Megasclere : A, large trachy style; B-C, trachy style; D, slender style.
Microsclere : E, large isochela; F, small isochela.

Figs. 3-5 *Axociella calla* (De Laubenfels).

3. Entire animal: A, front; B, back.
4. Surface of skeletal framework (SEM).
5. Megasclere : A, subtylostyle; B, slender style; C, small subtylostyle.
Microsclere : D, large toxa; E, small toxa; F, isochela.

Plate 6

Figs. 1-2 *Myxilla sigmatifera* (Levi).

1. entire animal.
2. megasclere : A, trachy style; B, smooth style; C, tornote.

microsclere : D, large sigma; E, small sigma;
F, large isochela; G, small isochela.

2. megasclere : A, oxea; B, slender oxea.
microsclere : C, toxo; D, sigma.

Figs. 3-5 *Haliclona perlucida* (Griessinger).

Figs. 3-5 *Reniera ventilabrum* Fristedt.

3. entire animal.
4. surface of skeletal framework (isodictyal reticulum) (SEM).
5. megasclere : A, oxea; B, slender Oxea.

3. entire animal.
4. surface of skeletal framework (SEM).
5. megasclere : A, oxea; B, slender oxea.

Plate 7

Figs. 1-4 *Haliclona ulreungia* sp. nov.

1. entire animal.
2. dermal membrane (SEM).
3. surface of skeletal framework (SEM).
4. megasclere : A, oxea; B, slender oxea.

Figs. 5-6 *Petrosia nigricans* Lindgren.

5. entire animal.
6. megasclere : A, stout oxea; B, micro oxea.

Plate 8

Figs. 1-2 *Gellius arcoferus* Vosmaer.

1. entire animal.

Plate 9

Figs. 1-4 *Reniera pigmentifera* Dendy.

1. entire animal.
2. surface of skeletal framework (SEM).
3. three hundred times enlarged skeletal framework (SEM).
4. megasclere : A, oxea; B, style; C, strongyle.

Plate 10

Figs. 1-6 *Coelosphaera physa* (Schmidt).

1. entire animal.
2. surface of skeletal framework (SEM).
3. 4. tylote (SEM). 5. isochela (SEM). 6. raphid (SEM).

PLATE 1




PLATE 2




PLATE 3




PLATE 4













PLATE 5




PLATE 6




PLATE 7




PLATE 8











PLATE 9









PLATE 10

