

**Sabatieria pellucida** n. sp.

Fig. 151 a, b

Locality and material. — Falkland Islands, St. 42: 1♀

Dimensions:  $L = 2,0$  mm.,  $a = 28,57$ ,  $\beta = 10,15$ ,  $\gamma = 13,07$ ,  $V = 50$  %

Body in its whole length about evenly thick and tapering only in the oesophageal region gradually. Cuticle very finely annulated and structured in usual manner by smallest granules. Head rounded, provided with slender, submedian bristles, measuring at least  $2/3$  of the corresponding head diameter. Lateral organs typically spiral-shaped. Oesophagus of about even thickness throughout, provided posteriorly with a moderately large, strongly muscular bulb. Tail in its greatest part conical and only in its posterior part rather uniformly thinned, on the whole similar to the tail of *Parasabatieria mortenseni* DITL., described below. Thickness of the thinned part =  $2/7$  of the anal body diameter: end swollen and provided with a pair of hairs. Vulva situated exactly in the middle of the body. Female organs slightly developed.

St. 42. Falkland Islands. Port Louis.  $51^{\circ} 33' S$ . —  $58^{\circ} 9' W$ . Ooze and shells. 8 m. 26. 7. 1962.

Number of species found: 55; Number of specimens found: 372.




Fig. 151. *Sabatieria pellucida* n. sp. a. Anterior end,  $\times 600$ , b. Tail,  $\times 364$