

(in) Herpyllobiidae - Leuraeopoda

Aphanodomus

Wilson, 1924

A. terebellae (M)

Syn. Crypsidomus Levinsen, 1878, p. 375.
Aphanodomus Wilson, 1924, p. 15

Not Crypsidomus Günther, 1864, Reptile Genus.

Ἀφανὸς = hidden and δόμος = house.

Body of ♀ fastened to Terebella cirrata, semi-
lunar, not jointed, very soft, destitute of ru-
dimentary appendages of any sort, fastened
by its center to the larger intestine of
the host, the lateral portions, dimin-
ishing gradually in size, lying next
to the intestine. Egg sacs filiform, a
little longer than the width of the body,
fastened to a small conical process,
penetrate the skin of the host and
hang free above its back. At the
point where it is fastened to host is an aper-
ture leading into the parasite's intestine
4 mm. long; 16 mm. wide; 3 mm. thick.

XENOCOELOMIDAE - SIPHONOSTOMATOIDA

APHANODOMUS WILSON 1924

CARD 2

—, BRESCIANI, 1986: 127, 135, 139, 142, FIG. 8 C - tegument

Notodelphyoida.

Aphanodomus terebellae (Levinson), ♀.

Crypsidomus t- Levinson, 1877, p. 375: pl. 6. f. 19-20.

— — Stephensen 1913, p. 343: 1916, p. 305

Aphanodomus t- Wilson, 1920, p. 92.

--, Bresciani and Lutzen, 1975, p. 39.

Habitat. Terebella cirrata & Thelepus
cinctus (Vermes).

-- Belamane-Debattville & Laubier, 1960: 1
Greenland.

--, Bresciani & Lutzen, 1974, pp. ~~3-5~~, 6 25-53, figs.
1-42

-- BRUNEL 1965: pp.? - GULF OF ST. LAWRENCE

APHANODOMUS TEREBELLAE WILSON CARD 2

--, BRESCIANI, 1986: 126, 127, 135, 139, 141, 142-tegument;
FIG. 7 A-C

-- BRESCIANI & LÜTZEN 1961: 368

-- IZAWA 1987: 154-

-- RAIBAUT 1985: 266-