

Lichomolgidae - Cyclopoida,

Boeckia = synonym.

Brady, 1872,

B. arenicola ^(M).

Preoccupied by Malm for a genus of
crustacea in 1870; changed to
Hermannella by Canu in 1891.

Boeckia, Brady, 1872, p. 93.

Hermannella, Canu, 1891, p.

BOECKIA, KABATA 1988b: 6, ORIGIN OF NAME

Body elongated, compressed; head not distinct from thorax. Abdomen 5-seg'd in ♂, 3-seg'd in ♀. Right ♂ antenna geniculate. Exopod of second antennae 4-seg'd. Mds. large with sharp teeth and broad palp, latter biramose, 4- and 3-seg'd. MX. with strong marginal plumose setae. Mxpds. much elongated, terminal part 5-seg'd. 5 pairs of swimming legs, all biramose. Rami of first 4 pairs 3-seg'd, of fifth pair 3-seg'd. in ♀, but with a 2-seg'd. exopod in ♂. Ovisac single and ventral.

Boeckia = synonym.

Calanidae - Calanoida.

Thomson, 1882.

B. triarticulata ^(M)

Syn. Boeckia Thomson, 1882, p. 93.

~~No. 1~~ Boeckia Malm, Crustacea, 1870

~~No. 2~~ Boeckia Brady, 1872, p. 93.

Boeckella Guerne & Richard, 1889, p. 151.

Boeckella Giesbrecht & Schmeil, 1898, p. 60

BOECKIA, KABATA 1988b: 6, ORIGIN OF NAME

Body elongated, compressed; head not distinct from thorax. Abdomen 5-seg'd. in ♂, 3-seg'd. in ♀. Right male antenna geniculate. Exopod of second antennae 4-seg'd. Mds. large with sharp teeth and broad palp, latter biramose, 4- and 3-seg'd. Mx. with strong marginal plumose setae. Mxpds. much elongated, terminal part 5-seg'd. 5 pairs of swimming legs, all biramose. Rami of first 4 pairs 3-seg'd., of fifth pair 3-seg'd. in ♀, but with a 2-seg'd. exopod in ♂. Ovisac single and ventral.

Lichomolgidae - Cyclopoida,

Boeckia arenicola G.S. Brady, ♀.

— — Brady, 1872, p. 430;

Boeckia preoccupied, so changed to
Hermannella arenicola by Canu in 1891.