

in)

Harpacticoida - Cerniniidae

Eucanvella valid

T. Scott, 1901 -

E. spinifera (M).

Syn. Eucanvella T. Scott, 1901, p. 245.

Eucanvella G.O. Sars, 1903, p. 25.

Prefix eu and Cannella.

Head partly separated from 1^o seg't.: gen. seg't.
produced on either side - 1^o antennae 7-joint-
ed: Exopod of 2^o antennae 4-jointed - Both
rami of swimming legs 3-jointed -

Eucanuella T. Scott

- Brodskaya 1963: 1786, 1793, 1802
- Montagna 1981: 1218, in key to Cerviniidae.

Eucanuella langi Por, 1964

-- -- Por, 1964, pp. 66-68, pl. 5 figs. 34-37

-- Por 1962: 18, 45-46, FIGS. 34-37, TAB. 5, N.S.P.,
MEDITERRANEAN COAST OF ISRAEL (THESIS)

Eucanuella longirostrata Ito, 1983

--, Ito, 1983, p. 177-184, figs. 17-20.

EUCANUELLA RETICULATA JOYER, 1969

-- COULL, 1973 c: 208; FIG. 27; TAB. 1

(E. AFF. R., OFF NORTH CAROLINA, USA) - ♂ ONLY

-- APOSTOLOV + PETKOVSKI 1980: 24-25.

Cerviniidae-Harpacticoida.

Eucanvella spinigera T. Scott, ♂, ♀.

— T. Scott, 1901, p. 245: pl. 18, fig. 1-10,

— T. Scott, 1902. b, p. 456

— G. O. Sars, 1904, p. 25: pl. 14, ♀.

— G. O. Sars, 1911, p. 352: suppl. pl. 2, fig. 1, ♂.

—, LANG, 1948, p. 184, Pl. 110: 2.

Habitat. British Seas: Norway Coast.

—, Jespersen, 1940, p. 72. Iceland.

—, COULL, 1973 c: 208 ~~1973 d: 208~~

E. spinifera, Wells, 1965. p. 2
--, Jakobi, 1959, p. 139, 279
--, Klie, 1941, p. 5
- Por 1962: 19, 45-46