

(in)
Homoiotēs, valid Catigoidae Caligidae.
Wilson, 1905. siphonostomatoida
H. palliata.

Syn. Homoiotēs Wilson, 1905, p. 661

- Dojiri, 1983: 370-371

- PRABHA 1983: 54, SYNONYM OF LEPEIOPHTHEIRUS

ὁμοιότης, similarity or likeness.

Carapace large, shield-shaped. No lunules.
Mtd. with saw-teeth on inner margin only.
Second mx. small & divided as in Leptophtheirus.
1st & 4th Thorax legs uniramose, 2nd. & 3rd. pairs
biramose. Gen. segt covered by a pair of
fused dorsal plates, which extend back over
base of abdomen in ♀, with a projection ending
in a spine on either side of abdomen.
5th. & 6th. legs plainly visible in ♂. & 5th pair much
better developed than usual in this family.
Abdomen unsegmented, without plates or processes;
anal laminae small, flattened, and armed
with plumose setae.

Homoiotes bermudensis Heegaard

_____, Heegaard, 1943, p. 21, figs. 56-64, ♀ ♂, Host:
Thynnus pelamys. Bermuda

A synonym of *Dentigryps* *curtus* Wilson - cf.
Heegard, 1945 (*Archiv für Zoologi* 35A, no. 18) p. 15.

Homoiotes bermudensis Heegaard (1943)

--, Kabata, 1973, p. 1893 (= Dentitgryps curtus)

Homoiotes palliata, gen. + sp. n. ^{Caligidae.} Wilson, ♂, ♀

—, Wilson, 1905, p. 661: pl. 29.

--, Kabata, 1973, p. 1892-1893 (= Lepeophtheirus
pravipes, L. breviventris & L. bifidus)

-- PRABHA 1983: 54

Hosts. Blue Cod (Ophiodon elongatus).

Locality unknown.