

in *Ternaeopodaida-Anneacheridae*
Ternaeascus, valid.
Claus, 1886, T. nematoxys.

Syn. Ternaeascus Claus, 1886, p. 231.

Λέρνη and ασκὸς = a bladder or bag.

Male. Well segmented, thickset, with broad, scale-like dorsal projections. Fourth & fifth abdominal segments fused inter se. Mouth parts with stylet-formed mandibles, without the second mxp'ds.

Female. Nematode-like, with a double row of dorsal & ventral scale-like skin-projections. The claw-antennae separated by quite a distance. Mouth-parts with an epipharyngeal beak and cheliform mxp'ds in a deeply, excavated pit. The 3 pairs of thorax legs very tiny rudiments.

(Claus. 1887.)

PHILICHTHYIDAE - POECILOSTOMATOIDEA

(2)

Lernaeascus

- Izawa, 1973, p. 84
- ESSAFI & RAIBAUT 1980: 357
- CASTRO ROMERO 1994a: 13-

Poecilostomatoida Philichthyidae.

Lernaeascus nematoxys, Claus, ♀.

—, Claus, 1886. p. 231.

—, —, 1887^a, p. 281; pl. XIV, figs. 1-12.
pl. XV, figs. 13-24; pl. XVI, figs. 25-32.

—, Claus, 1887^b, p. 241.

—, Delamare Deboutteville 1962. p. 43, fig. 17

Host. Solea monochir.

L. NEMATOXIS, KAZACHENKO 1994: 42, FIG. 9B

European Seas.

Philichthyidae: Proctostomidae

CARD 2

Lernaglascus nematoxys Claus, 1886

---, Kabata, 1970, fig. 64, 65, p. 76

-- IZAWA 1987: 153-