

in)

Pennellidae: Poecilostomatida ~~caligorda~~.

Peroderma, valid.

~~Dermatidae~~

(n)

Heller, 1865, P. cylindricum.

Syn. Peroderma Heller, 1865, p. 250.  
≡ Taphrobia Cornalia, 1875, p. 197

Τηρός = disabled or maimed, and Σέπα = skin.  
Τείρω = to pierce

Head globular, covered with numbers of closely packed fine-branching processes; united to long, sac-like gen. seg't. by a short, narrow slender neck, at a r. a., to v surface gen. seg't, and  $\frac{1}{3}$  v distance from its extremity.

First antennae 3-jointed, setiferous.

First 2-pairs of thorax legs triramous, others uniramous, each with 2-joints.

Egg-sacs long; eggs uniserial.

- 361 61 x 1

Peroderma

Kabata, 1972, p. 321, key to genera of lernaeoceridae  
Kabata 1979: 290 - 91, ... in Pennellidae

Peraderma bellottii, Richardson, ♀.  
Lernaeidae.

— —, Richardson, 1882 a, p. 149.

— —, Carus, 1885, p. 373.

Peraderma b — , B. Trinck. 1903, p. 109.

Host. Muscles of laryngeal apparatus in Sarcoptes  
benoiti.



(over)  
Mediterranean.

To the Milanese C. Bellotti who pro-  
cured the specimens for Richiardi.

Referred to the new genus Cardiodectes,

Lernaeidae.

Peroderma bruchiata, B.-Smitt. ♀.

—, B.-Smitt, 1898a, p. 13; pl. VII, fig. 2.

-- BEN HASSINE, RAIBAUT ET AL. 1990: 10

a synonym of Peroderma cylindricum.

Host, Buried in flesh of Nucrates ductor.

Hemaeidae.

Peroderm a cylindricum, Heller. ♀.

- —, Heller, 1865, p. 250; pl. XXV, fig. 6.  
— —, Richardsoni, 1876, p. 189; pl. VI, fig. 1.  
— brauchiata, B. Smith, 1898b, p. 13; pl. VII, fig. 2.  
— cylindricum, Brian, 1898, p. 20.

Host, Buried in side muscles of Coila dus-  
sumieri.



Mediterranean & Bombay.

- , Carus, 1885, p. 373.
- , B. Smith, 1899, p. 488.
- , Brian, 1908, p. 16
- , Brian, 1912, p. 31: pl. 4, figs. 5-6: pl. 8, figs. 5-6
- Taphrobia pilchardi, Cornalia, 1874, p. 197: pl. 6.
- P. cylindricum Giard, 1888, p. 929.
- , Wilson, 1917, a, p. 50.
- , Brian, 1924, p. 45.
- , Nunes-Ruivo, 1953. pp. 6-8, figs. 4. Sardina  
pilchardus. Mediterranean.
- , Nunes-Ruivo, 1956, p. 30- 31. Sardinella cameronensis Regan (?)  
+ S. eba C. & R. Angola.
- , Rodriguez-Roda, 1959, pp. 125-129. Sardina pilchardus, Barbate, Spain,  
Larache, Morocco.

Peroderma cylindricum Heller

(2)

--- --, Nunes Ruivo, 1962. pp. 77-78

--- --, Rodriguez-Roda, 1959. pp. 125-129

--- --, Ho Ju-- Shey 1966, p. 192.

---, Kabata, 1970, p. 35, 97-100, figs. 98-99, 109, 116

--- , Argilias, A., 1930, pp. 96

---, IZAWA, 1977: 253, 256, 258

--- CHANGEURX 1960: 82

--- EL GHARBI ET AL. 1983(85): 191

--- RAIBAUT ET AL. 1971: 189-190, HOST RECORDS, TUNISIA

--- HONMA + HO 1988: 41-

PERODERMA CYLINDRICUM HELLER

(3)

-- BEN HASSINE, RAIBAUT, BEN SOUSSI + ROUSSET

1990: 10-15, FIGS. 1-2.

PERODERMA PACIFICA (IZAWA, 1977)

-- IZAWA, 1977: 253-258 ♀; FIGS. 1-13.

EX BREGMACEROS JAPONICUS (FISH)

Lernaeidae. — Lernaeoida.

Peraderma petersii, Trichiardi ♀

— — Trichiardi, 1881, a, p. 247. nomen nudum.

— — Trichiardi, 1881, c, p. 387.

Peraderma petersi B-Smith, 1903, p. 109.

Host. Gobius buccatus, near pectoral fin.

— — —

Mediterranean