

in Canthocamptidae - Harpacticoida.

Pteropsyllus, valid -

(M).

T. Scott, 1906. a.

P. consimilis.

Syn. Pteropsyllus T. Scott, 1906,

Pteropsyllus G. O. Sars, 1911, p. 412.

Tetragoniceps T. Scott, pars, not Brady.

Πτέρον, a wing and ψυλλός, a glea.

General form like *Phyllopoedopsyllus*; gen. segt.
in ♀ large, depressed; caudal rami narrow lin-
ear. First antennae 8-jointed, 1st. joint much
elongated. Exopod of 2nd. pair rudimentary.
Endopod of 1st legs much longer than exopod,
3-jointed, of following legs 2-jointed.
Last legs in ♀ very large, forming beneath the
urosome a roomy incubatory chamber.

TETRAGONICIPITIDAE

PTEROPSYLLUS T. SCOTT

(2)

- Hays 1992: 71-
- Kunz 1984c: 33-

Canthocamptidae - Harpacticoida.

Pteropsyllus consimilis T. Scott, ♂.♀.

Tetragoniceps — T. Scott, 1894, p. 244: pl. 7, f. 4-12.

Pteropsyllus — T. Scott, 1906, a, p. 459: pl. 11, f. 7-8.

— — T. Scott, 1906, b, p. 343.

— — Sars, 1911, p. 413: Suppl. pl. 38.

— —, LANG, 1948, p. 592.

Habitat. British Seas: Norway Coast.

— —, Wells, 1970, p. 264

— —, Jakobi, 1959, p. 145

— Wells, 1967: 143

-- , Klie, 1950, p. 102

-- , COLL, 1973a: 11-12 ; SYN: P. PLEBIUS PLEBIUS;

P. P. FURCATUS

-- MIELKE 1989b: 116

Pteropsyllus furcatus

--, Jakobi, 1959, p. 280

--, Klie, 1950, p. 102

Pteropsyllus plebeius Monard

____ Monard 1935a, p. 62, fig. 164-171 ♀, Roscoff,
France

— —, HANT, 1946, p. 892.

--, Jakobi, 1959, p. 145, 280

--, Kli, 1950, p. 102

-- PLEBIUS, COULL, 1973a: 11-12 = P. CONSIMILIS.

-- MIELKE 1989b: 116

PTEROPSYLLUS

PLEBEIUS

FURCATUS

KUNZ

—, LANG, 1948, P. 893.

---, = P. CONSIMILIS, COULL, 1973a: 11-12

---, MIELKE 1989b: 116

HAB. GER. HELGOLAND.

Pteropsyllus trisetosus Mielke, 1989

-- Mielke 1989b: 109-116, Figs. 8-12, n. sp. female +
male, Galapagos

Pteropsyllus sp. Wells, 1961

-- Mielke, 1989b: 116