

in Arietellidae

Centropagidae

Rhapidophorus.

C.L. Edwards, 1891.

R. wilsoni. ^(M)

Syn. Rhapidophorus Edwards, 1891, p. 84.
Rhapidophorus Giesb. & Schmeil, '98, p. 125

ραπίς = whip-lash.

long and stout: carapace half, the
abdomen a third as long as middle body.
The chitini covering of the 5th thorax seg't.
extends back over the 10 abdomen seg't. and
wholly covers it. 1st antennae 22-jointed,
about as long as the carapace = exopod of 2.
antennae 3-jointed, endopod 6-jointed.
No mandibular palp: posterior nupds. 7-jointed
5th legs stunted: each made up of 2 basal joints,
a 2-jointed exopod and a spine-like one-
jointed endopod. Both rami of 10 to 40
legs 3-jointed - spines on last joint of
exopod 3 lateral & 1 terminal.

RHAPIDOPHORUS EDWARDS, 1891

2

- CAMPANER 1984a: 107
- MAUCHLINE 1988: 706
- OHTSUKA ET AL. 1991: 801
- OHTSUKA ET AL. 1994: 1317

Rhapidophorus wilsoni Edwards, 1891

card 2

Foss ^{hagen} & Go.
-- Ohtsuka, et al. 1991:801

Calanidae - Calanoidae
Rhapidophorus wilsoni Edwards, ♀.

- — C.L. Edwards, 1891, p. 84: pl. 4, fig. 1-11.
— — Giesbrecht, 1892, p. 423.
— — Giesbrecht & Schmeil, 1898, p. 125.

uncertain genus.

- — Fosshagen A., 1968 p. 61

Habitat. Bahama Islands.

-- CAMPANER, 1977: 817

-- CAMPANER 1984a: 106, 107, FIG. 3.

-- REID 1990 ♀: 185