

POECILOSTOMATOIDA.

Trichthacerus.

Krøyer, 1863 -

Chondracanthidae
Ternaeopodoidea

T. peristedii.

Syn. Trichthacerus, Krøyer, 1863, p. 269.
= Trichthacerus, Heller, 1865, p. 233.

TYPE: T. PERISTEDII KRØYER, 1863

Trixia, in 3 parts: Répas, a horn.

Female. Body stout, unsegmented; separated from a short & dilated head by a distinct neck -
Abdomen well defined, 2-jointed -

1st antennæ 3-jointed, setiferous: 2nd. pair horn-like; 3-parted, without terminal claws.

Gen. seg't. stout, saccate, without processes -

4 pairs rudimentary thorax legs, stout, sub-chelate -

1st. pair biramous, others more rudimentary -

Egg-sacs large, stout; eggs multiserial.

Male a pigmy like that of Chondrocanthus.

- Oakley 1930: 185, rejected genus
- Yamaguti 1963: 296, restored genus
- Ho 1970: 115, rejected genus; synonymized with
possibly Blias,

Trichthacerus molestus Heller - see Blias

Trichthacerus peristedii, Kröyer, ♂. ♀,
Chondracanthidae.

— —, Kröyer, 1863, p. 264; pl. XIV, fig. 7, a-f.
— —, B-Smith, 1899, p. 491.

-- Ho 1970: 115; POSSIBLY A SPECIES OF BLIAS

Host, gills of Peristethus.



Rio Janeiro.