

Taxonomy on Marine Sponges from Geojedo Island, Korea

Chung Ja Sim and Kyung Jin Lee

(Department of Biology, Han Man University, Taejeon 300-791, Republic of Korea)

ABSTRACT

The sponge specimens were collected from Geojedo Island, Korea during from 1994 to 1998 with fishing nets or by scuba divers. They were identified into 34 species of 25 genera in 15 families, of which the one species, *Tedania rhoi* is a new species, and *Clathria (Axociella) simae* is new to the Korean fauna. This species were redescribed.

Key words: taxonomy, marine sponges, Geojedo Island, Korea

INTRODUCTION

Sponges from Geojedo Island, Korea are so far poorly known. To date, only six species have been reported by Rho and Sim (1972, 1981) and Sim (1981). From 1994 through 1998, sponges were collected from Geojedo and its adjacent water by scuba divers and with fishing nets. 34 sponge species of 25 genera in 15 families were found from this Island, of which the one new species, *Tedania rhoi* was discovered. The species *Clathria (Axociella) simae* (Hooper, 1996) was redescribed.

Sponges were identified on the basis of overall shape, coloration and texture of the sponge, and on spicule shape and size. For skeletal arrangement, thin free-hand sections were made with a surgical blade from specimens hardened in alcohol. Spicules were examined by dissolving a piece of sponge in sodium hypochlorite (bleach). For the morphological analysis of microsclere, spicules were prepared using an AKASHI critical pointed drier, then viewed with an ISI-SS40 SEM at Hannam University. SEM analysis of spicules followed the procedure of Rützler (1978).

This study was supported by grant from Hannam University, 1997-1998.

SYSTEMATIC ACCOUNT

Phylum Porifera Grant, 1836 해면동물 문
 Class Demospongiae Sollas, 1885 보통해면 강
 Order Homosclerophorida Dendy, 1905 동골해면 목
 Family Plakinidae Schulze, 1880 관해면 과

1. *Plakortis simplex* Schulze, 1880 일삼해면

Previous records. Geojedo (Rho and Sim, 1981).

Distribution. Korea (Korea Strait), Japan (Hiwasa, Tokushima).

Order Astrophorida Levi, 1973 별해면 목
 Family Geodiidae Gray, 1867 조디아해면 과

2. *Caminus awashimensis* Tanita, 1969 카미너스해면

Material examined. Ssanggeun (scuba diver), 8 July 1996.

Distribution. Korea, Japan.

3. *Geodia reniformis* Thiele, 1898 앞조디아해면

Material examined. Daepo (fishing-net), 2 Oct. 1996.

Distribution. Korea (Cheju Island), Japan (Sagami Bay, Enoshima).

Order Hadromerida Topsent, 1894 경해면 목
 Family Clionidae Gray, 1867 호박해면 과

4. *Cliona celata* Grant, 1826 호박해면

Previous records. Geojedo (Rho and Sim, 1972).

Material examined. Ssanggeun (scuba diver), 8 July 1996.

Distribution. South Korea, Gulf of St. Lawrence to South Carolina, Gulf Coast of Louisiana and Texas, Pacific Coast of North America.

Family Suberitidae, Schmidt, 1870 코르크해면 과

5. *Suberites ficus* Johnston, 1842 집게코르크해면

Material examined. Daepo (fishing-net), 2 Oct. 1994; Ssanggeun (scuba diver), 8 July 1996.

Distribution. Korea (Korea Strait and East Sea), Japan (Seto Island Sea).

6. *Suberites japonicus* Thiele, 1898 왜코르크해면

Previous records. Geojedo (Sim, 1981).

Material examined. Ssanggeun (scuba diver), 8 July 1996; Heungnam (scuba diver), 30 Jan. 1997.

Distribution. Korea (Geojedo), Japan (Seto Island Sea).

7. *Terpios fugax* Dachassing and Michelotti, 1864 피터오즈해면

Material examined. Ssanggeun (scuba diver), 8 July 1996.

Distribution. Korea (Cheju Island), West Central Pacific (Marshall Island, Ponape Island), Caribbean Sea (Caragao Island, Carcas Bay), Bermuda (Bermudas).

Order Poecilosclerida Topsent, 1928 다골해면 목
 Family Microcionidae Carter, 1875 작은눈해면 과

8. *Clathria (Clathria) mosulpia* Sim and Byeon, 1989 모슬침유령해면

Material examined. Daepo (fishing-net), 3 Oct. 1994.

Distribution. Korea (Cheju Island).

9. *Clathria (Clathria) spinispicula* Tanita, 1968 침유령해면

Material examined. Daepo (fishing-net), 2 Oct. 1994, 3 Oct. 1994.

Distribution. Korea (Korea Strait, Cheju Island), Japan.

10. *Clathria (Axociella) simae* (Hooper, 1996) 등근축털해면 (Fig. 1 A-G)

Clathria (Microcionia) simae Hooper, 1996, p. 224

Axociella cylindrica Sim and Byeon, 1989, p. 39, Pl. 5, figs. 1-2

Material examined. Maemuldo (fishing-net), 9 July 1996.

Description. Sponge erect and dichotomous flattened branching, 17.5 cm high, 10 cm wide, 1.5 cm stem diameter, surface hispid with spicules. Oscules rare 0.4-0.8 mm in diameter. Texture firm, colour beige in spirit.

Ectosome: Erect spicule brushes.

Choanosome: Fibroreticulation fibres cored by one to two styles, thin subtylostyle found interstitially.

Spicules:	Megascleres, Thick styles	300-750 × 15-30 μm
	Subtylostyles	260-650 × 3-7 μm
	Microscleres, Toxas (large)	100-130 × 2-4 μm
	(medium)	70-90 × 1-2 μm
	(small)	25-40 × 0.5-1 μm
	Isochelas (large)	18-25 μm
	(small)	10-12 μm

Remarks: This species was described as a new species, *Clathria (Microcionia) simae* by Hooper (1996). However, subgenus *Microcionia* and *Axociella* have quite different skeletal architecture. *Axociella* displays well differentiated axial and extra-axial components of the spicules. Such axial differentiation is not found in *Microcionia*. This sponge has fibroreticulate cored thick style spicules. Echinating acanthostyles are absent. Thick styles have microspined or smooth bases. Thin tylostyles also have microspined bases. Accordingly, this *Clathria (Microcionia) simae* is reassigned to the

Table 1. Spicule dimensions of *Clathria (Axociella) simae*.

Localities	Spicules	Thick styles (μm)	Subtylostyles (μm)	Toxas (μm)			Isochelas (μm)	
				large	medium	small	large	small
Moseulpo (Cheju Island)				100-120	60-90	25-45	16-27	9-12
Fishing-net, 1987		290-730 × 15-30	265-610 × 3-8	× 2-4	× 1-2	× 1		
Maemuldo (Geojedo)				100-130	70-90	25-40	18-25	10-12
Fishing-net, 1996		300-750 × 15-30	260-650 × 3-7	× 2-4	× 1-2	× 0.5-1		




Fig. 1. *Clathria* (*Axociella*) *simae*. A, Side view; B, Styles: a, thick style, b, subtylostyle; C, Skeletal structure (SEM); D, Skeletal structure; E, Megascleres; a-b, head of thick style, c-d, head of subtylostyle (SEM); F-G, Microscleres; F, a, side view of large isochela, b, large isochela, c, small isochela (SEM); G, a, large toxa, b, medium toxa, c, small toxa (SEM). Scale bars = 2 cm (A), 100 μ m (B), 30 μ m (C), 200 μ m (D), 50 μ m (E-G).

subgenus *Axociella*. Hooper (1996) erroneously recorded the type locality of this sponge as the South China Sea. The actual type locality of this species was Cheju Island, in the South Sea of Korea. Table 1 shows that the spicule dimensions of these species is very similar in both localities.

Distribution. Korea (Korea Strait, Cheju Island), Port Jackson.

11. *Lissoplocamia tocushima* Tanita, 1970 미끈이해면

Material examined. Daepo (fishing-net), 2 Oct. 1994; 9 July 1996

Distribution. Korea (South Sea), Japan.

12. *Ophlitaspongia noto* Tanita, 1963 바늘뼈해면

Previous records. Geojedo (Rho and Sim, 1981).

Material examined. Gudo (scuba diver), 6 Feb. 1996.

Distribution. Korea (Korea Strait), Japan (Aikawa, Seto Island, Noto-peninsula).

Family Raspailiidae Hentschel, 1923 털해면 과

13. *Raspailia folium* Thiele, 1898 잎사귀해면

Material examined. Daepo (fishing-net), 2 Oct. 1994.

Distribution. Korea (Korea Strait, Cheju Island), Japan (Enoshima).

14. *Raspailia hirsuta* Thiele, 1898 털많은가지해면

Material examined. Daepo (fishing-net), 2 Oct. 1994; 9 July 1996.

Distribution. Korea (Korea Strait, Cheju Island), Japan (Sagami Bay).

15. *Ceratopsis ramosa* Thiele, 1898 가지뿔해면

Material examined. Daepo (fishing-net), 9 July 1996.

Distribution. Korea (Cheju Island), Japan (Sagami Bay).

Family Coelosphaeridae Hentschel, 1923 강해면 과

16. *Lissodendoryx firma* Lambe, 1895 뇌산호끈적해면

Material examined. Daepo (fishing-net), 3 Oct. 1994.

Distribution. Korea (Cheju Island), California, San Juan Archipelago, Washington, Western Canada.

17. *Lissodendoryx isodictyalis* Carter, 1882 두드럭끈적해면

Previous records. Geojedo (Rho and Sim, 1972).

Distribution. Korea (Korea Strait), North America, Caribbean Sea.

Family Myxillidae Hentschel, 1923 끈적해면 과

18. *Myxilla incrustans* Johnston, 1942 껍질끈적해면

Material examined. Daepo (fishing-net), 2 July 1994, 9 July 1996, Heungnam (scuba diver), 30 Jan. 1997, 13 Jan. 1998.

Distribution. Korea (Korea Strait, East Sea, South sea), Japan (Ariake Sea).

19. *Myxilla productus* Hoshino, 1981 긴끈적해면

Material examined. Daepo (fishing-net), 9 July 1996.

Distribution. Korea, Japan.

20. *Myxilla setoensis* Tanita, 1961 넓적끈적해면

Material examined. Daepo (fishing-net), 2 Oct. 1994.

Distribution. Korea (Korea Strait, East Sea), Japan (Seto Island Sea).

Family Tedaniidae Ridley and Dendy, 1886 테다니해면 과

21. *Tedania rhoi* n. sp. 로이테다니해면 (신칭) (Fig. 2 A-E)

Material examined. Holotype: Por. 25 (NHM, Hannam Univ.), Gudo (scuba diver 5 m), 6 Feb. 1996. Paratype: Por. 25-1, Por. 25-2, Por. 25-3 (Dept. of Biology, Hannam Univ.); Por. 25-4, Por. 25-5 (NHM, Ewha Womans Univ.), Ssanggeun (scuba diver 5 m), 8 July 1996.

Description. Commonly massive, sometimes cushion shaped measuring up to 75 × 77 × 20 mm; surface uneven bearing very low papillae. Smooth to the touch, sometimes very slimy. Rare oscules 0.5 mm in diameter. Texture hard and compressible. Live specimen appear fleshy-pink, creamy-beige in spirit.

Ectosome: Thin dermal membrane, 1 mm thick, smooth with a skin like membrane. Tylotes arranged at an angle to the surface, mixed with many onychaetes.

Choanosome: A series of ascending tracts of styles, connected by much spongin, tylota occur in small amounts.

- Spicules: Megascleres, Styles270-340 × 8-10 μm
- Tylotes230-290 × 4-5 μm
- Microscleres, Long thin onychaetes160-170 × 1 μm
- Short thin onychaetes.....40-50 × 1 μm
- Thick onychaetes120 × 3 μm

Ectymology. This species is named for Dr. Boon Jo Rho, who has contributed greatly to sponge taxonomy since 1966.

Remarks. This new species is similar in spicule morphology to *Tedania diversirhaphideophora* Brønsted, 1923. This species was redescribed by Bergquist (1961) and by Bergquist and Fromont (1988). *T. diversirhaphideophora* has two size categories of onychaetes, whereas this new species usually has three size categories of onychaetes. No other species of *Tedania* has been described that has three categories of onychaetes. We found 18 specimens from two localities (Gudo and Ssanggeun), all have papillae on the surface of the sponges. The ends of the tylotes are smooth, without acanthose. Table 2 shows the spicule dimension of this new species.

Table 2. Spicule dimensions of *Tedania rhoi* n. sp.

Localities	Spicules	Styles (μm)	Tylotes (μm)	Onychaetes (μm)		
				large thin	Small thin	Thick
Gudo						
SCUBA, 5m	270-340 × 8-10	230-290 × 4-5	160-170 × 1	40-50 × 1	120 × 3	
6 Feb. 1996						
Ssanggeun						
SCUBA, 5m	280-320 × 8-10	220-280 × 4-5	160-170 × 1	40-50 × 1	120-130 × 3	
8 July 1996						




Fig. 2. *Tedania rhoi* n. sp. A, Side view; B, Megascleres: a, style, b, tylote; C, Skeletal structure; D, a, head of style, b, head of tylote; E, Microscleres: a, base of thick onychaete, b, point of thick onychaete, c, base of long thin onychaete, d, point of long thin onychaete, e, point of short thin onychaete (SEM). Scale bars = 1 cm (A), 100 μm (B), 150 μm (C), 200 μm (D), 5 μm (E).

Family Mycalidae Lundbeck, 1905 깃해면 과

22. *Mycale plumosa* Carter, 1882 깃바늘뻘해면

Material examined. Ssanggeun (scuba diver), 8 July 1996; Hakdong (scuba diver), 29 Jan. 1997.

Distribution. Korea (Korea Strait), Japan (Matsushima Bay), Indian Ocean.

23. *Oxymycale rhoi* Sim and Lee, 1998 로이간상바늘뻘해면

Material examined. Daepo (fishing-net), 3 Oct. 1994, 9 July 1996.

Distribution. Korea (Geojedo).

24. *Oxymycale koreana* Sim, 1982 한국간상바늘뻘해면

Material examined. Daepo (fishing-net), 3 Oct. 1994.

Distribution. Korea (Cheju Island).

25. *Esperiopsis uncigera* Topsent, 1928 관발톱해면

Material examined. Daepo (fishing-net), 9 July 1996.

Distribution. Korea (Korea Strait, East Sea), Japan (Sagami Bay).

Order Halichondrida Vosmaer, 1885 해변해면 목

Family Dictyonellidae van Soest, Diaz and Pomponi, 1990 수정해면 과

26. *Dactylella hilgendorfi* Thiele, 1898 수정해면

Material examined. Daepo (fishing-net), 9 July 1996.

Distribution. Korea (Korea Strait, East Sea), Japan (Seto Island Sea).

Family Axinellidae Ridley and Dendy, 1888 축해면 과

27. *Acanthella vulgata* Thiele, 1899 보통가시해면

Material examined. Daepo (fishing-net), 2 Oct. 1994.

Distribution. Korea (Korea Strait), Japan (Sagami Bay).

28. *Axinella copiosa* Thiele, 1898 축해면

Material examined. Daepo (fishing-net), 9 July 1996.

Distribution. Korea (Cheju Island), Japan (Hakodate).

29. *Phakellia elegans* Thiele, 1898 맵시해면

Material examined. Daepo (fishing-net), 2 Oct. 1994.

Distribution. Korea (Korea Strait), Japan (Sagami Bay).

Family Halichondriidae Vosmaer, 1885 해변해면 과

30. *Halichondria oshoro* Tanita, 1961 황록해면해면

Previous records. Geojedo (Rho and Sim, 1972).

Distribution. Korea (Korea Strait), Bristol Bay.

31. *Halichondria panicea* Pallas, 1776 회색해면해면

Material examined. Gudo (scuba diver), 6 Feb. 1996.

Distribution. Korea (Korea Strait, East Sea), Japan (Matsushima, Oginohama Bays), Gulf of Mexico.

32. *Hymeniacion sinapium* De Laubenfels, 1930 주황해면해면

Material examined. Gudo (scuba diver), 6 Feb. 1996; Gabae, 8 July 1996; Haegeumgang, 4 Sep. 1997; Haksan (scuba diver), 14 Jan. 1998.

Distribution. Korea (Korea Strait, East Sea, West Sea), Japan.

Order Haplosclerida Topsent, 1928 단골해면 목

Family Callyspongiidae De Laubenfelds, 1936 예쁜이해면 과

33. *Ceraochalina differentiata* Dendy, 1921 뽕예쁜이해면

Material examined. Gabae, 6 Feb. 1996.

Distribution. Korea (Korea Strait), Japan (Funakawa, Akita Prefecture, Noto-peninsula).

Family Chalinidae Gray, 1867 고삐해면 과

34. *Gellius angulatus* Bowerbank, 1866 솜덩이해면

Material examined. Daepo (fishing-net), 2 Oct. 1994, 3 Oct. 1994.

Distribution. Korea (Cheju Island), Guernsey, Rev. A.M. Norman Bay of Naples.

ACKNOWLEDGEMENT

The authors are specially grateful to Dr. James T. Wetzel, Department of Biology, Presbyterian College, for review of this manuscript.

REFERENCES

- Bergquist, P.R. 1961. The Demospongiae (Porifera) of the Chatham Islands and Chatham Rise, collected by the Chatham Islands 1954 Expedition. N. Z. Department of Scientific and Industrial Research Bulletin No. **139**(5): 169-206, figs 1-6 (N.Z. Oceanographic Institute Memoir 13).
- Bergquist, P.R. and P.J. Fromont. 1988. The Marine Fauna of New Zealand: Porifera, Demospongiae, Part 4 (Poecilosclerida). New Zealand Oceanographic Institute Memoir 96. pp. 1-197.
- Birsted, H.V. 1923. Sponges from the Auckland and Campbell islands. Papers from Dr Th. Mortensen's Pacific Expedition 1914-16. XV. Videnskabelige Beddelelser from Dansk Naturhistorisk Forening i Kjobenhavn **75**: 117-167.
- Hooper, J.N.A. 1996. Memoirs of the Queensland Museum. Queensland Museum. **40**: 1-626.
- Rho, B.J. and C.J. Sim. 1972. Marine sponges in South Korea (3). J. Korean Research Institute for Better Living Ewha Womans University. **8**: 181-192.
- Rho, B.J. and Sim, C.J. 1981. A taxonomic study on the marine sponges in Korea 3. Choristida. J. Korean Res. Inst. Liv'. Ewha Womans Univ. **28**: 55-65.
- Rützler, K. 1978. Sponges in coral reefs. In: Stoddart DR and Johannes RE (eds), Coral Reefs: Research Methods. Monogr Oceanogr Neth UNESCO **5**: 299-313.
- Sim, C.J. 1981. A taxonomic study on the marine sponges in Korea 2. Hadromerida. Korean J. Zool. **24**(1): 9-17.
- Sim, C.J. and H.S. Byeon. 1989. A systematic study on the marine sponges in Korea 9. Ceractinomorpha. Korean J. Syst. Zool. **5**(1): 33-57.

RECEIVED: 18 August 1998

ACCEPTED: 9 September 1998

거제도 해산 해면류의 분류학적 연구

심 정 자·이 경 진
(한남대학교 이과대학 생물학과)

요 약

1994년부터 1998년까지 거제도에서 어망과 잠수부에 의해서 채집된 해면류는 15과 25속 34종으로 동정, 분류되었다. 이 중 *Tedania rhoi* 로이테다니해면은 신종으로 밝혀졌고, *Clathria (Axociella) simae* 동근축털해면은 한국 미기록종으로 판명되었다.