

Anticoma tenuis ALLGÉN

Fig. 1 a, b

ALLGÉN 1930, p. 249—250.

Localities and material. Coast of Argentina, St. 2: 1♂, 1♀; Falkland Islands, St. 58: 1♀, Graham Land, St. 6: 1♀

Dimensions: ♂ L = 2,655 mm., $a = 64,76$, $\beta = 5,31$, $\gamma = 17,13$

♀ L = 3,063 mm., $a = 65,17$, $\beta = 6,13$, $\gamma = 14,38$, V = 54,29 %

From the above-named localities some specimens of an *Anticomid* were recorded, which in their whole shape agree rather well with this small, southern species, described first by the present author (1930) from the Staten Island in the Fuegian Archipelago.

It is easily recognizable by its *elongated, thin, about evenly thick body*, its rounded head and its filiform tail, similar to that of *A. similis* COBB.





Fig. 1. *Anticoma tenuis* ALLGÉN a. Anterior end, $\times 450$, b. Tail and spicular app., $\times 270$

- St. 2. Off the Coast of North Argentina. $37^{\circ} 15' S.$ — $56^{\circ} 8' W.$ Sand-mixed gravel. 100 m. 23. 12. 1901.
Number of species found: 12; Number of specimens found: 26.
- St. 6. Graham Region. $64^{\circ} 36' S.$ — $57^{\circ} 42' W.$ Stones and gravel. Mud-sample. 125 m. 20. 1. 1902.
Number of species found: 29; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland. $52^{\circ} 29' S.$ — $60^{\circ} 36' W.$ Sand and gravel. 197 m. 11. 9. 1902.
Number of species found: 23; Number of specimens found: 93.