

DITLEVSEN 1921, p. 27—29, pl. II, fig. 10, pl. III, fig. 4—7.

Localities and material. — Fuegian Archipelago, St. 62: 2♂♂, 3 juv.; St. 64: 1♂, 1♀, 14 juv.; Falkland Islands, St. 39: 2 juv.; St. 40: 1♀, 2 juv.; St. 49: 1 juv.; St. 51: 1 juv.; St. 53: 1 juv.; St. 55: 2♀♀, 4 juv.; St. 58: 1♀, 13 juv.; St. 59: 3♀♀. South Georgia, St. 22c: 1♀; St. 34: 2♂♂

Dimensions: ♂ L = 8,200 mm., $\alpha = 53,59$, $\beta = 16,4$, $\gamma = 117,14$
 ♀ L = 10,750 mm., $\alpha = 55,33$, $\beta = 6,32$, $\gamma = 137,82$, V = 61,4 %

Among the numerous *Thoracostomes* in the Antarctic material there were found also some males, which seem to agree so well in their general shape with *Th. aucklandiae*, described by DITLEVSEN (1921) from the Auckland Islands, that I should like to regard them as synonymous with that southern species.

Body on the whole evenly thick. Cuticle moderately thick. Head truncate, provided at the middle of the cephalic mail with 2 lateral and 8 submedial very small bristles in usual arrangement. Cephalic mail relatively weakly developed. In the expanded posterior parts of the anchor-shaped lobules there are no locules, contrary to *Th. antarcticum*.

Lateral organs small and rounded situated at the level of the posterior part of the mail. Ocelles situated 0,180 mm behind the front end, brownish coloured, with a lens.

Tail short and clumsy, shorter than the anal body diameter, at the ventral side weakly curved, at the end rounded.

Spicules also short, clumsy, weakly bent, swollen at the middle, uniformly thinned in the anterior third, rounded at the proximal end.

Accessory pieces immediately behind the spicules and measuring about half their length.

The very small supplementary organ is about one tail-length in front of the anus. It consists of 2 plates, directed one against the other, provided with anteriorly and posteriorly directed toothed branches. Vulva behind the middle of the body.

Female organs paired, elongated, with reflexed ovaries. On both sides of the vulva 2 elongated shell eggs.

Geographical distribution. — Auckland Islands (DITLEVSEN 1921), Australia: Sydney, the harbour, on *Pennaria*, and Port Jackson, among stones and algae (ALLGÉN 1951), Antarctica: Discovery Bay (ALLGÉN 1929).

- St. 22a. South Georgia. Cumberland, May-Bay. Catching over stony bottom among algae in and under the tide zone. 5. 5. 1902.
Number of species found: 11; Number of specimens found 63.
- St. 22b. South Georgia. Grytviken. 22. 5. 1902 and 20 m. depth. 11. 6. 1902.
Number of species found: 16; Number of specimens found: 77.
- St. 22c. South Georgia. Grytviken, from old kelp-rhizoids. 23. 5. 1902.
Number of species found: 45; Number of specimens found: 350.
- St. 22d. South Georgia. Grytviken. Sample of fine washings from old kelp. 22. 5. 1902.
Number of species found: 22; Number of specimens found: 200.
- St. 23. South Georgia. Off the mouth of the Moraine-Bay. 54° 23' S. — 36° 26' W. Grey clay with gravel and stones. 64—74 m. Bottom temp. + 1,65 C. 16. 5. 1902.
Number of species found: 32; Number of specimens found: 147.
- St. 23a. South Georgia. Moraine-Fiord. 148 m. Bottom temp. — 0,35 C. 15. 2. 1902.
Number of species found: 14; Number of specimens found: 51.
- St. 23b. South Georgia. Moraine-Fiord. 14 m.
Number of species found: 12; Number of specimens found: 49.
- St. 24. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 37' W. Grey clay. 95 m. 20. 5. 1902.
Number of species found: 23; Number of specimens found: 120.
- St. 25. South Georgia. Off the "Kochtopf"-Bay 54° 22' S. — 36° 27' W. Grey clay with some algae. 24—52 m. 21. 5. 1902.
Number of species found: 29; Number of specimens found: 83.
- St. 26. South Georgia. Off the "Kochtopf"-Bay. 54° 22' S. — 36° 27' W. Stony bottom with algae off the Macrocystis-Formation. 30 m. 24. 5. 1902.
Number of species found: 11; Number of specimens found: 29.
- St. 28. South Georgia. Mouth of the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Sand and algae. 12—15 m. 24. 5. 1902.
Number of species found: 58; Number of specimens found: 338.
- St. 30. South Georgia. The Moraine-Fiord. 54° 24' S. — 36° 26' W. Clay with sparse stones. 125 m. Bottom temp. — 0,25 C. 26. 5. 1902.
Number of species found: 23; Number of specimens found: 247.
- St. 33. South Georgia, in the "Kochtopf"-Bay. 54° 22' S. — 36° 28' W. Clay and algae. 22 m. 30. 5. 1902.
Number of species found: 23; Number of specimens found: 106.
- St. 34. South Georgia. Off the mouth of the Cumberland-Bay. 54° 11' S. — 36° 18' W. Grey clay with a few stones. 250—310 m. Bottom temp. + 1,45 C. 5. 6. 1902.
Number of species found: 38; Number of specimens found: 224.
- St. 39. Falkland Islands. Port William. 51° 40' S. — 57° 41' W. Sand and small stones with algae. 40 m. 4. 7. 1902.
Number of species found: 11; Number of specimens found: 12.
- St. 40. Falkland Islands. Berkeley Sound. 51° 33' S. — 58° 0' W. Gravel and shells with algae. 16 m. Bottom temp. — 2,75 C. 19. 7. 1902.
Number of species found: 54; Number of specimens found: 291.
- St. 41. Falkland Islands. Port Louis, shallow water. 51° 33' S. — 58° 9' W.
Number of species found: 51; Number of specimens found: 310.
- St. 42. Falkland Islands. Port Louis. 51° 33' S. — 58° 9' W. Ooze and shells. 8 m. 26. 7. 1902.
Number of species found: 55; Number of specimens found: 372.
- St. 42a. Falkland Islands. Port Louis: Greenpatch. Material shaken up from algae and kelp-rhizoids, cast up on shore by storm. 30. 7. 1902.
Number of species found: 54; Number of specimens found: 150.
- St. 46. Falkland Islands. Port Louis. Carenage Creek. 51° 32' S. — 58° 7' W. Sandy bottom with quantities of *Codium*. 1 m. 9. 8. 1902.
Number of species found: 28; Number of specimens found: 103.
- St. 47. Falkland Islands. Port Louis. Mouth of the Carenage Creek. 51° 32' S. — 58° 7' W. Shells and stones. 3—4 m. 9. 8. 1902.
Number of species found: 63; Number of specimens found 247.

- St. 49. Falkland Islands. Berkeley Sound. 51° 35' S. — 57° 56' W. Shells and stones. 25—30 m. 10. 8. 1902.
Number of species found: 27; Number of specimens found: 58.
- St. 51. Falkland Islands. Port William. 51° 40' S. — 57° 42' W. Sand. 22 m. 3. 9. 1902.
Number of species: 45; Number of specimens found: 245.
- St. 53. Falkland Islands. Port William. 51° 40' S. — 57° 47' W. Sand and gravel. 12 m. 3. 9. 1902.
Number of species found: 65; Number of specimens found: 372.
- St. 54. Falkland Islands. Stanley Harbour. 51° 42' S. — 57° 50' W. Ooze with shells. 10 m. 3. 9. 1902.
Number of species found: 2; Number of specimens found: 7.
- St. 55. Falkland Islands. Port Albemarle. 52° 11' S. — 60° 26' W. Sandy bottom with algae. 40 m. 8. 9. 1902.
Number of species found: 33; Number of specimens found: 113.
- St. 56. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 9' S. — 60° 33' W. Sandy bottom with algae. 15 m. 8. 9. 1902.
Number of species found: 15; Number of specimens found: 40.
- St. 57. Falkland Islands. Port Albemarle. Albemarle Harbour. 52° 8' S. — 60° 33' W. Sand. 18—30 m. 11. 9. 1902.
Number of species found: 21; Number of specimens found: 40.
- St. 58. Falkland Islands. S. W. West Falkland. 52° 29' S. — 60° 36' W. Sand and gravel. 197 m. 11. 9. 1902.
Number of species found: 23; Number of specimens found: 93.
- St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank. 53° 45' S. — 61° 10' W.
Crushed shells with stones 137—150 m. 12. 9. 1902.
Number of species found: 20; Number of specimens found: 70.
- St. 62. Fuegian Archipelago. Beagle-Channel. 54° 53' S. — 67° 56' W. Sand-mixed clay. 140 m. 16. 9. 1902.
Number of species found: 12; Number of specimens found: 63.
- St. 64. Fuegian Archipelago. North side of the Beagle Channel between Ushuaia and Lapataia. 54° 52' S. — 68° 25' W. Shells and algae. 35 m. 13. 10. 1902.
Number of species found: 33; Number of specimens found: 192.
- St. 67. Fuegian Archipelago. Ushuaia. 54° 49' S. — 68° 18' W. Ooze. 6 m. 16. 10. 1902.
Number of species found: 9; Number of specimens found: 42.