

Enoplolaimus propinquus DE MAN

DE MAN 1922, p. 132 and 1922 a, p. 256—257, fig. 47 a—b.

Localities and material. — Fuegian Archipelago, St. 3: 1 juv.; Falkland Islands, St. 15: 3♀♀, 3 juv.

The first find of this species was described from the Zuider Sea. It differs from the above mentioned, elder closely related North-Sea species *E. vulgaris*, except in the tail and the spicular apparatus, in the more strongly developed cephalic bristles, being much more elongated than in the species of comparison.

Geographical distribution. — Sweden: Southern Kattegatt (ALLGÉN 1947), the Sound (ALLGÉN 1929, 1935), Holland: Zuider Sea (DE MAN 1922, SCHUURMANS-STEKHOFEN 1930), Belgium: (DE CONINCK and SCHUURMANS-STEKHOFEN 1933).

St. 3. Fuegian Archipelago. 54° 43' S. — 64° 8' W. Rubble stones and gravel. 36 m. 6. 1. 1902.
Number of species found: 20; Number of specimens found: 37.

St. 15. Falkland Islands. Port William. 51° 40' S. — 57° 49' W. Macrocystis-Formation. 10 m. 31. 3. 1902.
Number of species found: 44; Number of specimens: 150.