

***Halicylindrolaimus conicaudatus* n. sp.**

Fig. 240 a, b

Localities and material. — Falkland Islands, St. 49: 1♀; South Georgia, St. 28b: 3♂♂, 2 juv.

Dimensions: ♂ L = 3,317 mm.,  $\alpha = 26,85$ ,  $\beta = 7,06$ ,  $\gamma = 19,30$

Body moderately slender. Cuticle thin and smooth. Head truncate to rounded. Cephalic bristles not observed. Buccal cavity bordered by thin walls, broad, cylindrical. Lateral organs situated about  $10\ \mu$  behind the front end, rounded, large, measuring in section  $6\ \mu$ , i. e.  $2/5$  of the corresponding head diameter. Oesophagus short, probably bulb-like swollen posteriorly. Tail short, conical, posteriorly rounded, measuring  $1,6 \times$  the anal body diameter. Spicules short, strongly curved, thickest in their proximal part, posteriorly pointed. Accessory piece slender, its apophysis directed dorsally and posteriorly.

St. 23b. South Georgia. Moraine-Fiord. 14 m.

Number of species found: 12; Number of specimens found: 49.

St. 49. Falkland Islands. Berkeley Sound.  $51^{\circ} 35' S.$  —  $57^{\circ} 56' W.$  Shells and stones. 25—30 m. 10. 8. 190

Number of species found: 27; Number of specimens found: 58.




Fig. 240. *Halicylindrolaimus obtusicaudatus* n. sp. a. Anterior end,  $\times 450$ , b. Tail and spicules,  $\times 350$

