

**Oncholaimus rotundicaudatus** n. sp.

Fig. 53 a, b

Locality and material. — Falkland Islands, St. 40: 1♀

In one of the samples from St. 40 was found a single *female*, so strongly wounded, that its total length was not to be measured.

Body in its whole length evenly thick. Head truncate to rounded. Buccal cavity bordered by strongly chitinized walls, cylindrical, about twice deeper than broad; length  $37\ \mu$  and maximum thickness  $19\ \mu$ . Dorsal tooth  $20\ \mu$  from the anterior edge, i. e. immediately behind the middle of the buccal cavity. Subventral teeth about evenly large. Their points in a distance of  $13\text{--}15\ \mu$  from the anterior edge of the buccal cavity. Oesophagus elongated and about evenly thick. Ventral gland opening far anteriorly; its duct swollen to an ampulla before opening. The gland proper was not to be found. Lateral organs at the middle of the buccal cavity, thinwalled, circular. *Tail short, evidently rounded.*

St. 40. Falkland Islands. Berkeley Sound.  $51^{\circ} 33' S$ . —  $58^{\circ} 0' W$ . Gravel and shells with algae. 16 m. Bottom temp. —  $2,75\ C$ . 19. 7. 1902.



53. *Oncholaimus rotundicaudatus* n. sp. a. Anterior end, b. Tail,  $\times 450$