

Terschellingia baylisi n. sp.

Fig. 238 a, b

Locality and material. — Falkland Islands, St. 53: 1♂

Dimensions: L = 1,808 mm., $\alpha = 30,19$, $\beta = 11,74$, $\gamma = 13,55$

Body about evenly thick throughout. Cuticle finely annulated. Head rounded, with 4 rather short, slender submedial bristles.

Buccal cavity small, rather difficult to study, probably typical to this genus. Lateral organs 34μ behind (i. e. close to) the front end, thin-walled, circular, in section $6,7 \mu$, i. e. exactly 0,16 of the corresponding head diameter. Oesophagus very short and thick with a very large, about rounded, strongly muscular bulb. Tail short, cylindro-conical, weakly curved, posteriorly rounded. Spicules short, slender, strongly curved, at their proximal end expanded, obliquely-truncate, evenly thick, posteriorly pointed. Accessory piece partly in the shape of an elongated arch-like formation, at the distal part of the spicules, partly as an apophysis with a dorsal and posterior direction.

St. 53. Falkland Islands. Port William. $51^{\circ} 40' S.$ — $57^{\circ} 47' W.$ Sand and gravel. 12 m. 3. 9. 1902.

Number of species found: 65; Number of specimens found: 372.

